

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International Advanced Subsidiary and Advanced Level

## **MARK SCHEME for the March 2016 series**

### **9701 CHEMISTRY**

**9701/33**

Paper 3 (Advanced Practical Skills), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the March 2016 series for most Cambridge IGCSE<sup>®</sup> and Cambridge International A and AS Level components.

© IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – March 2016	9701	33

question	indicative material	mark	total																					
1 (a)	I All thermometer readings <b>and</b> mass of <b>FA 2</b> recorded. Do not award if mass of <b>FA 2</b> > 0.50 g.	1	[4]																					
	II All temperatures recorded to 0.5 °C.	1																						
	Award III and IV if within ranges given of supervisor's value.	2																						
	<table border="1"> <thead> <tr> <th>supervisor's <math>\Delta T / ^\circ\text{C}</math></th> <th>III</th> <th>IV</th> </tr> </thead> <tbody> <tr> <td><math>\geq 46.0</math></td> <td><math>\pm 5.0</math></td> <td><math>\pm 2.5</math></td> </tr> <tr> <td>36.0–45.5</td> <td><math>\pm 4.0</math></td> <td><math>\pm 2.0</math></td> </tr> <tr> <td>26.0–35.5</td> <td><math>\pm 3.0</math></td> <td><math>\pm 1.5</math></td> </tr> <tr> <td>16.0–25.5</td> <td><math>\pm 2.0</math></td> <td><math>\pm 1.0</math></td> </tr> <tr> <td>6.0–15.5</td> <td><math>\pm 1.0</math></td> <td><math>\pm 0.5</math></td> </tr> <tr> <td>&lt; 6.0</td> <td><math>\pm 0.5</math></td> <td>–</td> </tr> </tbody> </table>	supervisor's $\Delta T / ^\circ\text{C}$	III	IV	$\geq 46.0$	$\pm 5.0$	$\pm 2.5$	36.0–45.5	$\pm 4.0$	$\pm 2.0$	26.0–35.5	$\pm 3.0$	$\pm 1.5$	16.0–25.5	$\pm 2.0$	$\pm 1.0$	6.0–15.5	$\pm 1.0$	$\pm 0.5$	< 6.0	$\pm 0.5$	–		
	supervisor's $\Delta T / ^\circ\text{C}$	III	IV																					
	$\geq 46.0$	$\pm 5.0$	$\pm 2.5$																					
	36.0–45.5	$\pm 4.0$	$\pm 2.0$																					
	26.0–35.5	$\pm 3.0$	$\pm 1.5$																					
16.0–25.5	$\pm 2.0$	$\pm 1.0$																						
6.0–15.5	$\pm 1.0$	$\pm 0.5$																						
< 6.0	$\pm 0.5$	–																						
(b)	I Axes labelled with units and uniform scale chosen to use more than half of each axis including 10 °C above the highest recorded temperature.	1	[4]																					
	II All recorded points plotted (minimum 9).	1																						
	III Appropriate lines of best fit drawn: <ul style="list-style-type: none"> <li>best fit lines must be or a smooth curve;</li> <li>Points not on the line must be balanced on either side of the best-fit line and any points ringed or labelled as anomalous ignored.</li> </ul>	1																						
	IV Lines extrapolated and correct value (within 0.5 °C) of $\Delta T$ read from graph.	1																						
(c) (i)	Correctly calculates $Q = 25 \times 4.2 \times \Delta T$ from (b).	1	[3]																					
(ii)	Correct expression for value of enthalpy change $= \frac{-(c)(i) \times 24.3}{\text{mass in (a)} \times 1000}$ (ignore sign)	1																						
	Negative sign <b>and</b> both answers recorded to 2–4 sig. fig. <b>and</b> no rounding to 1 sig. fig. during calculation (unless exact value).	1																						
(d)	Incorrect, as the acid was in excess already.	1	[1]																					
(e)	Any one from: <ul style="list-style-type: none"> <li>use lid or use specified extra insulation to reduce heat losses (by convection or conduction);</li> <li>use a pipette or burette for <b>FA 1</b> to reduce % error/as more accurately calibrated (owtte);</li> <li>use magnesium turnings/powder so reaction complete sooner as there is heat loss while magnesium ribbon is still reacting;</li> <li>use lid or plastic cup with higher walls to reduce acid spray;</li> </ul>	1	[1]																					

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – March 2016	9701	33

question	indicative material	mark	total
2 (a)	<p><b>I</b> Initial and final burette readings and volume added recorded for rough titre <b>and</b> accurate titre details tabulated.</p> <p><b>II</b> Initial and final burette readings recorded and volume of <b>FA 4</b> added recorded for each accurate titration. All headings and units correct for accurate titrations:</p> <ul style="list-style-type: none"> <li>initial/final (burette) reading/volume <b>or</b> reading/volume at start/finish</li> <li>volume/<b>FA 4</b> added/used <b>or</b> titre</li> <li>(cm<sup>3</sup>) <b>or</b> /cm<sup>3</sup> <b>or</b> in cm<sup>3</sup> <b>or</b> cm<sup>3</sup> by every entry.</li> </ul> <p><b>III</b> All accurate burette readings are recorded to the nearest 0.05 cm<sup>3</sup>.</p> <p><b>IV</b> Has two uncorrected, accurate titres within 0.1 cm<sup>3</sup>.</p> <p><b>V, VI and VII</b> Award <b>V, VI and VII</b> for <math>\delta \leq 0.20 \text{ cm}^3</math> Award <b>V and VI</b> for <math>0.20 \text{ cm}^3 &lt; \delta \leq 0.30 \text{ cm}^3</math> Award <b>V</b> for <math>0.30 \text{ cm}^3 &lt; \delta \leq 0.50 \text{ cm}^3</math></p>	1  1  1  1	[7]
(b)	<p>Mean titre correctly calculated from clearly selected values.</p> <ul style="list-style-type: none"> <li>Candidates must average two (or more) titres where the <b>total</b> spread is <math>\leq 0.20 \text{ cm}^3</math>.</li> <li>Working must be shown or ticks must be put next to the two (or more) accurate readings selected.</li> <li>The mean should normally be quoted to 2 d.p. rounded to the nearest 0.01.</li> </ul> <p><i>Note: the candidate's mean will sometimes be marked as correct even if it is different from the mean calculated by the examiner for the purpose of assessing accuracy.</i></p>	1	[1]
(c)(i)(ii)	Correctly calculates $\frac{0.100 \times (\mathbf{b})}{1000}$ <b>and</b> <b>(ii) = (i)</b>	1	[5]
(iii)	Correct expression $\frac{(\mathbf{c})(\mathbf{ii}) \times 1000 \times 10}{25}$	1	
(iv)	mol Mg = mass in <b>1(a)</b> / 24.3 <b>and</b> mol HCl = <b>(c)(iii)</b> $\times 25 / 1000$	1	
	mol HCl > 2 $\times$ mol Mg (owtte) so the statement is correct. Allow ecf from incorrect <b>(iii)</b> .	1	
	Final answers <b>(i), (ii)</b> and <b>(iii)</b> to 3 or 4 sig. fig. <b>and</b> no rounding errors.	1	

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – March 2016	9701	33

question	indicative material	mark	total
(d)	Correct expression $\frac{0.1 \times 100}{(b)}$ and answer to minimum 2 sig. fig. / correct answer to minimum 2 sig.fig. <b>and</b> <b>FA 3</b> (is measured more accurately). Allow ecf from <b>(b)</b> > 41.67 cm <sup>3</sup> then <b>FA 4</b> (is measured more accurately).	1	[1]

test	observations	
	FA 5	FA 6
NaOH	no reaction / no change / no ppt	white ppt, soluble in excess
NH <sub>3</sub>	no reaction / no change / no ppt	white ppt, insoluble in excess
HCl (warm)	blue solution brown gas / gas turning brown / gas turns blue litmus red / bleaches	no reaction / no change
H <sup>+</sup> / MnO <sub>4</sub> <sup>-</sup>	decolourises / purple to colourless <b>or</b> (solution) stays colourless	stays purple / pink <b>or</b> changes to purple / pink
Ba <sup>2+</sup> / HCl	no reaction / no change / no ppt	white ppt, insoluble in HCl

question	indicative material	mark	total
<b>FA 5 is NaNO<sub>2</sub>; FA 6 is Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>; FA 7 is Na<sub>2</sub>SO<sub>3</sub> (Na<sub>2</sub>S<sub>2</sub>O<sub>5</sub>)</b>			
3 (a)	Observations fully correct for both <b>FA 5</b> and <b>FA6</b> for NaOH.	1	[8]
	Observations fully correct for both <b>FA 5</b> and <b>FA6</b> for NH <sub>3</sub> .	1	
	Observation of blue solution or brown gas with <b>FA 5</b> <b>and</b> no reaction with <b>FA 6</b> for HCl.	1	
	Observations fully correct for both <b>FA 5</b> and <b>FA6</b> for H <sup>+</sup> / MnO <sub>4</sub> <sup>-</sup> .	1	
	Observations fully correct for both <b>FA 5</b> and <b>FA6</b> for Ba <sup>2+</sup> / HCl.	1	
	Cations: <b>FA 5</b> unknown <b>and</b> <b>FA 6</b> Al <sup>3+</sup> / aluminium	1	
	Anions: <b>FA 5</b> NO <sub>2</sub> <sup>-</sup> / nitrite <b>FA 6</b> SO <sub>4</sub> <sup>2-</sup> / sulfate	1	

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – March 2016	9701	33

question	indicative material	mark	total
(b) (i)	(Warm with) Al and NaOH and test gas with (damp) red litmus paper.	1	[5]
	No reaction <b>and</b> not nitrate / N / same element as <b>FA 5</b> .	1	
(ii)	BaCl <sub>2</sub> / Ba(NO <sub>3</sub> ) <sub>2</sub> and HCl / HNO <sub>3</sub> <b>or</b> H <sup>+</sup> / KMnO <sub>4</sub> / acidified potassium manganate(VII) <b>or</b> any named acid, (warm) and test gas with H <sup>+</sup> / KMnO <sub>4</sub> .	1	
	Ba <sup>2+</sup> and acid: white ppt, soluble in acid <b>or</b> H <sup>+</sup> / MnO <sub>4</sub> <sup>-</sup> : solution decolourises / purple to colourless <b>or</b> acid and test gas with H <sup>+</sup> / KMnO <sub>4</sub> : gas (evolved with acid) which decolourises H <sup>+</sup> / MnO <sub>4</sub> <sup>-</sup> (paper).	1	
	<b>FA 7</b> contains sulfite / SO <sub>3</sub> <sup>2-</sup>	1	