



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

CHEMISTRY

9701/22

Paper 2 AS Level Structured Questions

February/March 2026

1 hour 15 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **16** pages. Any blank pages are indicated.

1 The Group 2 elements Mg to Ba show trends in their physical properties down the group.

(a) Table 1.1 shows some information about the elements Mg to Ba.

Table 1.1

element	proton number	shorthand electronic configuration	sum of first and second ionisation energies / kJ mol^{-1}	third ionisation energy / kJ mol^{-1}
Mg	12	[Ne] $3s^2$	2186	7740
Ca	20	[Ar] $4s^2$	1740	4940
Sr	38	[Kr] $5s^2$	1608	4120
Ba	56	[Xe] $6s^2$	1468	3390

(i) Give the full electronic configuration of a calcium atom.

..... [1]

(ii) Explain the trend shown in the sum of the first and second ionisation energies, as proton number increases.

.....

 [2]

(iii) Write an equation to represent the third ionisation energy of barium.

..... [1]

(iv) Explain why the third ionisation energy of magnesium is much greater than the sum of its first and second ionisation energies.

.....

 [2]



(b) When strontium reacts with water, an alkaline solution is produced.

(i) Write an equation for this reaction. Include state symbols.

..... [2]

(ii) Samples of the alkaline solution produced in **1(b)(i)** are tested with dilute HCl(aq) and separately with dilute $\text{H}_2\text{SO}_4\text{(aq)}$.

State what you would observe in each test.

observation with dilute HCl(aq)

.....

observation with dilute $\text{H}_2\text{SO}_4\text{(aq)}$

..... [2]

(c) The nitrates and carbonates of Group 2 elements decompose on heating.

(i) State the trend in the thermal stability of the Group 2 carbonates down the group.

..... [1]

(ii) Write an equation for the thermal decomposition of $\text{Ca(NO}_3)_2$.

..... [1]

(d) Calcium chlorate(I), Ca(ClO)_2 , is used for water purification in the same way as sodium chlorate(I), NaClO .

Ca(ClO)_2 can be prepared by the reaction of aqueous Ca(OH)_2 with Cl_2 .

(i) Write an ionic equation for the reaction of cold dilute OH^- ions with Cl_2 .

..... [1]

(ii) State why NaClO and Ca(ClO)_2 are used in water purification.

..... [1]

[Total: 14]



- 2 Nitrogen monoxide, NO, is an atmospheric pollutant.

(a) Figure 2.1 shows the dot-and-cross diagram of NO.




Figure 2.1

- (i) NO has an unpaired electron.

State the term for a molecule with an unpaired electron.

..... [1]

- (ii) In NO, the N and O atoms are bonded by a sigma (σ) and a pi (π) bond.

Complete Figure 2.2 to show the pi bond in NO that is formed from orbital overlap.





Figure 2.2

[1]

- (b) In the laboratory, NO is made in the reaction of copper metal with dilute nitric acid. The reaction is kept cold to prevent the oxidation of NO.



A student reacts 0.00393 mol Cu with excess dilute $\text{HNO}_3(\text{aq})$ at 5°C and 101 kPa.

Calculate the maximum volume, in cm^3 , of $\text{NO}(\text{g})$ that is produced under these conditions.

volume of $\text{NO}(\text{g})$ produced = ... cm^3 [4]



- (c) (i) State a natural process that creates atmospheric NO.


..... [1]

- (ii) Photochemical smog forms when NO reacts with unburned hydrocarbons.

Identify the major component of photochemical smog.

..... [1]


- (iii) NO has a catalytic role in the oxidation of atmospheric SO₂.



Show, using relevant equations, the catalytic role that NO has in this oxidation.

..... [2]

- (d) Reaction 1 occurs during the catalytic removal of NO from car exhaust gases.



The activation energy, E_a , of this reaction is 570 kJ mol^{-1} .

- (i) Complete Figure 2.3 to sketch the energy profile for reaction 1.

Show the enthalpy change, ΔH , and the activation energy, E_a .




Figure 2.3

[2]

- (ii) On your diagram in Figure 2.3, show how the addition of a catalyst affects reaction 1. [1]

[Total: 13]

- 3 Secondary halogenoalkanes are important building blocks in organic chemistry.

2-chlorobutane is an example of a secondary halogenoalkane.

2-chlorobutane




Figure 3.1

- (a) Explain why 2-chlorobutane is classified as a secondary halogenoalkane.

.....
 [1]

- (b) 2-chlorobutane can be formed by the electrophilic addition of HCl to different isomers of C_4H_8 . Compound **A** is one of these isomers.






Figure 3.2

- (i) Draw a structural isomer of **A** that can also form 2-chlorobutane when it reacts with HCl .



[1]

- (ii) Draw and give the systematic name of the geometrical isomer of **A**.



systematic name [2]




- (iii) State and explain the origin of geometrical isomerism in alkenes.

.....

 [2]

- (c) Butane, C_4H_{10} , reacts with Cl_2 in the presence of ultraviolet light to form 2-chlorobutane, which has molecular formula C_4H_9Cl .




- (i) Name the mechanism of this reaction.

..... [1]

- (ii) Complete the reaction scheme given to show the formation of 2-chlorobutane.

You may use molecular formulas in your answer.

initiation



propagation



termination [2]

- (d) The rates of hydrolysis of three secondary halogenoalkanes, **B**, **C** and **D**, are compared.




Figure 3.3

Each halogenoalkane is dissolved in ethanol and reacted with $AgNO_3(aq)$.

Identify the order of the rates of hydrolysis of compounds **B**, **C** and **D**. Explain your answer.



..... > >
 fastest slowest

explanation
 [2]

(e) The organic product of hydrolysis of each of **B**, **C** and **D** is butan-2-ol.

Complete Table 3.1 to give details of some typical chemical reactions of butan-2-ol.

Table 3.1

reagent and conditions	observation with butan-2-ol	organic product(s)
acidified $\text{K}_2\text{Cr}_2\text{O}_7$		
Na(s)	effervescence	
	yellow precipitate	 and CHI_3

[3]

[Total: 14]





Turn over



4 Cyanoethanoic acid, $\text{CH}_2(\text{CN})\text{COOH}$, is a weak organic acid.

(a) At room temperature, $\text{CH}_2(\text{CN})\text{COOH}$ undergoes an acid–base reaction with $\text{NaOH}(\text{aq})$.

(i) Define weak acid.

.....
 [1]

(ii) Sketch the pH titration curve on Figure 4.1 for the reaction of:

- 25.00 cm^3 of 0.10 mol dm^{-3} $\text{CH}_2(\text{CN})\text{COOH}(\text{aq})$ and
- 0.10 mol dm^{-3} $\text{NaOH}(\text{aq})$, added to excess.




Figure 4.1

[3]

(iii) Write the overall ionic equation for the neutralisation of $\text{CH}_2(\text{CN})\text{COOH}(\text{aq})$ by $\text{NaOH}(\text{aq})$ in 4(a)(ii).

Include state symbols.

..... [1]

(iv) Suggest the formula of the organic product when $\text{CH}_2(\text{CN})\text{COOH}(\text{aq})$ is heated under reflux with $\text{NaOH}(\text{aq})$.

..... [1]



- (b) $\text{CH}_2(\text{CN})\text{COOH}$ is very soluble in water because it can form hydrogen bonds with water.

Complete Figure 4.2 to show **one** hydrogen bond between a molecule of $\text{CH}_2(\text{CN})\text{COOH}$ and a molecule of H_2O .

Label the hydrogen bond clearly, and show any relevant partial charges and lone pairs of electrons.




Figure 4.2

[3]

- (c) Figure 4.3 shows a synthesis of $\text{CH}_2(\text{CN})\text{COOH}$.




Figure 4.3

- (i) The reaction mechanism of the synthesis in Figure 4.3 has a single step.

Complete Figure 4.3 to show this mechanism. Include lone pairs, charges, dipoles and curly arrows as appropriate.

[2]

- (ii) Identify the nucleophile in Figure 4.3.

..... [1]

- (iii) Identify the mechanism of the reaction in Figure 4.3.

..... [1]



(d) Figure 4.4 describes a sequence of reactions, starting with $\text{CH}_2(\text{CN})\text{COOH}$, to form **E**.




Figure 4.4

- (i) Identify the type of reaction that occurs in step 1.
 [1]
- (ii) Name HCHO , the reactant used in step 2.
 [1]
- (iii) Deduce the molecular formula of **E**.
 [1]
- (iv) **E** can be polymerised to form **F**.
 Draw the structure of **one** repeat unit of polymer **F**.

[1]



(v) Figure 4.5 shows the infrared spectrum of **E**.




Figure 4.5

Table 4.1

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers)/cm ⁻¹
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3650

Using the information given in Table 4.1, identify with a cross (X) on Figure 4.5 an absorption that can be used to monitor the progress of the polymerisation of **E** to form **F**.

Explain your answer with reference to the relevant bond.

.....

 [2]

[Total: 19]





Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.



The Periodic Table of Elements

Group																																			
1	2	1														13	14	15	16	17	18														
		Key																		2															
		atomic number atomic symbol name relative atomic mass																																	
3	Li lithium 6.9	4	Be beryllium 9.0											5	6	7	8	9	10	He helium 4.0															
11	Na sodium 23.0	12	Mg magnesium 24.3											13	14	15	16	17	18																
19	K potassium 39.1	20	Ca calcium 40.1	21	Sc scandium 45.0	22	Ti titanium 47.9	23	V vanadium 50.9	24	Cr chromium 52.0	25	Mn manganese 54.9	26	Fe iron 55.8	27	Co cobalt 58.9	28	Ni nickel 58.7	29	Cu copper 63.5	30	Zn zinc 65.4	31	Ga gallium 69.7	32	Ge germanium 72.6	33	As arsenic 74.9	34	Se selenium 79.0	35	Br bromine 79.9	36	Kr krypton 83.8
37	Rb rubidium 85.5	38	Sr strontium 87.6	39	Y yttrium 88.9	40	Zr zirconium 91.2	41	Nb niobium 92.9	42	Mo molybdenum 95.9	43	Tc technetium	44	Ru ruthenium 101.1	45	Rh rhodium 102.9	46	Pd palladium 106.4	47	Ag silver 107.9	48	Cd cadmium 112.4	49	In indium 114.8	50	Sn tin 118.7	51	Sb antimony 121.8	52	Te tellurium 127.6	53	I iodine 126.9	54	Xe xenon 131.3
55	Cs caesium 132.9	56	Ba barium 137.3	57–71 lanthanoids		72	Hf hafnium 178.5	73	Ta tantalum 180.9	74	W tungsten 183.8	75	Re rhenium 186.2	76	Os osmium 190.2	77	Ir iridium 192.2	78	Pt platinum 195.1	79	Au gold 197.0	80	Hg mercury 200.6	81	Tl thallium 204.4	82	Pb lead 207.2	83	Bi bismuth 209.0	84	Po polonium	85	At astatine	86	Rn radon
87	Fr francium	88	Ra radium	89–103 actinoids		104	Rf rutherfordium	105	Db dubnium	106	Sg seaborgium	107	Bh bohrium	108	Hs hassium	109	Mt meitnerium	110	Ds darmstadtium	111	Rg roentgenium	112	Cn copernicium	113	Nh nihonium	114	Fl flerovium	115	Mc moscovium	116	Lv livermorium	117	Ts tennessine	118	Og oganeson

lanthanoids

57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —

actinoids

