



# Cambridge International AS & A Level

CANDIDATE  
NAMECENTRE  
NUMBER

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

CANDIDATE  
NUMBER

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

## CHEMISTRY

9701/42

Paper 4 A Level Structured Questions

February/March 2026

2 hours

You must answer on the question paper.

No additional materials are needed.

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

### INFORMATION

- The total mark for this paper is 100.
- The number of marks for each question or part question is shown in brackets [ ].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **28** pages. Any blank pages are indicated.




1 Chromium is a transition element that forms coloured compounds.

(a) (i) Define transition element.

.....  
 ..... [1]

(ii) Complete the electrons-in-boxes diagram to show the electronic configuration of an isolated gaseous chromium atom.



[1]

(iii) Sketch the shape of a  $3d_{z^2}$  orbital in Figure 1.1.




Figure 1.1

[1]

(iv) State **two** reasons that explain why chromium compounds can behave as catalysts.

1 .....  
 .....  
 2 .....  
 .....

[2]

- (b) Table 1.1 lists standard electrode potentials,  $E^\ominus$ , for some electrode reactions of chromium and other species.

Table 1.1

| electrode reaction                                                                                                                                            | $E^\ominus/V$ |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------|
| $\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cr}(\text{s})$                                                                              | -0.91         |
| $\text{Cr}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Cr}^{2+}(\text{aq})$                                                                         | -0.41         |
| $\text{Cr}(\text{OH})_3(\text{s}) + \text{e}^- \rightleftharpoons \text{Cr}(\text{OH})_2(\text{s}) + \text{OH}^-(\text{aq})$                                  | -1.26         |
| $\text{Cr}_2\text{O}_7^{2-}(\text{aq}) + 14\text{H}^+(\text{aq}) + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+}(\text{aq}) + 7\text{H}_2\text{O}(\text{l})$ | +1.33         |
| $\text{CrO}_4^{2-}(\text{aq}) + 4\text{H}_2\text{O}(\text{l}) + 3\text{e}^- \rightleftharpoons \text{Cr}(\text{OH})_3(\text{s}) + 5\text{OH}^-(\text{aq})$    | -0.13         |
| $\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}(\text{l})$                                                | +1.23         |
| $\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightleftharpoons 4\text{OH}^-(\text{aq})$                                               | +0.40         |

- (i) Define standard electrode potential,  $E^\ominus$ , including a description of standard conditions.

.....

.....

.....

..... [2]

- (ii) Explain fully, with reference to relevant data in Table 1.1, why solutions of  $\text{Cr}^{2+}(\text{aq})$  need to be stored in an atmosphere of argon instead of in air.

.....

.....

..... [2]

- (iii) A green precipitate of  $\text{Cr}(\text{OH})_3$  forms when aqueous  $\text{NaOH}$  is added to aqueous  $\text{CrCl}_3$ .

Construct an ionic equation for the formation of this green precipitate.

..... [1]




- (iv) Green  $\text{Cr}(\text{OH})_3(\text{s})$  dissolves in alkali to form a yellow solution containing  $\text{CrO}_4^{2-}(\text{aq})$  when it is left in air for several hours.

Use Table 1.1 to construct an ionic equation for this reaction.

..... [2]

- (c) The equation for the decomposition of ammonium dichromate,  $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$ , is shown.



$$\Delta S^\ominus = +915 \text{ JK}^{-1} \text{ mol}^{-1}$$

- (i) Explain why the standard entropy change,  $\Delta S^\ominus$ , is positive in reaction 1.

.....  
 ..... [1]

- (ii) Table 1.2 gives some standard entropy values relevant to reaction 1.


**Table 1.2**

| substance                                        | standard entropy, $S^\ominus$<br>$/\text{JK}^{-1} \text{ mol}^{-1}$ |
|--------------------------------------------------|---------------------------------------------------------------------|
| $(\text{NH}_4)_2\text{Cr}_2\text{O}_7(\text{s})$ | 114                                                                 |
| $\text{Cr}_2\text{O}_3(\text{s})$                | 81                                                                  |
| $\text{N}_2(\text{g})$                           | 192                                                                 |

Use the information in this question to calculate the standard entropy,  $S^\ominus$ , in  $\text{JK}^{-1} \text{ mol}^{-1}$ , of  $\text{H}_2\text{O}(\text{g})$ .

$$S^\ominus \text{ of } \text{H}_2\text{O}(\text{g}) = \dots\dots\dots \text{ JK}^{-1} \text{ mol}^{-1} \quad [2]$$

(iii) Thermodynamic data for reaction 1 is shown.



$$\Delta H^\ominus = -341 \text{ kJ mol}^{-1}$$

$$\Delta S^\ominus = +915 \text{ J K}^{-1} \text{ mol}^{-1}$$

statement 1 Reaction 1 is feasible at all temperatures.

statement 2 Reaction 1 will **not** occur unless  $(\text{NH}_4)_2\text{Cr}_2\text{O}_7$  is heated strongly.

Explain statements 1 and 2. Refer to the Gibbs equation in your answer.

.....

.....

.....

..... [2]

(d) Figure 1.2 shows the Born–Haber diagram for the ionic solid  $\text{Cr}_2\text{O}_3$ . Relevant data is given in Table 1.3.

$\Delta H_{i1-3}$  represents the sum of the first, second and third ionisation energies of Cr.

$\Delta H_{\text{EA}1-2}$  represents the sum of the first and second electron affinities of O.




Figure 1.2

Table 1.3

| enthalpy change                   | value/kJ mol <sup>-1</sup> |
|-----------------------------------|----------------------------|
| $\Delta H_{\text{at}}(\text{Cr})$ | +398                       |
| $\Delta H_{\text{at}}(\text{O})$  | +248                       |
| $\Delta H_{\text{i}1-3}$          | +5233                      |
| $\Delta H_{\text{EA}1-2}$         | +702                       |
| $\Delta H_{\text{f}}$             | -1128                      |

(i) Define first electron affinity.

.....  
 .....  
 ..... [2]

(ii) Use Figure 1.2 and data in Table 1.3 to calculate the lattice energy,  $\Delta H_{\text{latt}}$ , in kJ mol<sup>-1</sup>, of Cr<sub>2</sub>O<sub>3</sub>.

$$\Delta H_{\text{latt}} = \dots\dots\dots \text{kJ mol}^{-1} \quad [2]$$

(iii) The ionic radius of Cr<sup>3+</sup> is 0.069 nm. The ionic radius of Al<sup>3+</sup> is 0.050 nm.

Suggest how the lattice energies of Cr<sub>2</sub>O<sub>3</sub> and Al<sub>2</sub>O<sub>3</sub> differ.

Explain your answer.


.....  
 .....  
 ..... [2]

[Total: 23]



- 2 The manganate(VII) ion,  $\text{MnO}_4^-$ , is used in chemistry as an oxidising agent.

Three reactions each involve  $\text{MnO}_4^-$  as a reagent.



- (a) Oxygen slowly oxidises  $\text{Fe}^{2+}(\text{aq})$ . Reaction 2 can be used to determine the percentage of  $\text{Fe}^{2+}(\text{aq})$  that is oxidised over a certain time.

A student prepares a  $250\text{ cm}^3$  solution using  $17.70\text{ g}$  of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ . After one week, the student titrates  $25.0\text{ cm}^3$  portions of this solution with acidified  $\text{KMnO}_4(\text{aq})$  of concentration  $0.0500\text{ mol dm}^{-3}$ .

The average titre is  $23.80\text{ cm}^3$ .

- (i) Calculate the percentage of  $\text{Fe}^{2+}$  ions in the original sample that were oxidised during one week.

$[M_r: \text{FeSO}_4 \cdot 7\text{H}_2\text{O}, 277.9]$

percentage of  $\text{Fe}^{2+}$  ions oxidised = ..... % [3]

- (ii) Explain why an indicator is **not** needed in this titration.

.....  
 ..... [1]



(b) Reaction 3 uses hot alkaline  $\text{KMnO}_4$  as the reagent. The reaction mixture is then acidified.

Draw the structure of the organic product,  $\text{C}_7\text{H}_6\text{O}_2$ , of reaction 3.

[1]

(c) Reaction 4 is called a reverse disproportionation reaction.

Suggest the meaning of reverse disproportionation. Use oxidation numbers to explain your reasoning.

.....  
 .....  
 .....  
 ..... [2]

(d) The solubility of  $\text{Mn}(\text{OH})_2$  in water is  $3.20 \times 10^{-3} \text{ g dm}^{-3}$  at 298 K.

(i) Write an expression for the solubility product,  $K_{\text{sp}}$ , of  $\text{Mn}(\text{OH})_2$ .

$$K_{\text{sp}} =$$

[1]

(ii) Calculate the value of  $K_{\text{sp}}$  of  $\text{Mn}(\text{OH})_2$  at 298 K. State its units.

value of  $K_{\text{sp}} =$  .....

units .....

[3]





(e) (i)  $\text{Mn}^{3+}(\text{aq})$  ions are coloured. Explain why.

.....

.....

.....

.....

.....

..... [3]

(ii)  $\text{Mn}^{3+}$  ions form the complex ion  $[\text{Mn}(\text{C}_2\text{O}_4)_3]^{3-}$ , which shows stereoisomerism.

Complete Table 2.1 to give details of this complex. Use  $\text{O} \quad \text{O}$  to represent the ligands in the three-dimensional structures of the stereoisomers of  $[\text{Mn}(\text{C}_2\text{O}_4)_3]^{3-}$ .

Table 2.1

|                                          |                                            |  |
|------------------------------------------|--------------------------------------------|--|
| formula of complex                       | $[\text{Mn}(\text{C}_2\text{O}_4)_3]^{3-}$ |  |
| formula of ligand                        |                                            |  |
| shape of complex                         |                                            |  |
| coordination number of Mn in complex     |                                            |  |
| type of stereoisomerism shown by complex | optical                                    |  |
| structures of stereoisomers              |                                            |  |

[4]

[Total: 18]



- 3 Figure 3.1 shows a reaction scheme starting from 4-hydroxybutanoic acid.






Figure 3.1

- (a) Under acidic conditions, 4-hydroxybutanoic acid exists in equilibrium with compound **G**,  $\text{C}_4\text{H}_6\text{O}_2$ , and water. **G** does **not** react with Na.

4-hydroxybutanoic acid can also react to form condensation polymer **H**, which has formula  $(\text{C}_4\text{H}_6\text{O}_2)_n$ .

Draw the structures of **G** and **H**. Show only **one** repeat unit of **H**.



[2]

- (b) Construct an equation for the reaction of 4-hydroxybutanoic acid with an excess of  $\text{PCl}_5$ , forming **J**.

..... [2]



(c) 4-hydroxybutanoic acid,  $\text{HO}(\text{CH}_2)_3\text{COOH}$ , is a weak acid with  $K_a = 1.91 \times 10^{-5} \text{ mol dm}^{-3}$  at  $25^\circ\text{C}$ .

(i) Calculate the pH of a  $0.150 \text{ mol dm}^{-3}$  solution of  $\text{HO}(\text{CH}_2)_3\text{COOH}$  at  $25^\circ\text{C}$ .

pH = ..... [2]

(ii) When  $\text{HO}(\text{CH}_2)_3\text{COOH}(\text{aq})$  is mixed with its conjugate base, a buffer solution can form.

Define buffer solution.

.....  
 .....  
 ..... [2]

(iii) State the structural formula of the conjugate base of  $\text{HO}(\text{CH}_2)_3\text{COOH}$ .

..... [1]

(iv) A student makes a buffer solution by reacting  $25.0 \text{ cm}^3$  of  $0.150 \text{ mol dm}^{-3}$   $\text{HO}(\text{CH}_2)_3\text{COOH}$  with  $10.0 \text{ cm}^3$  of  $0.200 \text{ mol dm}^{-3}$   $\text{NaOH}(\text{aq})$ .

Calculate the pH of this buffer solution.

$K_a$  of  $\text{HO}(\text{CH}_2)_3\text{COOH} = 1.91 \times 10^{-5} \text{ mol dm}^{-3}$  at  $25^\circ\text{C}$ .

pH of buffer solution = ..... [3]



- (d) 4-hydroxybutanoic acid,  $\text{HO}(\text{CH}_2)_3\text{COOH}$ , can be used to produce butanedioic acid,  $\text{HOOC}(\text{CH}_2)_2\text{COOH}$ .

Suggest a suitable reagent and conditions for this reaction.

..... [1]


- (e) A student does an experiment to determine the partition coefficient,  $K_{\text{pc}}$ , of butanedioic acid between the immiscible solvents ethoxyethane and water.

- (i) State what is meant by partition coefficient.

..... [1]

- (ii) The student adds butanedioic acid to a flask containing  $25.0 \text{ cm}^3$  of ethoxyethane and  $25.0 \text{ cm}^3$  of water. The mixture is shaken and left to separate into two layers.

- $10.00 \text{ cm}^3$  of the ethoxyethane layer requires  $7.50 \text{ cm}^3$  of  $0.100 \text{ mol dm}^{-3}$   $\text{NaOH}(\text{aq})$  for complete reaction.
- $10.00 \text{ cm}^3$  of the aqueous layer requires  $11.10 \text{ cm}^3$  of  $0.500 \text{ mol dm}^{-3}$   $\text{NaOH}(\text{aq})$  for complete reaction.



Calculate the concentration of butanedioic acid in each solvent.

Hence, determine the value of the partition coefficient,  $K_{\text{pc}}$ , of butanedioic acid between ethoxyethane and water.

concentration of butanedioic acid in ethoxyethane = .....  $\text{mol dm}^{-3}$

concentration of butanedioic acid in water = .....  $\text{mol dm}^{-3}$

$K_{\text{pc}} =$  ..... [2]


[Total: 16]

[Turn over]



- 4 A student performs a series of experiments to investigate the chemistry of the ester methyl ethanoate,  $\text{CH}_3\text{COOCH}_3$ .

(a)  $\text{CH}_3\text{COOCH}_3$  is hydrolysed under acidic conditions.



The rate of reaction 5 is investigated.

A small amount of  $\text{H}^+(\text{aq})$  is added to a dilute solution of  $\text{CH}_3\text{COOCH}_3(\text{aq})$ . The graph in Figure 4.1 shows how  $[\text{CH}_3\text{COOCH}_3]$  changes with time.




Figure 4.1

- (i) Use Figure 4.1 to show that reaction 5 is first order with respect to  $[\text{CH}_3\text{COOCH}_3]$ .

Show any working fully and explain your reasoning.

.....

.....

.....

..... [2]



(ii) Use Figure 4.1 to calculate the initial rate of reaction in  $\text{mol dm}^{-3} \text{s}^{-1}$ .

initial rate of reaction = .....  $\text{mol dm}^{-3} \text{s}^{-1}$  [1]

(iii) Reaction 5 is first order with respect to  $[\text{H}^+(\text{aq})]$ .

Use this statement and information from 4(a)(i) to construct the overall rate equation for the hydrolysis reaction.

State the units of the rate constant,  $k$ , of this reaction.

rate = .....

units of  $k$  = ..... [2]

(b) Methyl ethanoate,  $\text{CH}_3\text{COOCH}_3$ , forms when  $\text{CH}_3\text{OH}$  reacts with either  $\text{CH}_3\text{COCl}$  or  $\text{CH}_3\text{COOH}$ .


Table 4.1

| reaction | reactants                                                               | products                                                                 |
|----------|-------------------------------------------------------------------------|--------------------------------------------------------------------------|
| 6        | $\text{CH}_3\text{COCl}(\text{l})$ and $\text{CH}_3\text{OH}(\text{l})$ | $\text{CH}_3\text{COOCH}_3(\text{l})$ and $\text{HCl}(\text{g})$         |
| 7        | $\text{CH}_3\text{COOH}(\text{l})$ and $\text{CH}_3\text{OH}(\text{l})$ | $\text{CH}_3\text{COOCH}_3(\text{l})$ and $\text{H}_2\text{O}(\text{l})$ |

(i) Suggest why reaction 6 gives a higher yield of  $\text{CH}_3\text{COOCH}_3$  than reaction 7.

.....  
 .....  
 .....  
 ..... [2]

(ii) Complete Figure 4.2 to describe the addition–elimination mechanism of reaction 6.



Include all relevant curly arrows, charges and dipoles.





Figure 4.2


[4]

(c) The relative rate of hydrolysis of chloroethane, ethanoyl chloride and chlorobenzene is investigated.

chloroethane



ethanoyl chloride



chlorobenzene




Figure 4.3

State the relative rate of hydrolysis of these three compounds. Explain your answer.

..... < ..... < .....

slowest fastest

.....

.....

.....

.....

.....

.....

[3]

[Total: 14]





Turn over



- 5 (a) 4-methylphenol can be synthesised from methylbenzene in three steps, as shown in Figure 5.1.




Figure 5.1


- (i) In step 1, methylbenzene reacts with the  $\text{NO}_2^+$  electrophile.

Write an equation for the formation of  $\text{NO}_2^+$ .

..... [1]

- (ii) Draw the mechanism for the reaction, in step 1, of  $\text{NO}_2^+$  with methylbenzene.

Include all relevant curly arrows and charges. Draw the organic intermediate. Include the by-product of the reaction on the dotted line.



- (iii) 1-methyl-4-nitrobenzene,  $\text{CH}_3\text{C}_6\text{H}_4\text{NO}_2$ , is reduced in step 2.

Write an equation for step 2.

Use  $[\text{H}]$  to represent one atom of hydrogen from the reducing agent.

..... [1]

- (iv) Step 3 is a two-stage process. Identify the reagents and conditions necessary for the two stages.

1 .....

2 ..... [2]

- (v) A student suggests an alternative synthesis of 4-methylphenol from nitrobenzene. The student decides to use a Friedel–Crafts alkylation reaction as the first step, as shown in Figure 5.2.




**Figure 5.2**

Suggest the reagents for this Friedel–Crafts alkylation reaction.

Explain why this reaction will produce a low yield of 1-methyl-4-nitrobenzene.

reagents .....

explanation .....

.....

..... [2]

(b) The structures of three different phenols are shown in Figure 5.3.




**Figure 5.3**

A mixture of these three phenols is analysed by gas/liquid chromatography.

The mobile phase is helium gas.

Figure 5.4 shows the gas/liquid chromatogram that is produced.




**Figure 5.4**

(i) Explain why helium gas is used as the mobile phase.

..... [1]

(ii) Suggest a stationary phase for use in gas/liquid chromatography.

..... [1]








(iii) Suggest why 2-ethylphenol has the longest retention time.

..... [1]

(iv) Table 5.1 shows the area underneath each peak in the chromatogram in Figure 5.4.

### Table 5.1

| compound                  | phenol<br> | 4-methylphenol<br> | 2-ethylphenol<br> |
|---------------------------|---------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| area underneath each peak | 29                                                                                          | 34                                                                                                   | 22                                                                                                   |

The area underneath each peak is proportional to the mass of the respective compound in the mixture.

Calculate the percentage by mass of 2-ethylphenol in the mixture.

percentage by mass of 2-ethylphenol = ..... % [1]


**(c)** Describe the relative acidities of phenol, water and ethanol,  $\text{C}_2\text{H}_5\text{OH}$ . Explain your answer.

weakest acid , strongest acid



- 6 (a) Glutathione is a naturally occurring compound.



glutathione



Glutathione is hydrolysed to form three amino acids.

One of the amino acids is cysteine,  $\text{H}_2\text{NCH}(\text{CH}_2\text{SH})\text{COOH}$ .

- (i) Draw the structures of the two other amino acids that are formed by this hydrolysis.



[2]

- (ii) Cysteine has an isoelectric point of 5.0.

Explain what is meant by isoelectric point.

.....  
 ..... [1]

- (iii) Draw the structure of cysteine at its isoelectric point.






[1]



- (b) **X**, **Y** and **Z** are other amino acids that contain sulfur. Table 6.1 gives some details of these three compounds.

Table 6.1

| amino acid        | <b>X</b>                                                                          | <b>Y</b>                                                                           | <b>Z</b>                                                                            |
|-------------------|-----------------------------------------------------------------------------------|------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|
|                   |  |  |  |
| $M_r$             | 135.1                                                                             | 149.1                                                                              | 149.1                                                                               |
| isoelectric point | 6.0                                                                               | 5.7                                                                                | 4.9                                                                                 |

A mixture of these three amino acids, **X**, **Y** and **Z**, is analysed by electrophoresis, using a buffer solution at pH 5.5.




Figure 6.1

Draw and label **three** spots on Figure 6.1 to indicate the predicted position of each of these three amino acids, **X**, **Y** and **Z**, after electrophoresis at pH 5.5.

Explain your answer.

.....

.....


.....

.....

.....

[3]


(c) Anion  $Z^-$  is a bidentate ligand.



(i) Define bidentate ligand.

.....  
 .....  
 ..... [2]

(ii) Draw three-dimensional diagrams to show the two enantiomers of  $Z^-$ .



[1]

(iii) Suggest **one** reason why it is desirable to use a single enantiomer of  $Z^-$  as a drug.

.....  
 ..... [1]

Octahedral complex **C** forms when two moles of  $Z^-$  react with one mole of  $\text{Cu}^{2+}(\text{aq})$ , as shown in Figure 6.2.





Figure 6.2



(iv) Complete the equation to describe the formation of octahedral complex **C** in Figure 6.2.



[Total: 12]





### Important values, constants and standards

|                                 |                                                                                                                                           |
|---------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------|
| molar gas constant              | $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$                                                                                              |
| Faraday constant                | $F = 9.65 \times 10^4 \text{ C mol}^{-1}$                                                                                                 |
| Avogadro constant               | $L = 6.022 \times 10^{23} \text{ mol}^{-1}$                                                                                               |
| electronic charge               | $e = -1.60 \times 10^{-19} \text{ C}$                                                                                                     |
| molar volume of gas             | $V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K)<br>$V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions |
| ionic product of water          | $K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))                                                             |
| specific heat capacity of water | $c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g <sup>-1</sup> K <sup>-1</sup> )                                                    |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.



The Periodic Table of Elements

| Group                        |                               |                                                                       |                               |                                 |                                |                               |                                |                              |                                |                               |                               |                               |                               |                               |                                |                              |                             |                         |                               |                             |                               |                            |                              |
|------------------------------|-------------------------------|-----------------------------------------------------------------------|-------------------------------|---------------------------------|--------------------------------|-------------------------------|--------------------------------|------------------------------|--------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|--------------------------------|------------------------------|-----------------------------|-------------------------|-------------------------------|-----------------------------|-------------------------------|----------------------------|------------------------------|
| 1                            | 2                             | 1<br>H<br>hydrogen<br>1.0                                             |                               |                                 |                                |                               |                                |                              |                                |                               |                               |                               |                               |                               |                                | 13                           | 14                          | 15                      | 16                            | 17                          | 18                            |                            |                              |
| 3<br>Li<br>lithium<br>6.9    | 4<br>Be<br>beryllium<br>9.0   | Key<br>atomic number<br>atomic symbol<br>name<br>relative atomic mass |                               |                                 |                                |                               |                                |                              |                                |                               |                               |                               |                               |                               |                                |                              |                             | 5<br>B<br>boron<br>10.8 | 6<br>C<br>carbon<br>12.0      | 7<br>N<br>nitrogen<br>14.0  | 8<br>O<br>oxygen<br>16.0      | 9<br>F<br>fluorine<br>19.0 | 10<br>Ne<br>neon<br>20.2     |
|                              | 11<br>Na<br>sodium<br>23.0    | 12<br>Mg<br>magnesium<br>24.3                                         |                               |                                 |                                |                               |                                |                              |                                |                               |                               |                               |                               |                               |                                |                              |                             |                         | 13<br>Al<br>aluminium<br>27.0 | 14<br>Si<br>silicon<br>28.1 | 15<br>P<br>phosphorus<br>31.0 | 16<br>S<br>sulfur<br>32.1  | 17<br>Cl<br>chlorine<br>35.5 |
| 19<br>K<br>potassium<br>39.1 | 20<br>Ca<br>calcium<br>40.1   | 21<br>Sc<br>scandium<br>45.0                                          | 22<br>Ti<br>titanium<br>47.9  | 23<br>V<br>vanadium<br>50.9     | 24<br>Cr<br>chromium<br>52.0   | 25<br>Mn<br>manganese<br>54.9 | 26<br>Fe<br>iron<br>55.8       | 27<br>Co<br>cobalt<br>58.9   | 28<br>Ni<br>nickel<br>58.7     | 29<br>Cu<br>copper<br>63.5    | 30<br>Zn<br>zinc<br>65.4      | 31<br>Ga<br>gallium<br>69.7   | 32<br>Ge<br>germanium<br>72.6 | 33<br>As<br>arsenic<br>74.9   | 34<br>Se<br>selenium<br>79.0   | 35<br>Br<br>bromine<br>79.9  | 36<br>Kr<br>krypton<br>83.8 |                         |                               |                             |                               |                            |                              |
| 37<br>Rb<br>rubidium<br>85.5 | 38<br>Sr<br>strontium<br>87.6 | 39<br>Y<br>yttrium<br>88.9                                            | 40<br>Zr<br>zirconium<br>91.2 | 41<br>Nb<br>niobium<br>92.9     | 42<br>Mo<br>molybdenum<br>95.9 | 43<br>Tc<br>technetium<br>—   | 44<br>Ru<br>ruthenium<br>101.1 | 45<br>Rh<br>rhodium<br>102.9 | 46<br>Pd<br>palladium<br>106.4 | 47<br>Ag<br>silver<br>107.9   | 48<br>Cd<br>cadmium<br>112.4  | 49<br>In<br>indium<br>114.8   | 50<br>Sn<br>tin<br>118.7      | 51<br>Sb<br>antimony<br>121.8 | 52<br>Te<br>tellurium<br>127.6 | 53<br>I<br>iodine<br>126.9   | 54<br>Xe<br>xenon<br>131.3  |                         |                               |                             |                               |                            |                              |
| 55<br>Cs<br>caesium<br>132.9 | 56<br>Ba<br>barium<br>137.3   | lanthanoids<br>57–71                                                  |                               | 73<br>Ta<br>tantalum<br>180.9   | 74<br>W<br>tungsten<br>183.8   | 75<br>Re<br>rhenium<br>186.2  | 76<br>Os<br>osmium<br>190.2    | 77<br>Ir<br>iridium<br>192.2 | 78<br>Pt<br>platinum<br>195.1  | 79<br>Au<br>gold<br>197.0     | 80<br>Hg<br>mercury<br>200.6  | 81<br>Tl<br>thallium<br>204.4 | 82<br>Pb<br>lead<br>207.2     | 83<br>Bi<br>bismuth<br>209.0  | 84<br>Po<br>polonium<br>—      | 85<br>At<br>astatine<br>—    | 86<br>Rn<br>radon<br>—      |                         |                               |                             |                               |                            |                              |
| 87<br>Fr<br>francium<br>—    | 88<br>Ra<br>radium<br>—       | actinoids<br>89–103                                                   |                               | 104<br>Rf<br>rutherfordium<br>— | 106<br>Sg<br>seaborgium<br>—   | 107<br>Bh<br>bohrium<br>—     | 108<br>Hs<br>hassium<br>—      | 109<br>Mt<br>meitnerium<br>— | 110<br>Ds<br>darmstadtium<br>— | 111<br>Rg<br>roentgenium<br>— | 112<br>Cn<br>copernicium<br>— | 113<br>Nh<br>nihonium<br>—    | 114<br>Fl<br>flerovium<br>—   | 115<br>Mc<br>moscovium<br>—   | 116<br>Lv<br>livermorium<br>—  | 117<br>Ts<br>tennessine<br>— | 118<br>Og<br>ognesson<br>—  |                         |                               |                             |                               |                            |                              |

|             |  |                                |                              |                                   |                                |                             |                               |                               |                                 |                              |                                 |                              |                             |                               |                                |                               |
|-------------|--|--------------------------------|------------------------------|-----------------------------------|--------------------------------|-----------------------------|-------------------------------|-------------------------------|---------------------------------|------------------------------|---------------------------------|------------------------------|-----------------------------|-------------------------------|--------------------------------|-------------------------------|
| lanthanoids |  | 57<br>La<br>lanthanum<br>138.9 | 58<br>Ce<br>cerium<br>140.1  | 59<br>Pr<br>praseodymium<br>140.9 | 60<br>Nd<br>neodymium<br>144.2 | 61<br>Pm<br>promethium<br>— | 62<br>Sm<br>samarium<br>150.4 | 63<br>Eu<br>europium<br>152.0 | 64<br>Gd<br>gadolinium<br>157.3 | 65<br>Tb<br>terbium<br>158.9 | 66<br>Dy<br>dysprosium<br>162.5 | 67<br>Ho<br>holmium<br>164.9 | 68<br>Er<br>erbium<br>167.3 | 69<br>Tm<br>thulium<br>168.9  | 70<br>Yb<br>ytterbium<br>173.1 | 71<br>Lu<br>lutetium<br>175.0 |
| actinoids   |  | 89<br>Ac<br>actinium<br>—      | 90<br>Th<br>thorium<br>232.0 | 91<br>Pa<br>protactinium<br>231.0 | 92<br>U<br>uranium<br>238.0    | 93<br>Np<br>neptunium<br>—  | 94<br>Pu<br>plutonium<br>—    | 95<br>Am<br>americium<br>—    | 96<br>Cm<br>curium<br>—         | 97<br>Bk<br>berkelium<br>—   | 98<br>Cf<br>californium<br>—    | 99<br>Es<br>einsteinium<br>— | 100<br>Fm<br>fermium<br>—   | 101<br>Md<br>mendelevium<br>— | 102<br>No<br>nobelium<br>—     | 103<br>Lr<br>lawrencium<br>—  |

