

**MARK SCHEME for the May/June 2011 question paper  
for the guidance of teachers**

**9701 CHEMISTRY**

**9701/42**

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.


Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.





Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2011	9701	42

2 (a) (i)



[2] + [2]

(ii) m. pt. trend: (from) giant/macro molecular/covalent to metallic bonding (or implied from at least two specific examples, e.g. diamond and tin) [1]  
(mention of *simple* covalent anywhere negates this mark)

conductivity trend: increasing delocalisation of electrons (down the group) [1]  
or  $e^-$  are more free-moving  
(or implied from at least two examples, e.g. Si is semiconductor, lead has delocalised  $e^-$ ) [6]

(b) (i) heat  $PbO_2$ , or  $T > 200^\circ C$  or  $\Delta$  on arrow:  $PbO_2 \rightarrow PbO + \frac{1}{2}O_2$  (N.B.  $\frac{1}{2}O_2$  NOT [O]) [1]

(ii) (burning CO in air produces  $CO_2$ ):  $CO + \frac{1}{2}O_2 \rightarrow CO_2$  [1]  
blue flame (ignore ref to limewater test) [1]

(iii) e.g.  $SnCl_2(aq)$  will turn  $KMnO_4$  from purple to colourless [1]  
 $5Sn^{2+} + 2MnO_4^- + 16H^+ \rightarrow 5Sn^{4+} + 2Mn^{2+} + 8H_2O$  [1]

or  $SnCl_2(aq)$  will turn  $K_2Cr_2O_7$  from orange to green [1]  
 $3Sn^{2+} + Cr_2O_7^{2-} + 14H^+ \rightarrow 3Sn^{4+} + 2Cr^{3+} + 7H_2O$  [1]

or  $SnCl_2(aq)$  will turn  $Fe^{3+}$  from orange/brown/yellow to green/colourless [1]  
 $Sn^{2+} + 2Fe^{3+} \rightarrow Sn^{4+} + 2Fe^{2+}$  [1]

or  $SnCl_2(aq)$  will turn  $Cu^{2+}(aq)$  from blue to colourless or give a pink/brown/copper-coloured ppt. [1]  
 $Sn^{2+} + Cu^{2+} \rightarrow Sn^{4+} + Cu$  [1]


Other possible oxidants ( $E^\ominus$  must be  $> +0.2V$ ) include:  $S_2O_8^{2-}$ ,  $H_2O_2$ ,  $Cl_2$ ,  $Br_2$ ,  $I_2$  and  $Ag^+$ . No observations with the first three of these, but this should be stated explicitly, e.g. "no colour change". [5]



[Total: 11 max 10]

3 (a)  $L = F/e$  or  $F = Le$

[1]  
[1]

(b) (i)



[ allow the conventional symbol  to represent  (the "P.S." is not required) ]

correct cell (2 electrodes + PS circuit) [1]

ammeter in series [1]

anode and cathode of the right polarity [IN WORDS] [1]

$\text{CuSO}_4(\text{aq})$  or  $\text{CuCl}_2(\text{aq})$  or  $\text{Cu}^{2+}(\text{aq})$  or soln or  $1 \text{ mol dm}^{-3}$  [1]

(ii)  $n(\text{Cu}) = (52.542 - 52.243) / 63.5 = 4.71 \times 10^{-3} \text{ mol}$  ( $4.67 \times 10^{-3}$ ) [1]  
 $n(e^-)$  required =  $4.71 \times 10^{-3} \times 2 = 9.42 \times 10^{-3} \text{ mol}$  ( $9.34 \times 10^{-3}$ ) ecf [1]

amount of electricity passed =  $0.5 \times 30 \times 60 = 900 \text{ C}$  [1]

no. of electrons passed =  $900 / 1.6 \times 10^{-19} = 5.625 \times 10^{21}$  ecf [1]

no of electrons/ $n(e^-) = L = 5.625 \times 10^{21} / 9.42 \times 10^{-3} = 5.97 \times 10^{23} \text{ mol}^{-1}$  ( $6.02 \times 10^{23}$ ) ecf [1]

(values in italics are if candidate has used  $A_r = 64$ , not 63.5. No last mark if not 3 s.f.:  
 correct ans = [5]) [9]

(c)

compound	product at anode	product at cathode
AgF	$\text{O}_2$	Ag
$\text{FeSO}_4$	$\text{O}_2$	$\text{H}_2$
$\text{MgBr}_2$	$\text{Br}_2$	$\text{H}_2$

6 correct  $\Rightarrow$  [5]  
 5 correct  $\Rightarrow$  [4] etc.


Names can be used instead of symbols. If the atomic symbol (e.g. Br or H or O) is used instead of the molecular formula (e.g.  $\text{Br}_2$  etc.) then deduct [1] mark only for the whole table.

[5]

[Total: 15]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2011	9701	42

4 (a) (i) (allow displayed, structural or skeletal formula)



chain [1]  
repeat unit [1]


(ii) **C** should be  $\text{CH}_2=\text{CHOH}$  (or skeletal formula) [1]

(iii) **C** is  $\text{CH}_3\text{CH}=\text{O}$  (or skeletal formula) [1]

(iv) e.g. add (2,4-)DNPH or DNP or Brady's reagent ecf [1]  
orange or red ppt forms (NOT yellow) ecf [1]  
(or could use Fehling's or Tollens',  
or  $\text{H}^+ + \text{Cr}_2\text{O}_7^{2-}$ : orange to green, or  $\text{H}^+ + \text{MnO}_4^-$ : purple to colourless)

[6]

(b) (i) (allow displayed, structural or skeletal formula)



**D**

correct repeat unit bracketed (any 3 atoms in chain) [1]

(ii) ester [1]

(iii) **E** is  $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CO}_2\text{H}$  (or skeletal structure etc.) (2-hydroxybutanoic acid) [1]  
allow ecf here from the formula of the repeat unit shown in (b)(i)


(iv) condensation (polymerisation) [1]

(v) they have the same "molecular" formula or  $\text{C}_4\text{H}_6\text{O}_2$  (do **NOT** allow empirical formula) or  
same no. and type of atoms or same functional group or both are esters or they are  
isomers [1]  
[5]

<b>Page 6</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – May/June 2011</b>	<b>9701</b>	<b>42</b>

(c) (i) optical isomerism (*or* chiral) [1]

(ii)



**F**

**G**

(letters may be reversed)(allow ecf from **E**, also allow ecf for **G** from **F**) [1] + [1]

cis-trans *or* geometrical isomerism [1]


**[4]**

**[Total: 15]**

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2011	9701	42


- 5 (a) acidity: ethanol < water [1]  
 due to +ve inductive effect of C<sub>2</sub>H<sub>5</sub> group or C<sub>2</sub>H<sub>5</sub> gives e<sup>-</sup> to oxygen or intensifies e<sup>-</sup> (in O-H bond) [1]  
 acidity: phenol > water [1]  
 due to stabilisation of the anion/anionic charge or makes the anion less basic [1]  
**[4]**

(b)



**[5]**

(c) H is



[1]

reagents & conditions:

step 1 **dilute** HNO<sub>3</sub> (dilute, not just 'aq'. H<sub>2</sub>SO<sub>4</sub> negates)

[1]

step 2 Sn/SnCl<sub>2</sub>/Fe + HCl or H<sub>2</sub> + Ni/Pd (NOT H<sub>2</sub> + Pt. NOT LiAlH<sub>4</sub> or NaBH<sub>4</sub>)

[1]

step 3 CH<sub>3</sub>COCl or (CH<sub>3</sub>CO)<sub>2</sub>O ('aq.' negates)

[1]

**[4]**

**[Total: 13]**

Page 8	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2011	9701	42


- 6 (a) They are polar/ionic *or* can hydrogen-bond *or* are hydrophilic. [1]  
(NOT 'contain the –OH group', on its own) [1]

- (b) (i) Primary structure is the sequence/order of amino acids [1]  
Secondary structure is the H-bonding between C=O & N-H *or* peptide group/bonds [1]  
Tertiary structure gives the (overall) 3D structure/shape/folding/globularity  
(not 'coiling' on its own)  
*or* mention of at least one method of forming the 3° structure, e.g.; hydrogen bonding  
**between R-groups/side chains**; –S-S- bridges; van der Waals forces; ionic interactions [1]

- (ii) The 3° structure provides a complementary shape to that of the substrate  
*or* it provides the right/specifically shaped cavity for the substrate. (NOT just 'a cleft')  
*or* provides nearby groups to aid the reactions of the substrate (owtte) [1]

- (iii) Two conditions out of the following:  
(a) Increased temperature  
(b) Decreased temperature  
(c) Change in pH  
(d) Addition of heavy metals (*or* specified, e.g. Hg/Ag)  
(e) Addition of inhibitors (competitive or non-competitive)  
Suitable reasons:  
(i) 3D structure changes shape/is deformed/is broken *or* R-R interactions (or a specific example, e.g. H-bonding) are broken  
(ii) inhibitor occupies active site. [2]  
(iii) *either* fewer substrate molecules with  $E > E_a$  *or* fewer successful collisions [6]

- (c) (i)



- left hand peak labelled as pepsin [1]  
right hand peak labelled as trypsin [1]  
(Correct enzymes, but wrong way round, scores [1] only)

- (ii) Peak between pH 6 and pH 8, **and** correct name (amylase) [1]  
[3]

[Total: 10]





Page 10	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – May/June 2011	9701	42

- 8 (a) Range should be from  $10^{-6}$ – $10^{-7}$  (the left hand arrow) [1]  
to  $10^{-8}$ – $10^{-9}$  (the right hand arrow) [1]  
[2]
- (b) Forms of the **same element** (or of **carbon**, since carbon is the context of the question) [1]  
with different structures/arrangements of atoms [1]  
allow 'different molecular structure', but not structural formula. Any mention of 'compound'  
negates the mark. [2]
- (c) Nanoparticles are smaller than (animal) cells or they can pass through the cell membrane  
or pass into/between cells [1]  
Drugs can be bound to/enclosed by the nanoparticle [1]  
[2]
- (d) (i) Reduction/redox [1]
- (ii)  $M_r$  of chalcopyrite is  $63.5 + 56 + 64 = 183.5$   
Mass of copper present is 63.5
- Hence percentage of copper present =  $\frac{63.5 \times 100}{183.5} = 34.6\%$  [1]  
(if  $A_r(\text{Cu}) = 64$  is used, ans = **34.8%**. allow **34–35%**)
- (iii) *If the ore contains 2% of chalcopyrite by mass, calculate how much copper is produced  
from each tonne of ore.*
- 1 tonne = 1000 kg  
1 tonne of chalcopyrite would produce 346 kg of copper  
1 tonne of 2 % ore would produce  $346 \times 0.02$  or **6.9** kg of copper ecf from (d)(ii) [1]  
(accept **7.0** or 7 kg)  
answer may be given as 7000 g or  $7 \times 10^{-3}$  tonnes. If no units are given, assume they  
are tonnes, and mark accordingly)
- (iv) By displacement with a metal (the following specified metals higher than Cu in the ECS  
may be used: Fe, Zn, Sn, Pb, Al, Mg. (NOT Ca, Li, Na, K etc.) or with a suitable non-  
metallic reducing agent, e.g.  $\text{SO}_2$  or  $\text{Sn}^{2+}$ , but not something that wouldn't react, like  $\text{H}_2$   
or By electrolysis (with carefully controlled voltage) [1]  
[4]

[Total: 10]