

---

**CHEMISTRY**

**9701/41**

Paper 4 A Level Structured Questions

**October/November 2019**

MARK SCHEME

Maximum Mark: 100

---

**Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

---

This document consists of **14** printed pages.

**PUBLISHED****Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

**GENERIC MARKING PRINCIPLE 1:**

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

**GENERIC MARKING PRINCIPLE 2:**

Marks awarded are always **whole marks** (not half marks, or other fractions).

**GENERIC MARKING PRINCIPLE 3:**

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

**GENERIC MARKING PRINCIPLE 4:**

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

**GENERIC MARKING PRINCIPLE 5:**


Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

**GENERIC MARKING PRINCIPLE 6:**

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.







**PUBLISHED**



Question	Answer	Marks
1(a)	Platinum / Pt      Aluminium / Al <b>BOTH</b>	<b>1</b>
1(b)(i)	<b>M1:</b> use of or quoting a valid Nernst equation $E = E^{\circ} + 0.0590 / z \log [\text{ox}] / [\text{red}]$ <b>OR</b> $E = 0.15 + (0.0590 / 2)\log 2$ <b>M2:</b> $E = (+)0.16$ (0.159) V minimum 2 sig. fig. correct answer scores 2 marks	<b>2</b>
1(b)(ii)	$E_{\text{cell}} = 0.16 - (-1.66) = +1.82 \text{ V}$ minimum 3 sig. fig.	<b>1</b>
1(b)(iii)	$2\text{Al} + 3\text{Sn}^{4+} \rightarrow 2\text{Al}^{3+} + 3\text{Sn}^{2+}$ <b>M1:</b> species <b>M2:</b> balancing	<b>2</b>
1(c)	<b>M1:</b> number of C ( $= 300\,000 \times 60 \times 60 \times 24$ ) $= 2.59 \times 10^{10}$ (C) <b>M2:</b> number of F ( $= 2.592 \times 10^{10} / 9.65 \times 10^4$ ) $= 2.69 \times 10^5$ (moles of electrons) <b>M3:</b> moles of Al ( $= 2.69 \times 10^5 / 3$ ) $= 8.95 \times 10^4$ <b>M4:</b> mass of Al ( $= 8.95 \times 10^4 \times 27$ ) $= 2420 \text{ kg}$ correct answer scores 4 marks	<b>4</b>
1(d)	<b>M1:</b> ( $\text{Cr}^{2+} + 2\text{e}^{-} \rightleftharpoons \text{Cr}$ ) $E^{\circ} = -0.91$ <b>and</b> ( $2\text{H}^{+} + 2\text{e}^{-} \rightleftharpoons \text{H}_2$ ) $E^{\circ} = 0.00$ seen <b>M2:</b> hydrogen formed instead / hydrogen (ions) easier to reduce / hydrogen has more positive $E^{\circ}$	<b>2</b>

Question	Answer	Marks
2(a)	 <p><b>M1:</b> eight electrons around N atom [N=O, N-O, N-Cl with N-O as dative] <b>M2:</b> all other electrons correct</p>	2
2(b)(i)	(rate =) $k[\text{ClNO}_2][\text{NO}]$	1
2(b)(ii)	$\text{mol}^{-1} \text{ dm}^3 \text{ s}^{-1}$	1
2(b)(iii)	Yes <b>AND</b> number of moles of reactants in overall equation is the same as order in rate equation	1
2(c)(i)	<ul style="list-style-type: none"> <li>• straight line with a negative gradient</li> <li>• starting at <math>2.0 \times 10^{-4}</math></li> <li>• reaches at <math>1.8 \times 10^{-4}</math> at 0.2 seconds</li> </ul> <p>Award 1 mark for two points, award 2 marks for all three points</p>	2
2(c)(ii)	$2 \times 10^{-5} \text{ (mol dm}^{-3}\text{)}$	1
2(c)(iii)	The reaction has reached equilibrium	1

**PUBLISHED**

Question	Answer	Marks
3(a)	a <b>measure / degree of disorder / randomness</b> of a system	<b>1</b>
3(b)	<b>M1:</b> negative – molecules have less energy in the system <b>M2:</b> positive – solid being converted into an aqueous solution <b>M3:</b> negative – gaseous ions being converted into a solid	<b>3</b>
3(c)(i)	(standard) Gibbs free energy <u>change</u>	<b>1</b>
3(c)(ii)	<b>M1:</b> $(\Delta)G = \Delta H - T\Delta S$ <b>M2:</b> description of calculating the minimum value of T for which $\Delta G$ is zero / becomes negative <b>OR</b> $T = \Delta H / \Delta S$ [1]	<b>2</b>

Question	Answer			Marks												
4(a)	<b>M1:</b> $\text{CH}_3\text{COCl}$ or ethanoyl chloride <b>M2:</b> $\text{AlCl}_3$ catalyst			<b>2</b>												
4(b)	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th data-bbox="338 347 562 413">reagent</th> <th data-bbox="562 347 958 413">organic product</th> <th data-bbox="958 347 1256 413">name of mechanism</th> </tr> </thead> <tbody> <tr> <td data-bbox="338 413 562 655">Cl</td> <td data-bbox="562 413 958 655">             chlorine atom(s) in side chain only         </td> <td data-bbox="958 413 1256 655">free radical substitution</td> </tr> <tr> <td data-bbox="338 655 562 959">nitric / sulfuric</td> <td data-bbox="562 655 958 959">             one only <math>-\text{NO}_2</math> group added at 3 position         </td> <td data-bbox="958 655 1256 959">electrophilic substitution</td> </tr> <tr> <td data-bbox="338 959 562 1023">Br</td> <td data-bbox="562 959 958 1023">no reaction with <math>\text{Br}_2</math></td> <td data-bbox="958 959 1256 1023" style="background-color: #cccccc;"></td> </tr> </tbody> </table> <p data-bbox="338 1054 1016 1091">Award 1 mark for each correct entry to the table [5]</p>			reagent	organic product	name of mechanism	Cl	 chlorine atom(s) in side chain only	free radical substitution	nitric / sulfuric	 one only $-\text{NO}_2$ group added at 3 position	electrophilic substitution	Br	no reaction with $\text{Br}_2$		<b>5</b>
reagent	organic product	name of mechanism														
Cl	 chlorine atom(s) in side chain only	free radical substitution														
nitric / sulfuric	 one only $-\text{NO}_2$ group added at 3 position	electrophilic substitution														
Br	no reaction with $\text{Br}_2$															
4(c)(i)	nucleophilic addition			<b>1</b>												

Question	Answer	Marks
4(c)(ii)	 <p><b>M1 and M2:</b> first structure – award one mark for two points correct, award two marks for four points correct.</p> <ul style="list-style-type: none"> <li>• correct polarity shown on C=O bond</li> <li>• curly arrow from double bond to O of the C=O</li> <li>• lone pair shown on C of CN<sup>-</sup> ion</li> <li>• curly arrow from C (on CN<sup>-</sup> ion) to C of C=O</li> </ul> <p><b>M3:</b> middle structure</p> <ul style="list-style-type: none"> <li>• correct intermediate <b>AND</b> curly arrow from lone pair on O<sup>-</sup> to H<sup>+</sup> <b>or</b> H of H<sub>2</sub>O or HCN</li> </ul> <p><b>M4:</b> third structure</p> <ul style="list-style-type: none"> <li>• correct product <b>AND</b> either CN<sup>-</sup> reformed (if HCN seen in step 2) <b>OR</b> curly arrow on H-CN bond towards CN in step 2</li> </ul>	4
4(d)(i)		1
4(d)(ii)	LiAlH <sub>4</sub> or NaBH <sub>4</sub>	1
4(d)(iii)	<b>conc</b> H <sub>2</sub> SO <sub>4</sub> / <b>conc</b> H <sub>3</sub> PO <sub>4</sub> / Al <sub>2</sub> O <sub>3</sub>	1
4(d)(iv)	6	1



**PUBLISHED**

Question	Answer	Marks
4(d)(v)	<ul style="list-style-type: none"> <li>• 25–50</li> <li>• 110–160</li> <li>• 190–220</li> </ul> <p>Award 1 mark for two points, award 2 marks for three points</p>	<b>2</b>

Question	Answer	Marks
5(a)(i)	$K_{sp} = [Ag^+]^2[S^{2-}]$	<b>1</b>
5(a)(ii)	<ul style="list-style-type: none"> <li>• <math>[S^{2-}] = 1.16 \times 10^{-17}</math></li> <li>• <math>[Ag^+] = 2.32 \times 10^{-17}</math></li> <li>• <math>K_{sp} = \mathbf{6.2(4)} \times 10^{-51}</math> minimum 2 sig. fig.</li> </ul> <p>correct answer scores 2 marks Award 1 mark for two points, award 2 marks for three points</p>	<b>2</b>
5(a)(iii)	<p><b>M1:</b> moles <math>Ag_2S = 1 / 247.9 = 0.00403</math> moles [1] 2sf min</p> <p><b>M2:</b> <math>1.16 \times 10^{-17} = 0.0040 / V</math> so <math>V = \mathbf{3.5 \times 10^{14}}</math> (dm<sup>3</sup>) [1] 2sf min ecf on M1</p> <p>correct answer scores 2 marks</p>	<b>2</b>
5(b)(i)	<p><b>M1:</b> <math>[H^+] = \sqrt{2.0 \times 10^{-9} \times 0.20}</math> <math>[H^+] = 2.0 \times 10^{-5}</math> (<math>1.9976 \times 10^{-5}</math>)</p> <p><b>M2:</b> pH = <b>4.7</b> (4.699) minimum 2 sig. fig. min</p> <p>correct answer scores 2 marks</p>	<b>2</b>


**PUBLISHED**

Question	Answer	Marks
5(b)(ii)	<p><b>M1:</b> Both equilibria correctly stated  moles KOH = <math>0.005 \times 0.2 = 1 \times 10^{-3}</math>  moles HOBr(initial) = <math>0.020 \times 0.2 = 4 \times 10^{-3}</math>  moles HOBr(eqm) = <math>4 \times 10^{-3} - 1 \times 10^{-3} = 3 \times 10^{-3}</math>  moles BrO<sup>-</sup>(eqm) = <math>1 \times 10^{-3}</math></p> <p><b>M2:</b> ratio [OBr<sup>-</sup>]/[HOBr] = 1/3  [H<sup>+</sup>] = <math>3 \times 2.0 \times 10^{-9} = 6 \times 10^{-9}</math>  pH = <b>8.2(2)</b></p> <p>correct answer scores 2 marks</p>	<b>2</b>

Question	Answer	Marks
6(a)(i)	$\text{Mg(g)} \rightarrow \text{Mg}^{\text{+}}(\text{g}) + \text{e}^{-}$	<b>1</b>
6(a)(ii)	$\text{Sr(s)} + 2\text{H}_2\text{O(l)} \rightarrow \text{Sr(OH)}_2(\text{aq}) + \text{H}_2(\text{g})$	<b>1</b>
6(a)(iii)	more reactive <b>and</b> easier to ionise down the group <b>OR</b> more reactive <b>and</b> ionisation energies decrease down the group	<b>1</b>
6(b)(i)	brown gas <b>and white</b> solid	<b>1</b>
6(b)(ii)	$2\text{Ca(NO}_3)_2 \rightarrow 2\text{CaO} + 4\text{NO}_2 + \text{O}_2$	<b>1</b>
6(b)(iii)	<p><b>M1:</b> more stable (down the group)</p> <p><b>M2:</b> cationic radius <b>increases</b> / charge density of M<sup>2+</sup> decreases (down the group)</p> <p><b>M3:</b> NO<sub>3</sub><sup>-</sup> anion is less polarised / distorted</p>	<b>3</b>

**PUBLISHED**

Question	Answer	Marks
6(c)	<p><b>M1:</b> less soluble / decreases (down the group)</p> <p><b>M2:</b> <math>\Delta H_{\text{latt}}</math> and <math>\Delta H_{\text{hyd}}</math> <b>both</b> decrease / less exothermic down the group</p> <p><b>M3:</b> <math>\Delta H_{\text{hyd}}</math> decreases by more (than <math>\Delta H_{\text{latt}}</math>)</p> <p><b>M4:</b> <math>\Delta H_{\text{sol}}</math> becomes more endothermic / less exothermic</p>	<b>4</b>

Question	Answer	Marks
7(a)	forms one or more stable ions with incomplete / partially filled d-orbitals / d-subshell	<b>1</b>
7(b)(i)	purple to pale pink / colourless <b>AND</b> orange to green	<b>1</b>
7(b)(ii)	$3\text{Sn}^{2+} + \text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ \rightarrow 3\text{Sn}^{4+} + 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	<b>1</b>
7(c)(i)	<ul style="list-style-type: none"> <li>• six</li> <li>• coordinate bonds / dative bonds / lone pairs donated</li> <li>• to the (central) metal ion</li> </ul> <p>award 1 mark for all three points</p>	<b>1</b>
7(c)(ii)	$[\text{Ru}(\text{NH}_3)_4\text{Cl}(\text{SO}_2)]^+$	<b>1</b>
7(c)(iii)	 <p><b>M1:</b> six correct ligands around Ru, bonds are shown from S of <math>\text{SO}_2</math></p> <p><b>M2:</b> 3-D bonds used correctly for octahedral</p> <p><b>M3:</b> cis and trans isomers shown</p>	<b>3</b>


Question	Answer	Marks
7(c)(iv)	cis-trans or geometric(al) [1]	1
7(c)(v)	<p><b>M1:</b> complexes have two sets of <b>d orbital(s)</b> of different energy <b>OR d-orbitals</b> splits into two sets (of orbitals)</p> <p><b>M2:</b> visible light <b>absorbed</b> (and complementary colour observed)</p> <p><b>M3:</b> <b>electron(s)</b> promoted / excited <b>OR electron(s)</b> moves to higher (d–) orbital</p>	3

Question	Answer	Marks
8(a)(i)	<p>any one from:</p> <ul style="list-style-type: none"> <li>OH<sup>-</sup> / NaOH; aqueous / dilute; heat under reflux</li> <li>H<sup>+</sup> / HCl / H<sub>2</sub>SO<sub>4</sub>, aqueous / dilute; heat under reflux</li> <li>protease or named protease; water; T = 30° – 40°C</li> </ul> <p>all three points in each bullet [1]</p>	1
8(a)(ii)	<p><b>M1:</b> three amino acids in any ionic / non-ionic states [1]</p> <p><b>M2:</b> three amino acids in the correct ionic state for their conditions [1]</p>	2

**PUBLISHED**

Question	Answer	Marks
8(b)	<ul style="list-style-type: none"> <li>permanent dipole-dipole</li> <li>one group that will have a <math>\delta^+</math> and another with <math>\delta^-</math> e.g. CO, NH, COOH, OH      <b>BOTH</b> [1]</li> <li>hydrogen bonds</li> <li>one group that will have a <math>H^{\delta+}</math>, e.g. NH, OH and another with lone pair, e.g. NH, COOH, OH, CONH<sub>2</sub>      <b>BOTH</b> [1]</li> <li>ionic bonding</li> <li>NH<sub>3</sub><sup>+</sup> <b>and</b> COO<sup>-</sup>      <b>BOTH</b> [1]</li> </ul> <p><b>ALLOW</b></p> <ul style="list-style-type: none"> <li>London forces</li> <li>C<sub>4</sub>H<sub>9</sub> groups or parts of these alkyl groups</li> </ul>	<b>3</b>
8(c)(i)	any structure containing one COOH / COCl <b>and</b> NH <sub>2</sub> groups in the same molecule [1]	<b>1</b>
8(c)(ii)	HOCH <sub>2</sub> CH <sub>2</sub> OH [1] ethan(e)-1,2-diol [1] <b>ecf</b> for diols  HO <sub>2</sub> CCO <sub>2</sub> H or ClOCCOCl [1] ethan(e)dioic acid or ethan(e)diol chloride [1] <b>ecf</b> for diacids / diacyl chlorides	<b>4</b>

Question	Answer	Marks
9(a)(i)	RNH <sub>2</sub> + H <sup>+</sup> → RNH <sub>3</sub> <sup>+</sup> <b>OR</b> RNH <sub>2</sub> + HCl → RNH <sub>3</sub> Cl [1]	<b>1</b>
9(a)(ii)	weaker <b>AND</b> lone pair of N delocalised into benzene ring [1]	<b>1</b>

Question	Answer	Marks
9(b)	 <p>[1]</p> <p>[1]</p> <p>[1]</p>	3
9(c)(i)	2 [1]	1
9(c)(ii)	CH <sub>2</sub> next to ester <b>and</b> terminal CH <sub>3</sub> are circled [1]	1
9(c)(iii)	<ul style="list-style-type: none"> <li>• one less peak</li> <li>• the lost peak is NH<sub>2</sub> / aryl amine</li> <li>• protons exchange with D <b>OR</b> protons are labile <b>OR</b> valid equation               <ul style="list-style-type: none"> <li>• ✓✓ for two marks [2]</li> </ul> </li> </ul>	2
9(d)	C <sub>6</sub> H <sub>4</sub> NH <sub>2</sub> <sup>+</sup> <b>and</b> CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> <sup>+</sup> [1]	1