

**Cambridge  
International  
AS & A Level**

# **Data Booklet**

**Cambridge International Advanced Subsidiary and Advanced Level  
in Chemistry (9701)**

**For use from 2016 in all papers for the above  
syllabus, except practical examinations.**

CSTxxx



## **Contents: Tables of Chemical Data**

	<b>Page no.</b>
1      Important values, constants and standards	3
2      Ionisation energies (1 <sup>st</sup> , 2 <sup>nd</sup> , 3 <sup>rd</sup> and 4 <sup>th</sup> ) of selected elements in kJ mol <sup>-1</sup>	4
3      Bond energies	5
4      Standard electrode potential and redox potentials, $E^\ominus$ at 298K (25 °C)	7
5      Atomic and ionic radii	10
6      Typical proton ( <sup>1</sup> H) chemical shift values ( $\delta$ ) relative to TMS = 0	12
7      Typical carbon ( <sup>13</sup> C) chemical shift values ( $\delta$ ) relative to TMS = 0	13
8      Characteristic infra-red absorption frequencies for some selected bonds	14
9      The orientating effect of groups in aromatic substitution reactions	15
10     Names, structures and abbreviations of some amino acids	16
11     The Periodic Table of Elements	17

## 1 Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
the Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
the Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
the Planck constant	$h = 6.63 \times 10^{-34} \text{ Js}$
speed of light in a vacuum	$c = 3.00 \times 10^8 \text{ m s}^{-1}$
rest mass of proton, ${}_1^1\text{H}$	$m_p = 1.67 \times 10^{-27} \text{ kg}$
rest mass of neutron, ${}_0^1\text{n}$	$m_n = 1.67 \times 10^{-27} \text{ kg}$
rest mass of electron, ${}_{-1}^0\text{e}$	$m_e = 9.11 \times 10^{-31} \text{ kg}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ under room conditions (where s.t.p. is expressed as 101 kPa, approximately, and 273 K [0 °C])
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K [25 °C])
specific heat capacity of water	$= 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (= 4.18 J g <sup>-1</sup> K <sup>-1</sup> )

**2 Ionisation energies (1st, 2nd, 3rd and 4th) of selected elements, in  $\text{kJ mol}^{-1}$**

	Proton number	First	Second	Third	Fourth
H	1	1310	—	—	—
He	2	2370	5250	—	—
Li	3	519	7300	11800	—
Be	4	900	1760	14800	21000
B	5	799	2420	3660	25000
C	6	1090	2350	4610	6220
N	7	1400	2860	4590	7480
O	8	1310	3390	5320	7450
F	9	1680	3370	6040	8410
Ne	10	2080	3950	6150	9290
Na	11	494	4560	6940	9540
Mg	12	736	1450	7740	10500
Al	13	577	1820	2740	11600
Si	14	786	1580	3230	4360
P	15	1060	1900	2920	4960
S	16	1000	2260	3390	4540
Cl	17	1260	2300	3850	5150
Ar	18	1520	2660	3950	5770
K	19	418	3070	4600	5860
Ca	20	590	1150	4940	6480
Sc	21	632	1240	2390	7110
Ti	22	661	1310	2720	4170
V	23	648	1370	2870	4600
Cr	24	653	1590	2990	4770
Mn	25	716	1510	3250	5190
Fe	26	762	1560	2960	5400
Co	27	757	1640	3230	5100
Ni	28	736	1750	3390	5400
Cu	29	745	1960	3350	5690
Zn	30	908	1730	3828	5980
Ga	31	577	1980	2960	6190
Br	35	1140	2080	3460	4850
Rb	37	403	2632	3900	5080
Sr	38	548	1060	4120	5440
Ag	47	731	2074	3361	5000
I	53	1010	1840	3000	4030
Cs	55	376	2420	3300	4400
Ba	56	502	966	3390	4700

### 3 Bond Energies

#### 3(a) Bond energies in diatomic molecules (these are exact values)

##### Homonuclear

Bond	Energy / $\text{kJ mol}^{-1}$
H—H	436
D—D	442
N≡N	944
O=O	496
P≡P	485
S=S	425
F—F	158
Cl—Cl	242
Br—Br	193
I—I	151

##### Heteronuclear

Bond	Energy / $\text{kJ mol}^{-1}$
H—F	562
H—Cl	431
H—Br	366
H—I	299
C≡O	1077

**3(b) Bond energies in polyatomic molecules (these are average values)**

**Homonuclear**

Bond	Energy / $\text{kJ mol}^{-1}$
C—C	350
C=C	610
C≡C	840
C—C (benzene)	520
N—N	160
N=N	410
O—O	150
Si—Si	225
P—P	200
S—S	265

**Heteronuclear**

Bond	Energy / $\text{kJ mol}^{-1}$
C—H	410
C—Cl	340
C—Br	280
C—I	240
C—N	305
C=N	610
C≡N	890
C—O	360
C=O	740
C=O in $\text{CO}_2$	805
N—H	390
N—Cl	310
O—H	460
Si—Cl	360
Si—H	320
Si—O (in $\text{SiO}_2(\text{s})$ )	460
Si=O (in $\text{SiO}_2(\text{g})$ )	640
P—H	320
P—Cl	330
P—O	340
P=O	540
S—H	340
S—Cl	250
S—O	360
S=O	500

## 4 Standard electrode potential and redox potentials, $E^\ominus$ at 298 K (25 °C)

For ease of reference, two tables are given:

- (a) an extended list in alphabetical order;
- (b) a shorter list in decreasing order of magnitude, i.e. a redox series.

### (a) $E^\ominus$ in alphabetical order

Electrode reaction	$E^\ominus/V$
$\text{Ag}^+ + \text{e}^- \rightleftharpoons \text{Ag}$	+0.80
$\text{Al}^{3+} + 3\text{e}^- \rightleftharpoons \text{Al}$	-1.66
$\text{Ba}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ba}$	-2.90
$\text{Br}_2 + 2\text{e}^- \rightleftharpoons 2\text{Br}^-$	+1.07
$\text{Ca}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ca}$	-2.87
$\text{Cl}_2 + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$	+1.36
$2\text{HOCl} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Cl}_2 + 2\text{H}_2\text{O}$	+1.64
$\text{ClO}^- + \text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{Cl}^- + 2\text{OH}^-$	+0.89
$\text{Co}^{2+} + 2\text{e}^- \rightleftharpoons \text{Co}$	-0.28
$\text{Co}^{3+} + \text{e}^- \rightleftharpoons \text{Co}^{2+}$	+1.82
$[\text{Co}(\text{NH}_3)_6]^{2+} + 2\text{e}^- \rightleftharpoons \text{Co} + 6\text{NH}_3$	-0.43
$\text{Cr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cr}$	-0.91
$\text{Cr}^{3+} + 3\text{e}^- \rightleftharpoons \text{Cr}$	-0.74
$\text{Cr}^{3+} + \text{e}^- \rightleftharpoons \text{Cr}^{2+}$	-0.41
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1.33
$\text{Cu}^+ + \text{e}^- \rightleftharpoons \text{Cu}$	+0.52
$\text{Cu}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu}$	+0.34
$\text{Cu}^{2+} + \text{e}^- \rightleftharpoons \text{Cu}^+$	+0.15
$[\text{Cu}(\text{NH}_3)_4]^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu} + 4\text{NH}_3$	-0.05
$\text{F}_2 + 2\text{e}^- \rightleftharpoons 2\text{F}^-$	+2.87
$\text{Fe}^{2+} + 2\text{e}^- \rightleftharpoons \text{Fe}$	-0.44
$\text{Fe}^{3+} + 3\text{e}^- \rightleftharpoons \text{Fe}$	-0.04
$\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$	+0.77
$[\text{Fe}(\text{CN})_6]^{3-} + \text{e}^- \rightleftharpoons [\text{Fe}(\text{CN})_6]^{4-}$	+0.36
$\text{Fe}(\text{OH})_3 + \text{e}^- \rightleftharpoons \text{Fe}(\text{OH})_2 + \text{OH}^-$	-0.56
$2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2$	0.00
$2\text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{H}_2 + 2\text{OH}^-$	-0.83
$\text{I}_2 + 2\text{e}^- \rightleftharpoons 2\text{I}^-$	+0.54
$\text{K}^+ + \text{e}^- \rightleftharpoons \text{K}$	-2.92
$\text{Li}^+ + \text{e}^- \rightleftharpoons \text{Li}$	-3.04
$\text{Mg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mg}$	-2.38
$\text{Mn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mn}$	-1.18
$\text{Mn}^{3+} + \text{e}^- \rightleftharpoons \text{Mn}^{2+}$	+1.49
$\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$	+1.23
$\text{MnO}_4^- + \text{e}^- \rightleftharpoons \text{MnO}_4^{2-}$	+0.56
$\text{MnO}_4^- + 4\text{H}^+ + 3\text{e}^- \rightleftharpoons \text{MnO}_2 + 2\text{H}_2\text{O}$	+1.67
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1.52
$\text{NO}_3^- + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{NO}_2 + \text{H}_2\text{O}$	+0.81
$\text{NO}_3^- + 3\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{HNO}_2 + \text{H}_2\text{O}$	+0.94
$\text{NO}_3^- + 10\text{H}^+ + 8\text{e}^- \rightleftharpoons \text{NH}_4^+ + 3\text{H}_2\text{O}$	+0.87

Electrode reaction	$E^\ominus/V$
$\text{Na}^+ + \text{e}^- \rightleftharpoons \text{Na}$	-2.71
$\text{Ni}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ni}$	-0.25
$[\text{Ni}(\text{NH}_3)_6]^{2+} + 2\text{e}^- \rightleftharpoons \text{Ni} + 6\text{NH}_3$	-0.51
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1.77
$\text{HO}_2^- + \text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons 3\text{OH}^-$	+0.88
$\text{O}_2 + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1.23
$\text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^- \rightleftharpoons 4\text{OH}^-$	+0.40
$\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2$	+0.68
$\text{O}_2 + \text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{HO}_2^- + \text{OH}^-$	-0.08
$\text{Pb}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pb}$	-0.13
$\text{Pb}^{4+} + 2\text{e}^- \rightleftharpoons \text{Pb}^{2+}$	+1.69
$\text{PbO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Pb}^{2+} + 2\text{H}_2\text{O}$	+1.47
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{SO}_2 + 2\text{H}_2\text{O}$	+0.17
$\text{S}_2\text{O}_8^{2-} + 2\text{e}^- \rightleftharpoons 2\text{SO}_4^{2-}$	+2.01
$\text{S}_4\text{O}_6^{2-} + 2\text{e}^- \rightleftharpoons 2\text{S}_2\text{O}_3^{2-}$	+0.09
$\text{Sn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sn}$	-0.14
$\text{Sn}^{4+} + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}$	+0.15
$\text{V}^{2+} + 2\text{e}^- \rightleftharpoons \text{V}$	-1.20
$\text{V}^{3+} + \text{e}^- \rightleftharpoons \text{V}^{2+}$	-0.26
$\text{VO}^{2+} + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{V}^{3+} + \text{H}_2\text{O}$	+0.34
$\text{VO}_2^+ + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{VO}^{2+} + \text{H}_2\text{O}$	+1.00
$\text{VO}_3^- + 4\text{H}^+ + \text{e}^- \rightleftharpoons \text{VO}^{2+} + 2\text{H}_2\text{O}$	+1.00
$\text{Zn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Zn}$	-0.76

All ionic states refer to aqueous ions but other state symbols have been omitted.

**(b)  $E^\ominus$  in decreasing order of oxidising power**

(a selection only – see also the extended alphabetical list on the previous pages)

Electrode reaction	$E^\ominus / \text{V}$
$\text{F}_2 + 2\text{e}^- \rightleftharpoons 2\text{F}^-$	+2.87
$\text{S}_2\text{O}_8^{2-} + 2\text{e}^- \rightleftharpoons 2\text{SO}_4^{2-}$	+2.01
$\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1.77
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$	+1.52
$\text{PbO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Pb}^{2+} + 2\text{H}_2\text{O}$	+1.47
$\text{Cl}_2 + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$	+1.36
$\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$	+1.33
$\text{O}_2 + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$	+1.23
$\text{Br}_2 + 2\text{e}^- \rightleftharpoons 2\text{Br}^-$	+1.07
$\text{ClO}^- + \text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{Cl}^- + 2\text{OH}^-$	+0.89
$\text{NO}_3^- + 10\text{H}^+ + 8\text{e}^- \rightleftharpoons \text{NH}_4^+ + 3\text{H}_2\text{O}$	+0.87
$\text{NO}_3^- + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{NO}_2 + \text{H}_2\text{O}$	+0.81
$\text{Ag}^+ + \text{e}^- \rightleftharpoons \text{Ag}$	+0.80
$\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$	+0.77
$\text{I}_2 + 2\text{e}^- \rightleftharpoons 2\text{I}^-$	+0.54
$\text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^- \rightleftharpoons 4\text{OH}^-$	+0.40
$\text{Cu}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu}$	+0.34
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{SO}_2 + 2\text{H}_2\text{O}$	+0.17
$\text{Sn}^{4+} + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}$	+0.15
$\text{S}_4\text{O}_6^{2-} + 2\text{e}^- \rightleftharpoons 2\text{S}_2\text{O}_3^{2-}$	+0.09
$2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2$	0.00
$\text{Pb}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pb}$	-0.13
$\text{Sn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sn}$	-0.14
$\text{Fe}^{2+} + 2\text{e}^- \rightleftharpoons \text{Fe}$	-0.44
$\text{Zn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Zn}$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{H}_2 + 2\text{OH}^-$	-0.83
$\text{V}^{2+} + 2\text{e}^- \rightleftharpoons \text{V}$	-1.20
$\text{Mg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mg}$	-2.38
$\text{Ca}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ca}$	-2.87
$\text{K}^+ + \text{e}^- \rightleftharpoons \text{K}$	-2.92

## 5 Atomic and ionic radii

<b>(a) Period 1</b>	<b>atomic / nm</b>	<b>ionic / nm</b>
single covalent	H 0.037	$\text{H}^-$ 0.208
van der Waals	He 0.140	
<b>(b) Period 2</b>	<b>atomic / nm</b>	<b>ionic / nm</b>
metallic	Li 0.152	$\text{Li}^+$ 0.060
	Be 0.112	$\text{Be}^{2+}$ 0.031
single covalent	B 0.080	$\text{B}^{3+}$ 0.020
	C 0.077	$\text{C}^{4+}$ 0.015 $\text{C}^{4-}$ 0.260
	N 0.074	$\text{N}^{3-}$ 0.171
	O 0.073	$\text{O}^{2-}$ 0.140
	F 0.072	$\text{F}^-$ 0.136
van der Waals	Ne 0.160	
<b>(c) Period 3</b>	<b>atomic / nm</b>	<b>ionic / nm</b>
metallic	Na 0.186	$\text{Na}^+$ 0.095
	Mg 0.160	$\text{Mg}^{2+}$ 0.065
	Al 0.143	$\text{Al}^{3+}$ 0.050
single covalent	Si 0.117	$\text{Si}^{4+}$ 0.041 $\text{Si}^{4-}$ 0.271
	P 0.110	$\text{P}^{3-}$ 0.212
	S 0.104	$\text{S}^{2-}$ 0.184
	Cl 0.099	$\text{Cl}^-$ 0.181
van der Waals	Ar 0.190	
<b>(d) Group 2</b>	<b>atomic / nm</b>	<b>ionic / nm</b>
metallic	Be 0.112	$\text{Be}^{2+}$ 0.031
	Mg 0.160	$\text{Mg}^{2+}$ 0.065
	Ca 0.197	$\text{Ca}^{2+}$ 0.099
	Sr 0.215	$\text{Sr}^{2+}$ 0.113
	Ba 0.217	$\text{Ba}^{2+}$ 0.135
	Ra 0.220	$\text{Ra}^{2+}$ 0.140


<b>(e) Group 14</b>	<b>atomic / nm</b>	<b>ionic / nm</b>	
single covalent	C 0.077		
	Si 0.117	Si <sup>4+</sup> 0.041	
	Ge 0.122	Ge <sup>2+</sup> 0.093	
metallic	Sn 0.162	Sn <sup>2+</sup> 0.112	
	Pb 0.175	Pb <sup>2+</sup> 0.120	
<b>(f) Group 17</b>	<b>atomic / nm</b>	<b>ionic / nm</b>	
single covalent	F 0.072	F <sup>-</sup> 0.136	
	Cl 0.099	Cl <sup>-</sup> 0.181	
	Br 0.114	Br <sup>-</sup> 0.195	
	I 0.133	I <sup>-</sup> 0.216	
	At 0.140		
<b>(g) First row transition elements</b>	<b>atomic / nm</b>	<b>ionic / nm</b>	
metallic	Sc 0.164	Sc <sup>3+</sup> 0.081	
	Ti 0.146	Ti <sup>2+</sup> 0.090	Ti <sup>3+</sup> 0.067
	V 0.135	V <sup>2+</sup> 0.079	V <sup>3+</sup> 0.064
	Cr 0.129	Cr <sup>2+</sup> 0.073	Cr <sup>3+</sup> 0.062
	Mn 0.132	Mn <sup>2+</sup> 0.067	Mn <sup>3+</sup> 0.062
	Fe 0.126	Fe <sup>2+</sup> 0.061	Fe <sup>3+</sup> 0.055
	Co 0.125	Co <sup>2+</sup> 0.078	Co <sup>3+</sup> 0.053
	Ni 0.124	Ni <sup>2+</sup> 0.070	Ni <sup>3+</sup> 0.056
	Cu 0.128	Cu <sup>2+</sup> 0.073	
	Zn 0.135	Zn <sup>2+</sup> 0.075	

## 6 Typical proton ( $^1\text{H}$ ) chemical shift values ( $\delta$ ) relative to TMS = 0

type of proton	environment of proton	example structures	chemical shift range ( $\delta$ )
C–H	alkane	$-\text{CH}_3$ , $-\text{CH}_2-$ , $>\text{CH}-$	0.9–1.7
	alkyl next to C=O	$\text{CH}_3\text{—C=O}$ , $-\text{CH}_2\text{—C=O}$ , $>\text{CH—C=O}$	2.2–3.0
	alkyl next to aromatic ring	$\text{CH}_3\text{—Ar}$ , $-\text{CH}_2\text{—Ar}$ , $>\text{CH—Ar}$	2.3–3.0
	alkyl next to electronegative atom	$\text{CH}_3\text{—O}$ , $-\text{CH}_2\text{—O}$ , $-\text{CH}_2\text{—Cl}$ , $>\text{CH—Br}$	3.2–4.0
	attached to alkyne	$\equiv\text{C—H}$	1.8–3.1
	attached to alkene	$=\text{CH}_2$ , $=\text{CH—}$	4.5–6.0
	attached to aromatic ring		6.0–9.0
O-H (see note below)	aldehyde		9.3–10.5
	alcohol	$\text{RO—H}$	0.5–6.0
	phenol		4.5–7.0
N-H (see note below)	carboxylic acid		9.0–13.0
	alkyl amine	$\text{R—NH—}$	1.0–5.0
	aryl amine		3.0–6.0
	amide		5.0–12.0

Note:  $\delta$  values for  $-\text{O-H}$  and  $-\text{N-H}$  protons can vary depending on solvent and concentration

## 7 Typical carbon ( $^{13}\text{C}$ ) chemical shift values ( $\delta$ ) relative to TMS = 0


hybridisation of the carbon atom	environment of carbon atom	example structures	chemical shift range ( $\delta$ )
$\text{sp}^3$	alkyl	$\text{CH}_3-$ , $-\text{CH}_2-$ , $-\text{CH}-$ , $>\text{C}-$	0–50
$\text{sp}^3$	next to alkene/arene	$\begin{array}{c}   \\ -\text{C}-\text{C}=\text{C}- \\   \end{array}$ , $\begin{array}{c}   \\ -\text{C}-\text{C}_6\text{H}_5- \\   \end{array}$	10–40
$\text{sp}^3$	next to carbonyl/carboxyl	$\begin{array}{c}   \\ -\text{C}-\text{COR}, -\text{C}-\text{CO}_2\text{R}, \\   \end{array}$	25–50
$\text{sp}^3$	next to nitrogen	$\begin{array}{c}   \\ -\text{C}-\text{NH}_2, -\text{C}-\text{NR}_2, -\text{C}-\text{NHCO} \\   \end{array}$	30–65
$\text{sp}^3$	next to chlorine ( $-\text{CH}_2\text{-Br}$ and $-\text{CH}_2\text{-I}$ are in the same range as alkyl)	$\begin{array}{c}   \\ -\text{C}-\text{Cl} \\   \end{array}$	30–60
$\text{sp}^3$	next to oxygen	$\begin{array}{c}   \\ -\text{C}-\text{OH}, -\text{C}-\text{O}-\text{CO}- \\   \end{array}$	50–70
$\text{sp}^2$	alkene or arene	$>\text{C}=\text{C}<$ , 	110–160
$\text{sp}^2$	carboxyl	$\text{R}-\text{CO}_2\text{H}$ , $\text{R}-\text{CO}_2\text{R}$	160–185
$\text{sp}^2$	carbonyl	$\text{R}-\text{CHO}$ , $\text{R}-\text{CO}-\text{R}$	190–220
$\text{sp}$	alkyne	$\text{R}-\text{C}\equiv\text{C}-$	65–85
$\text{sp}$	nitrile	$\text{R}-\text{C}\equiv\text{N}$	100–125

## 8 Characteristic infra-red absorption frequencies for some selected bonds

bond	functional groups containing the bond	absorption range (in wavenumbers) /cm <sup>-1</sup>	appearance of peak ( <i>s</i> = strong, <i>w</i> = weak)
C–O	alcohols, ethers, esters	1040–1300	<b>s</b>
C=C	aromatic compounds, alkenes	1500–1680	<i>w</i> unless conjugated
C=O	amides, ketones and aldehydes, carboxylic acids, esters	1640–1690 1670–1740 1680–1730 1710–1750	<b>s</b> <b>s</b> <b>s</b> <b>s</b>
C≡C	alkynes	2150–2250	<i>w</i> unless conjugated
C≡N	nitriles	2200–2250	<b>w</b>
C–H	alkanes, CH <sub>2</sub> –H alkenes/arenes, =C–H	2850–2950 3000–3100	<b>s</b> <i>w</i>
N–H	amines, amides	3300–3500	<i>w</i>
O–H	carboxylic acids, RCO <sub>2</sub> –H H–bonded alcohol, RO–H free alcohol, RO–H	2500–3000 3200–3600 3580–3650	<b>s</b> and very broad <b>s</b> <b>s</b> and sharp

## 9 The orientating effect of groups in aromatic substitution reactions.

The position of the incoming group, **Y**, is determined by the nature of the group, **X**, already bonded to the ring, and not by the nature of the incoming group **Y**.



<b>X</b> - groups that direct the incoming <b>Y</b> group to the 2- or 4- positions	<b>X</b> - groups that direct the incoming <b>Y</b> group to the 3- position
$-\text{NH}_2$ , $-\text{NHR}$ or $-\text{NR}_2$	$-\text{NO}_2$
$-\text{OH}$ or $-\text{OR}$	$-\text{NH}_3^+$
$-\text{NHCOR}$	$-\text{CN}$
$-\text{CH}_3$ , $-\text{alkyl}$	$-\text{CHO}$ , $-\text{COR}$
$-\text{Cl}$	$-\text{CO}_2\text{H}$ , $-\text{CO}_2\text{R}$

## 10 Names, structures and abbreviations of some amino acids

name	3-letter abbreviation	1-letter symbol	structure of side chain R- in
			$  \begin{array}{c}  \text{NH}_2 \\    \\  \text{R} - \text{CH} \\    \\  \text{CO}_2\text{H}  \end{array}  $
alanine	Ala	A	$\text{CH}_3-$
aspartic acid	Asp	D	$\text{HO}_2\text{CCH}_2-$
cysteine	Cys	C	$\text{HSCH}_2-$
glutamic acid	Glu	E	$\text{HO}_2\text{CCH}_2\text{CH}_2-$
glycine	Gly	G	$\text{H}-$
lysine	Lys	K	$\text{H}_2\text{NCH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$
phenylalanine	Phe	F	
serine	Ser	S	$\text{HOCH}_2-$
tyrosine	Tyr	Y	
valine	Val	V	

# The Periodic Table of Elements

Group																											
1	2											13	14	15	16	17	18										
<b>Key</b>																											
3 Li lithium 6.9	4 Be beryllium 9.0											1 H hydrogen 1.0															
11 Na sodium 23.0	12 Mg magnesium 24.3											5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2										
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8										
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium –	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3										
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids –	72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium –	85 At astatine –	86 Rn radon –										
87 Fr francium –	88 Ra radium –	89–103 actinoids –	104 Rf rutherfordium –	105 Db dubnium –	106 Sg seaborgium –	107 Bh bohrium –	108 Hs hassium –	109 Mt meitnerium –	110 Ds darmstadtium –	111 Rg roentgenium –	112 Cn copernicium –	114 Fl flerovium –	116 Lv livermorium –	–	–	–	–										

lanthanoids	57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.4	61 Pm promethium –	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
	89 Ac actinium –	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium –	94 Pu plutonium –	95 Am americium –	96 Cm curium –	97 Bk berkelium –	98 Cf californium –	99 Es einsteinium –	100 Fm fermium –	101 Md mendelevium –	102 No nobelium –	103 Lr lawrencium –

**BLANK PAGE**

## **BLANK PAGE**

## **BLANK PAGE**