

MARK SCHEME for the May/June 2013 series

9709 MATHEMATICS

9709/12

Paper 1, maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9709	12

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9709	12

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only - often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR -1	A penalty of MR -1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through ✓" marks. MR is not applied when the candidate misreads his own figures - this is regarded as an error in accuracy. An MR-2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA -1	This is deducted from A or B marks in the case of premature approximation. The PA -1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9709	12

<p>1 $\frac{dy}{dx} = \frac{6}{x^2}$ $y = -6x^{-1} + c$ Uses (2, 9) $\rightarrow c = 12$ $y = -6x^{-1} + 12$</p>	<p>B1 M1 A1 [3]</p>	<p>Integration only – unsimplified Uses (2, 9) in an integral</p>
<p>2 $\left(2x - \frac{1}{2x}\right)^6$ (i) Coeff of $x^2 = 15 \times 16 \times (-\frac{1}{2})^2 = 60$ (ii) Constant term is $20 \times 8x^3 \times (-1 \div 8x^3) \times (1 + x^2)$ needs to consider 2 terms $\rightarrow 60 - 20 = 40$</p>	<p>B1 B1 [2] B1 M1 A1 [3]</p>	<p>B1 for 2/3 parts. B1 B1 unsimplified Needs to consider the constant term</p>
<p>3 $mx + 14 = \frac{12}{x} + 2 \rightarrow mx^2 + 12x - 12 = 0$ Uses $b^2 = 4ac \rightarrow m = -3$ $-3x^2 + 12x - 12 = 0 \rightarrow P(2, 8)$ [Or $m = -12x^{-2}$ M1 Sub M1 $x = 2$ A1] [$\rightarrow m = -3$ and $y = 8$ M1 A1]</p>	<p>M1 M1 A1 DM1 A1 [5]</p>	<p>Eliminates x (or y) Any use of discriminant Any valid method.</p>
<p>4 (i) $BOC = 2 \tan^{-1} \frac{1}{2} = 0.9273$ (ii) $OB = \sqrt{10^2 + 5^2}$ or $11.2 = r$ Arc $BXC = \sqrt{125} \times 0.9273$ \rightarrow Perimeter = 20.4 cm (iii) Area = $\frac{1}{2}r^2\theta$ $-\frac{1}{2} \cdot 10 \cdot 10 \rightarrow 7.96 \text{ cm}^2$.</p>	<p>M1 A1 [2] B1 M1 A1 [3] M1 A1 [2]</p>	<p>Correct trigonometry. (ans given) Use of trig (or Pyth) for the $OB = \sqrt{125}$. Use of $s = r\theta$ with θ in rads, $r \neq 10$ Correct formula used with rads, $r \neq 10$. Allow 7.95 or 7.96</p>
<p>5 $a = \sin \theta - 3 \cos \theta$, $b = 3 \sin \theta + \cos \theta$ (i) $a^2 + b^2 =$ $(s^2 + 9c^2 - 6sc) + (9s^2 + c^2 + 6sc)$ $10c^2 + 10s^2 = 10$ (ii) $2s - 6c = 3s + c \rightarrow s = -7c$ $\rightarrow \tan \theta = -7$ $\rightarrow 98.1^\circ$ and 278.1°</p>	<p>B1 M1 A1 [3] M1 A1 A1 A1 [4]</p>	<p>Correct squaring Use of $s^2 + c^2 = 1$ to get constant. (can get 2/3 for missing $6sc$) Collecting and $t = s \div c$ For 180° + first answer, providing no extra answers in the range.</p>

Page 5	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9709	12

<p>6 $\vec{OA} = \mathbf{i} - 2\mathbf{j} + 2\mathbf{k}$, $\vec{OB} = 3\mathbf{i} + p\mathbf{j} + q\mathbf{k}$</p> <p>(i) $p = -6, q = 6$</p> <p>(ii) dot product = 0 $\rightarrow 3 - 2p + 4p = 0$ $\rightarrow p = -1.5$</p> <p>(iii) $\vec{AB} = \mathbf{b} - \mathbf{a} = 2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}$ Unit vector = $(2\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}) \div 7$</p>	<p>B1 B1 [2]</p> <p>M1 A1 [2]</p> <p>B1 M1 A1[†] [3]</p>	<p>Use of $x_1x_2 + y_1y_2 + z_1z_2 = 0$</p> <p>not for $\mathbf{b} - \mathbf{a}$.</p> <p>M1 for division by modulus. [†] on B1.</p>
<p>7 $3y + 2x = 33$.</p> <p>Gradient of line = $-\frac{2}{3}$</p> <p>Gradient of perpendicular = $3/2$</p> <p>Eqn of perp $y - 3 = \frac{3}{2}(x + 1)$</p> <p>Sim Eqns $\rightarrow (3, 9)$</p> <p>$(-1, 3) \rightarrow (3, 9) \rightarrow (7, 15)$</p>	<p>B1 M1 M1 M1 A1</p> <p>M1 A1 [7]</p>	<p>Use of $m_1m_2 = -1$ with gradient of line</p> <p>Correct form of perpendicular eqn.</p> <p>Sim eqns.</p> <p>Vectors or other method.</p>
<p>8 (i) $\pi r^2 h = 250\pi \rightarrow h = \frac{250}{r^2}$ $\rightarrow S = 2\pi r h + 2\pi r^2$ $\rightarrow S = 2\pi r^2 + \frac{500\pi}{r}$</p> <p>(ii) $\frac{dS}{dr} = 4\pi r - \frac{500\pi}{r^2}$ = 0 when $r^3 = 125 \rightarrow r = 5$ $\rightarrow S = 150\pi$</p> <p>(iii) $\frac{d^2S}{dr^2} = 4\pi + \frac{1000\pi}{r^3}$ This is positive \rightarrow Minimum</p>	<p>M1</p> <p>M1 [2]</p> <p>B1 B1</p> <p>M1 A1 [4]</p> <p>M1 A1 [2]</p>	<p>Makes h the subject. $\pi r^2 h$ must be right</p> <p>Ans given – check all formulae..</p> <p>B1 for each term</p> <p>Sets differential to 0 + attempt at soln</p> <p>Any valid method. 2nd differential must be correct – no need for numerical answer or correct r.</p>
<p>9 $f(x) = \frac{5}{1-3x}, x \geq 1$</p> <p>(i) $f'(x) = \frac{-5}{(1-3x)^2} \times -3$</p> <p>(ii) $15 > 0$ and $(1-3x)^2 > 0, f'(x) > 0$ \rightarrow increasing</p> <p>(iii) $y = \frac{5}{1-3x} \rightarrow 3x = 1 - \frac{5}{y}$ $\rightarrow f^{-1}(x) = \frac{x-5}{3x}$ or $\frac{1}{3} - \frac{5}{3x}$</p> <p>Range is ≥ 1 Domain is $-2.5 \leq x < 0$</p>	<p>B1 B1 [2]</p> <p>B1[†] [1]</p> <p>M1 A1</p> <p>B1 B1 B1 [5]</p>	<p>B1 without $\times -3$. B1 for $\times -3$, even if first B mark is incorrect</p> <p>[†] providing $()^2$ in denominator.</p> <p>Attempt to make x the subject. Must be in terms of x.</p> <p>must be \geq condone $<$</p>

Page 6	Mark Scheme	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2013	9709	12

<p>10 (a) $57 = 2(24 + 3d) \rightarrow d = 1.5$ $48 = 12 + (n - 1)1.5 \rightarrow n = 25$</p> <p>(b) $ar^2 = 4a \quad r = \pm 2$ $\frac{a(r^6 - 1)}{r - 1} = ka$ $\rightarrow k = 63 \text{ or } k = -21$</p>	<p>M1 A1 M1 A1 [4] B1 B1 B1 B1 [4]</p>	<p>Use of correct S_n formula. Use of correct T_n formula. (allow for $r = 2$)</p>
<p>11 $y = \sqrt{1 + 4x}$</p> <p>(i) $\frac{dy}{dx} = \frac{1}{2}(1 + 4x)^{-\frac{1}{2}} \times 4$ $= 2$ at $B(0, 1)$ Gradient of normal $= -\frac{1}{2}$ Equation $y - 1 = -\frac{1}{2}x$</p> <p>(ii) At $A \quad x = -\frac{1}{4}$ $\int \sqrt{1 + 4x} dx = \frac{(1 + 4x)^{\frac{3}{2}}}{\frac{3}{2}} \div 4$ Limits $-\frac{1}{4}$ to $0 \rightarrow \frac{1}{6}$ Area $BOC = \frac{1}{2} \times 2 \times 1 = 1$ \rightarrow Shaded area $= \frac{7}{6}$</p>	<p>B1 B1 M1 M1 A1 [5] B1 B1 B1 B1 B1^h [5]</p>	<p>B1 Without “$\times 4$”. B1 for “$\times 4$” even if first B mark lost. Use of $m_1 m_2 = -1$ Correct method for eqn. B1 Without the “$\div 4$”. For “$\div 4$” even if first B mark lost. For 1 + his “$1/6$”.</p>