

Cambridge International AS & A Level

MATHEMATICS**9709/51**

Paper 5 Probability & Statistics 1

May/June 2025**MARK SCHEME**

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2025 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

This document consists of **19** printed pages.

PUBLISHED
Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics-Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

Annotations guidance for centres

Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

Annotation	Meaning
	More information required
	Accuracy mark awarded zero
	Accuracy mark awarded one
	Independent accuracy mark awarded zero
	Independent accuracy mark awarded one
	Independent accuracy mark awarded two
	Benefit of the doubt
	Blank Page
	Incorrect
Dep	Used to indicate DM0 or DM1

Annotation	Meaning
DM1	Dependent on the previous M1 mark(s)
FT	Follow through
	Indicate working that is right or wrong
Highlighter	Highlight a key point in the working
ISW	Ignore subsequent work
J	Judgement
JU	Judgement
M0	Method mark awarded zero
M1	Method mark awarded one
M2	Method mark awarded two
MR	Misread
O	Omission or Other solution
Off-page comment	Allows comments to be entered at the bottom of the RM marking window and then displayed when the associated question item is navigated to.
On-page comment	Allows comments to be entered in speech bubbles on the candidate response.
PE	Judgment made by the PE
Pre	Premature approximation
SC	Special case
SEEN	Indicates that work/page has been seen

Annotation	Meaning
SF	Error in number of significant figures
	Correct
TE	Transcription error
XP	Correct answer from incorrect working

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

DM or DB When a part of a question has two or more ‘method’ steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

FT Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.

- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
- For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
- The total number of marks available for each question is shown at the bottom of the Marks column.
- Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
- Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no ‘follow through’ from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To

Question	Answer	Marks	Guidance
1	$P(Z < \frac{1.48 - \mu}{0.14}) = 0.22$	B1	$0.771 < z < 0.773$ or $-0.773 < z < -0.771$ seen.
	$\frac{1.48 - \mu}{0.14} = -0.772$	M1	Use of the \pm standardisation formula with μ , 1.48 and 0.14 and equating to a z -value (not 0.78, 0.22, 0.5871, 0.7823, 0.228). Condone σ^2 , $\sqrt{\sigma}$ and continuity correction ± 0.005 .
	$\mu = 1.59$	A1	$1.585 < \mu \leq 1.59$. If M0 scored SC B1 for correct answer WWW.
		3	

Question	Answer	Marks	Guidance
2(a)	Method 1 total arrangements with As together – arrangements with As together and Os together		
	$\frac{7!}{2!} - 6!$	M1	$\frac{7!}{2} - k$, where k is an integer ≥ 1 .
	$n - 6!$	M1	$n - 6!$, where n is an integer > 720 .
	1800	A1	
		3	
	Method 2 arrangements of other 6 letters with the As together and then the Os placed		
	$5! \times \frac{6 \times 5}{2}$	M1	$5! \times p$, where p is an integer > 1 .
		M1	$q \times \frac{6 \times 5}{2}$, $q \times {}^6C_2$, $q \times {}^6P_2$, $q \times 6 \times 5$, q an integer > 1 .
	1800	A1	
		3	

Question	Answer	Marks	Guidance
2(b)	Method 1 $[P(X < 6) =] 1 - \left(\frac{3}{4}\right)^5$	M1	$1 - \left(\frac{3}{4}\right)^a \quad a = 5, 6.$
	$= 0.763, \frac{781}{1024}$	A1	$0.7626 < p \leq 0.763.$
		2	
	Method 2 $[P(X < 6) =] \frac{1}{4} + \left(\frac{1}{4}\right)\left(\frac{3}{4}\right) + \left(\frac{1}{4}\right)\left(\frac{3}{4}\right)^2 + \left(\frac{1}{4}\right)\left(\frac{3}{4}\right)^3 + \left(\frac{1}{4}\right)\left(\frac{3}{4}\right)^4$	M1	$\frac{1}{4} + \left(\frac{1}{4}\right)\left(\frac{3}{4}\right) + \left(\frac{1}{4}\right)\left(\frac{3}{4}\right)^2 + \left(\frac{1}{4}\right)\left(\frac{3}{4}\right)^3 + \left(\frac{1}{4}\right)\left(\frac{3}{4}\right)^4 \left[+ \left(\frac{1}{4}\right)\left(\frac{3}{4}\right)^5 \right].$
	$= 0.763, \frac{781}{1024}$	A1	$0.7626 < p \leq 0.763.$
		2	
2(c)	$\left(\frac{1}{4}\right)^2 \left(\frac{3}{4}\right)^4 \times 5 = \frac{405}{4096}$	M1	$p^2 (1-p)^4 \times 5, \quad 0 < p < 1, \quad p \neq 1-p.$
	$= 0.0989$	A1	AWRT
		2	

Question	Answer			Marks	Guidance
3(a)	Gulls		Herons	B1	Correct stem, ignore extra values (not in reverse, not split). If a split stem-and-leaf plot is used (i.e. stem values are repeated), the remaining B marks are available.
	9 7			B1	Correct Gulls labelled on left, leaves in order from right to left and lined up vertically (less than halfway to next column), no commas or other punctuation.
	8 6 6 3 2 8	1 3 5 7		B1	Correct Herons labelled on same diagram, leaves in order and lined up vertically (less than halfway to next column), no commas or other punctuation.
	8 7 2 9	2 3 5 7 9 9		B1	Penalise each error only once in question. E.g. commas in both sets of data
	4 0 10	8		B1	If the correct data for Gulls & Herons is transposed, treat as a single error in Gulls and condone in Herons.
				B1	Correct key for their diagram, need both clubs labelled and ‘sec’ or ‘s’ stated at least once here, or in leaf headings or title. SC: If 2 separate diagrams drawn, max marks: B1 if both stems correct, B1 if Gulls is correct to the left of the stem B0 B1 if both keys correct including ‘sec’ or ‘s’
				4	

Question	Answer	Marks	Guidance
3(b)	Median = 8.8 (seconds)	B1	Clearly identified, e.g. Q2, med, m..
	[LQ 8.3, UQ 9.8] 9.8 – 8.3	M1	$9.7 \leq UQ \leq 10.0 - 8.2 \leq LQ \leq 8.6$. Implied if both quartile values are stated and the appropriate IQR is calculated accurately.
	1.5	A1	WWW If M0 scored SCB1 for 1.5 WWW.
		3	
3(c)	$\text{Mean} = \frac{175.0 + 30 \times 8.4}{50} \text{ or } \frac{175.0 + 252.0}{50}$	M1	
	= 8.54	A1	If M0 scored, SCB1 for 8.54 WWW.
		2	
3(d)	$1.38^2 = \frac{(1823.0 + \sum y^2)}{50} - \text{their} \left(\frac{427.0}{50} \right)^2$	M1	Substitute values into correct variance formula, any form.
	Solve to find $\sum y^2$	DM1	Rearrange equation to obtain $\sum y^2$. $\frac{\sum y^2}{50} = 1.38^2 + \text{their} \left(\frac{427.0}{50} \right)^2 - \frac{1823.0}{50}.$
	$\sum y^2 = 1918.8$	A1	Mark final answer. If one or both M marks not awarded, SCB1 for 1918.8 seen as final answer WWW mark final answer.
		3	

Question	Answer	Marks	Guidance
4(a)	$P\left(\frac{-1.2}{2.5} < Z < \frac{1.2}{2.5}\right)$	M1	OE Using \pm standardisation formula, not σ^2 , not σ , no continuity correction Use of \pm standardisation formula once with 32.4, 2.5 and either 31.2 or 33.6. No continuity correction, not σ^2 , not $\sqrt{\sigma}$ Implied by either $\frac{-1.2}{2.5}$ or $\frac{1.2}{2.5}$ seen
	$[\Phi(0.48) + \Phi(-0.48) = 2\Phi(0.48) - 1]$ $2 \times 0.6844 - 1$ or $2 \times (0.6844 - 0.5)$ or $0.6844 - 0.3156$	M1	Calculating the correct probability area (leading to their final probability). This may be implied by the correct or appropriate probability area.
	0.3688	A1	
	Expected number = $0.3688 \times 600 = 221.28$ so 221 or 222	B1 FT	FT their 4-figure probability to obtain a single integer answer. No approximation indicated, condone use of 3sf probability here if more accurate answer seen earlier.
		4	

Question	Answer	Marks	Guidance
4(b)	Method 1 $[P(2 \leq X < 8) = 1 - P(0, 1, 8, 9) =]$ $1 - (0.4^9 + {}^9C_1 0.4^8 0.6 + {}^9C_8 0.4^1 0.6^8 + 0.6^9) =$ $[1 - 0.000262144 - 0.00353894 - 0.060466176 - 0.010077696 =]$	M1 A1	One term of the form ${}^9C_x (p)^x (1-p)^{9-x}$, $0 < p < 1, x \neq 0$ or 9 Correct un-simplified expression. Condone omission of last bracket only. If both brackets omitted in un-simplified expression, allow recovery for final stated calculation of $1 - 0.07434\dots$ or better.
	0.926	B1	$0.925 < p \leq 0.926$ from correct working.
	Method 2 $[P(2 \leq X < 8) = P(2, 3, 4, 5, 6, 7) =]$ ${}^9C_2 0.4^7 0.6^2 + {}^9C_3 0.4^6 0.6^3 + {}^9C_4 0.4^5 0.6^4 + {}^9C_5 0.4^4 0.6^5 + {}^9C_6 0.4^3 0.6^6 + {}^9C_7 0.4^2 0.6^7$	M1 A1	One term of the form ${}^9C_x (p)^x (1-p)^{9-x}$, $0 < p < 1, x \neq 0$ or 9. Correct un-simplified expression.
	0.926	B1	$0.925 < p \leq 0.926$ from correct working.
		3	

Question	Answer	Marks	Guidance
4(c)	Mean = $80 \times 0.6 = 48$ Variance = $80 \times 0.6 \times 0.4 = 19.2$	B1	48 and 19.2 (CAO) seen, allow un-simplified. May be in standardisation formula. (4.38178... to at least 4SF identified as σ implies correct variance). Do not condone clear incorrect identification.
	$[P(X > 50) =] P(Z > \frac{50.5 - 48}{\sqrt{19.2}})$	M1	Substituting <i>their</i> 48 and <i>their</i> 19.2 into the \pm standardising formula (any number for 50.5), allow σ^2 or $\sqrt{\sigma}$.
		M1	Use continuity correction 49.5 or 50.5 in <i>their</i> standardisation formula. Note: If no working $\pm \left(\frac{2.5}{\sqrt{19.2}} \right)$ or $\pm \left(\frac{2.5}{4.382} \right)$ seen gains B2 .
	$[P(Z > 0.571) = 1 - \Phi(0.571) =]$ $1 - 0.7160 =$	M1	Appropriate probability area, from final process, must be a probability. May be implied by a sketch of the required probability area. Expect final answer < 0.5 .
	0.284	A1	AWRT
		5	

Question	Answer	Marks	Guidance
5(a)	3G 2P 1D: ${}^9C_3 \times {}^6C_2 \times {}^5C_1 = 6300$ 4G 1P 1D: ${}^9C_4 \times {}^6C_1 \times {}^5C_1 = 3780$ 5G 0P 1D: ${}^9C_5 [\times {}^6C_0] \times {}^5C_1 = 630$ Condone ${}^5C_1 = 5$ and ${}^6C_1 = 6$	B1	Correct outcome/value for either the 1st or 2nd scenario clearly identified, accept un-simplified, WWW.
	M1 2 correct outcomes/values obtained, accept un-simplified		
	M1	Sum of 3 correct scenarios, may be identified by un-simplified expression.	
	Total: 10710	A1	CAO If one or both M marks not awarded, SCB1 for 10710 WWW.
		4	
5(b)	Ways of selecting 1 st band: ${}^9C_3 \times {}^6C_1 \times {}^5C_1 = 2520$ Ways of selecting 2 nd band: ${}^6C_3 \times {}^5C_1 \times {}^4C_1 = 400$ Ways of selecting 3 rd band: $1 \times {}^4C_1 \times {}^3C_1 = 12$ [Total number of ways =] $2520 \times 400 \times 12 =$	M1	${}^9C_3 \times {}^6C_1 \times {}^5C_1$ or ${}^9C_3 \times 6 \times 5$ seen, condone $\times 3$ or $\times 3!$.
	M1 <i>their</i> $2520 \times$ <i>their</i> $400 \times$ <i>their</i> 12 seen, accept un-simplified, condone $\times 3$ or $\times 3!$.		
	12096000	A1	Condone 12100000. If one or both M marks not awarded, SCB1 for 12096000 (CAO) WWW.
		3	

Question	Answer	Marks	Guidance
6(a)	Method 1		
	$P(X=2) = \frac{6}{10} \times \frac{5}{9} \times \frac{4}{8} \times \frac{3}{7} \times \frac{4!}{2!2!} = \frac{3}{7}$	M1	AG. $\frac{6}{10} \times \frac{5}{9} \times \frac{4}{8} \times \frac{3}{7} \times k \text{ or } \frac{4}{10} \times \frac{3}{9} \times \frac{6}{8} \times \frac{5}{7} \times k \text{ for } k \text{ an integer, } k > 1.$
		A1	4C_2 may be seen for $\frac{4!}{2!2!}$.
	Method 2		
	$\frac{{}^6C_2 \times {}^4C_2}{10C4} = \left[\frac{15 \times 6}{210} \right] \frac{3}{7}$	M1	AG. ${}^6C_2 \times {}^4C_2$ seen as the numerator of a fraction. Condone use of permutations if used consistently.
		A1	
		2	

Question	Answer	Marks	Guidance												
6(b)	<table border="1"> <tr> <td>x</td><td>0</td><td>1</td><td>2</td><td>3</td><td>4</td></tr> <tr> <td>$P(X=x)$</td><td>$\frac{1}{210}$, 0.00476</td><td>$\frac{4}{35}$, 0.114</td><td>$\frac{3}{7}$</td><td>$\frac{8}{21}$, 0.381</td><td>$\frac{1}{14}$, 0.0714</td></tr> </table>	x	0	1	2	3	4	$P(X=x)$	$\frac{1}{210}$, 0.00476	$\frac{4}{35}$, 0.114	$\frac{3}{7}$	$\frac{8}{21}$, 0.381	$\frac{1}{14}$, 0.0714	B1	Table with correct values of x and at least one further non-zero probability correct. Condone extra x values if probability stated as 0.
x	0	1	2	3	4										
$P(X=x)$	$\frac{1}{210}$, 0.00476	$\frac{4}{35}$, 0.114	$\frac{3}{7}$	$\frac{8}{21}$, 0.381	$\frac{1}{14}$, 0.0714										
		B1	Third probability correct. Accept probabilities not in table if clearly identified.												
		B1	Fourth probability correct. Accept probabilities not in table if clearly identified.												
		B1	Fifth probability correct. Accept probabilities not in table if clearly identified. SCB1 for 4 further non-zero probabilities adding to $\frac{4}{7}$, 0.5712 if B2 max scored.												
		4													
6(c)	$[P(2B, 3B 3B1R or 2B2R or 1B 3R) =] \frac{\frac{3}{7} + \frac{8}{21}}{\frac{4}{35} + \frac{3}{7} + \frac{8}{21}}$ $= \left[\frac{17}{97} \right] = \frac{170}{194}, \frac{85}{97}, 0.876$	M1 B1 FT	$\frac{3}{7} + \text{their } \frac{8}{21}$ seen as the numerator of a fraction. $\text{their } \frac{4}{35} + \frac{3}{7} + \text{their } \frac{8}{21}$ seen as the denominator of a fraction.												
		A1	Accept 0.87628...to at least 3SF.												
		3													