



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS**9709/45**

Paper 4 Mechanics

May/June 2025**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 m s^{-2} .

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.





- Find the work done by the pulling force.


[4]

[illegible]



2

4



A machine for driving a nail into a block of wood causes a hammerhead to drop vertically onto the top of the nail. The mass of the hammerhead is 1.5 kg and the mass of the nail is 0.005 kg (see diagram). The hammerhead hits the nail with speed 32 m s^{-1} and remains in contact with the nail after the impact.

- (a) Calculate the speed with which the combined hammerhead and nail move immediately after the impact. Give your answer correct to 3 decimal places. [2]

.....

.....

.....

.....

.....

.....

There is a constant force resisting the motion of magnitude 25 000 N.

- (b) Calculate the distance the nail is driven into the wood. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





- Find the value of T and hence find the distance moved by the train while travelling at the constant speed of 30 ms^{-1} . [5]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.



(b) Find the value of P . [2]

This image shows a full page of white paper with horizontal dashed lines, typical of primary-ruled notebook paper. The lines are evenly spaced and run across the entire width of the page. There are no margins, text, or other markings present.

- (a)** It is given that there is a constant resistance to motion. The engine of the car is working at 24 kW while the car is travelling at a constant speed of 32 m s^{-1} . The power is now increased to 28 kW.

Find the acceleration of the car at the instant it is travelling at a speed of 36 ms^{-1} . [4]

[illegible]



- When the engine is working at 20kW, the car is travelling at constant speed.

Find this constant speed.

[3]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.



- 6 A particle P moves in a straight line starting from a point O . At time t s after leaving O , the velocity, $v \text{ m s}^{-1}$, of P is given by $v = (15 - 2t)^2$.

(a) Find the values of t when the velocity of P is 100 m s^{-1} . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Show that there is a particular value of t for which the velocity and acceleration of the particle are both zero. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....



[illegible]



(a) Given that the system is in equilibrium, find the set of possible values of m . [7]

[illegible]



Use an energy method to find the speed of the particles when each particle has moved 0.6 m. You may assume that this occurs before A reaches the pulley or B reaches the ground. [5]

[illegible]

[illegible]





Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

