

**MARK SCHEME for the October/November 2009 question paper
for the guidance of teachers**

9709/12	9709 MATHEMATICS Paper 12, maximum raw mark 75
----------------	--

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	12

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	12

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through ✓" marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	12

<p>1 $\frac{dy}{dx} = \frac{3}{\sqrt{x}} - x$</p> <p>$(y) = 6\sqrt{x} - \frac{x^2}{2} (+c)$</p> <p>(4, 6) fits $6 = 12 - 8 + c$ $\rightarrow c = 2$</p>	<p>B1, B1</p> <p>M1</p> <p>A1</p> <p>[4]</p>	<p>B1 for each term</p> <p>Uses (4, 6) in an integration with +c co</p>
<p>2 $(x + k)^8$</p> <p>(i) $k^8 + 8k^7x + 28k^6x^2 + 56k^5x^3$</p> <p>(ii) $28k^6 = 56k^5$ $\rightarrow k = 2$</p>	<p>B3, 2, 1</p> <p>[3]</p> <p>M1 A1</p> <p>[2]</p>	<p>Loses 1 for each error. He can gain these marks if appropriate in (ii).</p> <p>Correct method of solving. co. nb $k = 2x$ gets M1 A0.</p>
<p>3 (i) $a + d = 96$ and $a + 3d = 54$ $\rightarrow d = -21$ $a = 117$</p> <p>(ii) $ar = 96$ and $ar^3 = 54$ $\rightarrow r^2 = \frac{54}{96} \rightarrow r = \frac{3}{4}$ $\rightarrow a = 128$</p>	<p>B1</p> <p>M1A1</p> <p>[3]</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>For both expressions. Correct method of solution. co (nb no working, d correct, a wrong 0/3)</p> <p>For both expressions. Correct method of solution. co. $r = \pm\frac{3}{4}$, no penalty.</p>
<p>4 (i) $2 \leq f(x) \leq 8$</p> <p>(ii) $x \mapsto 5 - 3\sin 2x$</p> <p>(iii) No inverse – not 1 : 1.</p>	<p>B1, B1</p> <p>[2]</p> <p>B1</p> <p>DB1</p> <p>B1</p> <p>[3]</p> <p>B1</p> <p>[1]</p>	<p>B1 for 2, B1 for 8. Must be stated, not on graph.</p> <p>1 complete oscillation 1st quadrant, not touching x-axis. Needs to be “down” first and curves. nb If no labels, assume 0 to π.</p> <p>co. Independent of graph.</p>

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	12

<p>5</p> <p>(i) $(\sin x + \cos x)(1 - \sin x \cos x)$ $= \sin x + \cos x - \sin^2 x \cos x - \cos^2 x \sin x$ $\sin^2 x = 1 - \cos^2 x$ and $\cos^2 x = 1 - \sin^2 x$ $\rightarrow \sin^3 x + \cos^3 x$</p> <p>(ii) $(\sin x + \cos x)(1 - \sin x \cos x) = 9 \sin^3 x$ Uses part (i) $\rightarrow 8 \sin^3 x = \cos^3 x$ $\rightarrow \tan^3 x = \frac{1}{8} \rightarrow \tan x = \frac{1}{2}$ $\rightarrow x = 26.6^\circ$ and 206.6°</p>	<p>M1 M1 A1 [3]</p> <p>M1 A1 B1√ [3]</p>	<p>Needs 4 terms from the product. Needs to be used once. All ok.</p> <p>Uses $\tan x = \sin x \div \cos x \rightarrow \tan^3 x = k$. Co. $\sqrt{\quad}$ for $180^\circ +$ first answer and providing there are no other answers in range.</p>
<p>6</p> <p>(i) $\vec{OQ} = 3\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}$ $\vec{PQ} = -3\mathbf{i} + \mathbf{j} + 6\mathbf{k}$</p> <p>(ii) $(3\mathbf{i} + 3\mathbf{j} + 6\mathbf{k}) \cdot (-3\mathbf{i} + \mathbf{j} + 6\mathbf{k})$ $= -9 + 3 + 36 = 30$ $30 = \sqrt{54}\sqrt{46}\cos\theta$ $\theta = 53.0^\circ$</p> <p>Cosine rule M1 modulus M1 attempt at 3 sides M1 A1 answer.</p>	<p>B1 B2, 1 [3]</p> <p>M1 M1 M1A1 [4]</p>	<p>co Loses one for each error.</p> <p>Use of $x_1x_2 + y_1y_2 + z_1z_2$ co. Correct method for modulus (once) and all correctly linked. co.</p> <p>nb $\vec{OQ} \cdot \vec{QP}$ can gain 4/4. but $\vec{OQ} \cdot \vec{PO}$ can only gain 3/4. Use of other vectors (e.g. $\vec{OP} \cdot \vec{OQ}$) M3 ok.</p>
<p>7</p> <p>(i) $2r + r\theta = 50$ $\theta = \frac{1}{r}(50 - 2r)$ $A = \frac{1}{2}r^2\theta$ $\rightarrow A = 25r - r^2$</p> <p>(ii) $\frac{dA}{dr} = 25 - 2r$ $= 0$ when $r = 12.5$ $A = 156\frac{1}{4}$</p> <p>2^{nd} differential negative \rightarrow Maximum</p>	<p>M1 A1 M1 A1 [4]</p> <p>B1 M1 A1 B1 [4]</p>	<p>Must use $s = r\theta$ and link with perimeter co Used with θ as $f(r)$ co (answer given)</p> <p>co sets differential to 0 + solution co Could be quoted directly from quadratic.</p>

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	12

<p>8 $x \mapsto \frac{3}{2x+5}$</p> <p>(i) $f'(x) = -3(2x+5)^{-2} \times 2$ $f'(x)$ is negative \rightarrow decreasing</p> <p>(ii) $y = \frac{3}{2x+5} \rightarrow 2x+5 = \frac{3}{y}$ $\rightarrow f^{-1}(x) = \frac{1}{2}\left(\frac{3}{x} - 5\right)$ or $\frac{3-5x}{2x}$</p> <p>(iii) $\int \pi \frac{9}{(2x+5)^2} dx$ $= (-9\pi(2x+5)^{-1} \div 2)$ Limits 0 to 2 $\rightarrow \pi(-\frac{1}{2} - -0.9)$ $\rightarrow = 0.4\pi$ (or 1.26)</p>	<p>B1 B1 B1 $\sqrt{\quad}$ [3]</p> <p>M1</p> <p>A1 [2]</p> <p>B1 B1 M1 A1 [4]</p>	<p>B1 for $-3(2x+5)^{-2}$. B1 for $\times 2$ $\sqrt{\quad}$ providing bracket is squared. (using value or values only B0)</p> <p>Attempt at making x the subject.</p> <p>co including $f(x)$ not $f(y)$</p> <p>For $-9(2x+5)^{-1}$ For $\div 2$ in \int of y^2 Use of correct limits with \int of y^2. co</p>
<p>9 (i) y-coordinate same as the y-coordinate of the mid-point of AC.</p> <p>(ii) m of $AD = \frac{8}{h}$ or $\frac{h-12}{8}$ m of $CD = \frac{8}{12-h}$ or $\frac{-h}{8}$</p> <p>nb $AC = 20$, $M(6, 6)$ $MD = 10 \rightarrow$ $D(16, 6)$ and $B(-4, 6)$</p> <p>(iii) Product of gradients = -1 $\rightarrow h^2 - 12h - 64 = 0$ $\rightarrow h = 16$ or -4 so $x_D = 16$ and $x_B = -4$ or Pyth $h^2 + 8^2 + 8^2 + (12-h)^2 = 400$</p> <p>(iv) Area = $\sqrt{320} \times \sqrt{80}$ $\rightarrow 160$</p> <p>(or Area = $2 \times$ area of a triangle with base = BD, $\rightarrow 2 \times \frac{1}{2} \times 20 \times 8$ $= 160$) (or matrix method)</p>	<p>B1 [1]</p> <p>M1 A1</p> <p>A1 [3]</p> <p>M1 M1 DM1A1 [4]</p> <p>M1 M1 A1 [3]</p>	<p>co</p> <p>any use of y-step \div x-step for M mark</p> <p>co</p> <p>Used correctly with the two gradients Forming a quadratic equation Solution of equation. co</p> <p>M1 for method for one of the lengths M1 for base \times height. co</p>

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2009	9709	12

<p>10</p> <p>(i) (a) $2y = x + 5, y = x^2 - 4x + 7$ Sim equations $\rightarrow 2x^2 - 9x + 9 = 0$ $\rightarrow x = 3$ or $x = 1\frac{1}{2}$.</p> <p>(b) $\frac{dy}{dx} = 2x - 4$ $\rightarrow y - 4 = 2(x - 3)$</p> <p>nb use of $y + 4$ or x, y interchanged M1 A0</p> <p>(c) $m = 2 \rightarrow$ angle of 63.4° $m = \frac{1}{2} \rightarrow$ angle of 26.6° \rightarrow angle between = 37°</p> <p>(i+2j).(2i+j) $\rightarrow 4 = \sqrt{5}\sqrt{5}\cos\theta$ M1M1A1 or use of $\tan(A-B)$ M2A1 or Cosine rule with 3 sides found.</p> <p>(ii) $y = x^2 - 4x + 7$ $2y = x + k$ Sim eqns $\rightarrow 2x^2 - 9x + 14 - k = 0$ Uses $b^2 - 4ac, 81 - 8(14 - k)$ Key value is $k = 3.875$ or $31/8$. $k < 3.875$</p>	<p>M1 DM1 A1 [3]</p> <p>B1 M1 A1 [3]</p> <p>M1 M1A1 [3]</p> <p>M1 A1 M1 A1 [4]</p>	<p>Complete elimination of x or y Correct method for quadratic. co.</p> <p>co</p> <p>Correct form of eqn with m numeric. co</p> <p>Finds angle with x-axis once. Subtracts two angles. co.</p> <p>Eliminates y or x completely. Co (= 0) Uses $b^2 - 4ac = 0, \text{ or } < 0 \text{ or } > 0$</p> <p>Co condone \leq.</p>
---	---	---