
MATHEMATICS

9709/62

Paper 6

October/November 2019

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **12** printed pages.

PUBLISHED**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

PUBLISHED**Mark Scheme Notes**

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B** Mark for a correct result or statement independent of method marks.
- DM or DB** When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- FT** Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.

Abbreviations

AEF/OE	Any Equivalent Form (of answer is equally acceptable) / Or Equivalent
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
CWO	Correct Working Only
ISW	Ignore Subsequent Working
SOI	Seen Or Implied
SC	Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)
WWW	Without Wrong Working
AWRT	Answer Which Rounds To
NFWW	Not From Wrong Working

PUBLISHED

Question	Answer	Marks	Guidance
1(i)	Median = 51 UQ = 57.5, LQ = 40	B1	
	IQR = UQ – LQ	M1	$55 \leq \text{UQ} \leq 62 - 38 \leq \text{LQ} \leq 45$
	17.5	A1	NFWW
		3	
1(ii)	Result will be disproportionately affected by 110	B1	Affected by an extreme/large value There is a large outlier ...contains outliers such as 110... Not 'mean affected by extreme values'
		1	

Question	Answer	Marks	Guidance
2(i)	$0.4x + 0.6 \times 2x = 0.36$ or $0.4(1 - x) + 0.6(1 - 2x) = 0.64$	M1	$0.4a + (1 - 0.4)b = 0.36$ or 0.64 , a, b terms involving x
	$1.6x = 0.36$ $x = 0.225$	A1	Fully justified by algebra AG
		2	

PUBLISHED

Question	Answer	Marks	Guidance
2(ii)	$P(H L) = \frac{0.4(1-x)}{1-0.36} = \frac{0.4 \times (1-0.225)}{0.64} = \frac{0.4 \times 0.775}{0.4 \times 0.775 + 0.6 \times 0.55}$	M1	Correct numerical numerator of a fraction. Allow unsimplified.
		M1	Denominator 0.36 or 0.64. Allow unsimplified.
	$\frac{31}{64}$ or 0.484	A1	
		3	

Question	Answer	Marks	Guidance
3(i)	0.5 2.4 3 1.4 0.4	M1	At least 3 frequency densities calculated (frequency \div class width) e.g. $\left(\frac{10}{20}, \frac{10}{19} \text{ or } \frac{10}{19.5}\right)$ may be read from graph using <i>their</i> scale, 3SF or exact
	All heights correct on graph.	A1	
	Bar ends of 9.5, 29.5, 39.5, 59.5, 89.5	B1	
	Axes labelled: Frequency density (fd) and speed/km h ⁻¹ (or appropriate title). Linear scales $9.5 \leq \text{horizontal axis} \leq 89.5$, $0 \leq \text{vertical axis} \leq 3$, 5 bars with no gaps	B1	
		4	

PUBLISHED

Question	Answer	Marks	Guidance
3(ii)	$\frac{19.5 \times 10 + 34.5 \times 24 + 44.5 \times 30 + 54.5 \times 14 + 74.5 \times 12}{90}$ <p style="text-align: center;"><i>their 90</i></p> $= \frac{195 + 828 + 1335 + 763 + 894}{90}$ $= \frac{4015}{90} \text{ or } \frac{803}{18}$	M1	Uses at least 4 midpoint attempts (e.g. 19.5 ± 0.5). Allow unsimplified expression.
	$44 \frac{11}{18}$ or 44.6 (km h ⁻¹)	A1	Final answer not an improper fraction NFWW
		2	

Question	Answer	Marks	Guidance
4(i)	$P(8, 9, 10) = {}^{10}C_8 0.66^8 0.34^2 + {}^{10}C_9 0.66^9 0.34^1 + 0.66^{10}$	M1	Correct binomial term, ${}^{10}C_a 0.66^a (1-0.66)^b$ $a+b = 10, 0 < a, b < 10$
		A1	Correct unsimplified expression
	0.284	B1	CAO
		3	

PUBLISHED

Question	Answer	Marks	Guidance
4(ii)	$np = 0.66 \times 150 = 99$ $npq = 0.66 \times (1 - 0.66) \times 150 = 33.66$	B1	Accept evaluated or unsimplified μ, σ^2 numerical expressions, condone $\sigma = \sqrt{33.66} = 5.8017$ or 5.802 CAO
	$P(X > 84) = P\left(Z > \frac{84.5 - 99}{\sqrt{33.66}}\right)$	M1	\pm Standardise, $\frac{x - \text{their } 99}{\sqrt{\text{their } 33.66}}$, condone σ^2, x a value
		M1	84.5 or 83.5 used in <i>their</i> standardisation formula
	$(= P(Z > -2.499))$	M1	Correct final area
	0.994	A1	Final answer (accept 0.9938) SC if no standardisation formula seen, B2 $P(Z > -2.499) = 0.994$
		5	

PUBLISHED

Question	Answer	Marks	Guidance														
5(i)	<table border="1"> <tr> <td>x</td> <td>-1</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> </tr> <tr> <td>p</td> <td>$\frac{1}{12}$</td> <td>$\frac{1}{12}$</td> <td>$\frac{3}{12}$</td> <td>$\frac{2}{12}$</td> <td>$\frac{3}{12}$</td> <td>$\frac{2}{12}$</td> </tr> </table>	x	-1	0	1	2	3	4	p	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{3}{12}$	$\frac{2}{12}$	$\frac{3}{12}$	$\frac{2}{12}$	B1	Table with correct values of x , at least 1 probability, all probabilities ≤ 1
	x	-1	0	1	2	3	4										
	p	$\frac{1}{12}$	$\frac{1}{12}$	$\frac{3}{12}$	$\frac{2}{12}$	$\frac{3}{12}$	$\frac{2}{12}$										
			B1	2 probabilities correct, may not be in table													
			B1	2 more probabilities correct, may not be in table													
		B1	All correct, values in table SC1 No more than 1 correct probability and at least 5 probabilities summing to 1 in table														
		4															
5(ii)	$[E(X)] = \left(\frac{-1+0+3+4+9+8}{12} \right) = \frac{23}{12}$	M1	May be implied by use in variance. Allow unsimplified expression														
	$[\text{Var}(X)] = \frac{1+0+3+8+27+32(=71)}{12} - \left(\frac{23}{12} \right)^2$	M1	Appropriate variance formula using <i>their</i> $E(X)^2$														
	2.24 or $\frac{323}{144}$ or $2\frac{35}{144}$	A1	CAO														
		3															

PUBLISHED

Question	Answer	Marks	Guidance
6(i)	$P(X < 45) = P\left(Z < \frac{45 - 40}{8}\right)$ $= P(Z < 0.625)$	M1	\pm Standardise, no continuity correction, σ^2 or $\sqrt{\sigma}$, formula must be seen
	0.734(0)	A1	CAO
		2	
6(ii)	$1 - 2(1 - (i)) = 2(i) - 1 = 2((i) - 0.5)$	M1	Use result of part (i) or recalculated to find area OE
	0.468	A1ft	$0 < \text{FT from (i)} < 1$ or correct.
		2	
6(iii)	$P(X < 10) = 48/500 = 0.096$ $z = -1.305$	B1	$z = \pm 1.305$
	$P(X > 24) = 76/500 = 0.152$ $z = 1.028$	B1	$z = \pm 1.028$
	$10 - \mu = -1.305\sigma$ $24 - \mu = 1.028\sigma$	M1	Form 1 equation using 10 or 24 with μ, σ, z -value. Allow continuity correction, not $\sigma^2, \sqrt{\sigma}$
	$14 = 2.333\sigma$	M1	OE Solve two equations in σ and μ to form equation in one variable
	$\sigma = 6.[00], \mu = 17.8[3]$	A1	CAO, WWW
		5	

PUBLISHED

Question	Answer	Marks	Guidance
7(i)	6! = 720	B1	Evaluated
		1	
7(ii)	Total no of arrangements: $\frac{9!}{2!3!} = 30240$	B1	Accept unevaluated
	No with Ts together = $\frac{8!}{3!} = 6720$	B1	Accept unevaluated
	With Ts not together: 30 240 – 6720	M1	correct or $\frac{9!}{m} - \frac{8!}{n}$, m, n integers > 1 or <i>their</i> identified total – <i>their</i> identified Ts together
	23 520	A1	CAO
	Alternative method for question 7(ii)		
	$\frac{7!}{3!} \times \frac{8 \times 7}{2}$	B1	$7! \times (k > 0)$ in numerator, cannot be implied by 7P_2 , etc.
		B1	$3! \times (k > 0)$ in denominator
		M1	$\frac{\textit{their } 7!}{\textit{their } 3!} \times {}^8C_2$ or 8P_2
	23 520	A1	CAO
		4	

PUBLISHED

Question	Answer	Marks	Guidance
7(iii)	Number of arrangements = $\frac{7!}{3!}$ Probability = $\frac{\text{their } \frac{7!}{3!}}{\text{their } \frac{9!}{3!2!}} = \frac{840}{30240}$	M1	$\frac{\text{their identified number of arrangements with T at ends}}{\text{their identified total number of arrangements}}$ $\frac{7!}{9!}$ or $\frac{m}{n}$, m, n integers > 1
	$\frac{1}{36}$ or 0.0278	A1	Final answer
		2	
7(iv)	OOT__ ${}^4C_2 = 6$ OOTT_ ${}^4C_1 = 4$ OOOT_ ${}^4C_1 = 4$ OOOT_ $= 1$	M1	4C_x seen alone or ${}^4C_x \times k \geq 1$, k an integer, $0 < x < 4$
		A1	${}^4C_2 \times k$, $k = 1$ oe or ${}^4C_1 \times m$, $m = 1$ oe alone
		M1	Add 3 or 4 identified correct scenarios only, accept unsimplified
	(Total) = 15	A1	CAO, WWW Only dependent on 2nd M mark
		4	