



## Cambridge International AS & A Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**MATHEMATICS**

**9709/41**

Paper 4 Mechanics

**October/November 2022**

**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity ( $g$ ) is needed, use  $10 \text{ m s}^{-2}$ .

### INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **12** pages.





3 A constant resistance of magnitude 1400 N acts on a car of mass 1250 kg.

(a) The car is moving along a straight level road at a constant speed of  $28 \text{ m s}^{-1}$ .

Find, in kW, the rate at which the engine of the car is working. [2]

.....  
.....  
.....  
.....  
.....  
.....

(b) The car now travels at a constant speed up a hill inclined at an angle of  $\theta$  to the horizontal, where  $\sin \theta = 0.12$ , with the engine working at 43.5 kW.

Find this speed. [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....





.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The block is initially at rest.

Find the distance travelled by the block during the fourth second of motion. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 A particle  $P$  moves on the  $x$ -axis from the origin  $O$  with an initial velocity of  $-20 \text{ m s}^{-1}$ . The acceleration  $a \text{ m s}^{-2}$  at time  $t \text{ s}$  after leaving  $O$  is given by  $a = 12 - 2t$ .

(a) Sketch a velocity-time graph for  $0 \leq t \leq 12$ , indicating the times when  $P$  is at rest. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....







(b)




Fig. 6.2

It is given instead that the plane is smooth and the particles are released from rest when the difference in the vertical heights of the particles is 1 m (see Fig. 6.2).

Use an energy method to find the speed of the particles at the instant when the particles are at the same horizontal level. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

