



Cambridge International AS & A Level

CANDIDATE
NAME



CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

October/November 2024

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.



1 Expand $(9-3x)^{\frac{1}{2}}$ in ascending powers of x , up to and including the term in x^2 , simplifying the coefficients. [4]





2 (a) By sketching a suitable pair of graphs, show that the equation $\cot 2x = \sec x$ has exactly one root in the interval $0 < x < \frac{1}{2}\pi$. [2]

(b) Show that if a sequence of real values given by the iterative formula

$$x_{n+1} = \frac{1}{2} \tan^{-1}(\cos x_n)$$

converges, then it converges to the root in part (a).

[1]





3 The square roots of $6 - 8i$ can be expressed in the Cartesian form $x + iy$, where x and y are real and exact.

By first forming a quartic equation in x or y , find the square roots of $6-8i$ in exact Cartesian form. [5]

[5]





4 Solve the equation $5^x = 5^{x+2} - 10$. Give your answer correct to 3 decimal places. [3]





5 (a) The complex number u is given by

$$u = \frac{(\cos \frac{1}{7}\pi + i \sin \frac{1}{7}\pi)^4}{\cos \frac{1}{7}\pi - i \sin \frac{1}{7}\pi}.$$

Find the exact value of $\arg u$.


[2]

The complex numbers u and u^* are plotted on an Argand diagram.

Describe the single geometrical transformation that maps u onto u^* and state the exact value of $\arg u^*$. [2]

DO NOT WRITE IN THIS MARGIN





The variables x and y satisfy the equation $ay = b^x$, where a and b are constants. The graph of $\ln y$ against x is a straight line passing through the points $(0.50, 2.24)$ and $(3.40, 8.27)$, as shown in the diagram.

Find the values of a and b . Give each value correct to 1 significant figure.

[4]





7 (a) Show that the equation $\tan^3 x + 2 \tan 2x - \tan x = 0$ may be expressed as

$$\tan^4 x - 2 \tan^2 x - 3 = 0$$

for $\tan x \neq 0$.

[3]





(b) Hence solve the equation $\tan^3 2\theta + 2 \tan 4\theta - \tan 2\theta = 0$ for $0 < \theta < \pi$. Give your answers in exact form. [3]





8 The parametric equations of a curve are

$$x = \tan^2 2t, \quad y = \cos 2t,$$

for $0 < t < \frac{1}{4}\pi$.

(a) Show that $\frac{dy}{dx} = -\frac{1}{2}\cos^3 2t$.

[4]

DO NOT WRITE IN THIS MARGIN





(b) Hence find the equation of the normal to the curve at the point where $t = \frac{1}{8}\pi$. Give your answer in the form $y = mx + c$. [4]





9 With respect to the origin O , the points A , B and C have position vectors given by

$$\overrightarrow{OA} = \begin{pmatrix} 2 \\ 1 \\ -3 \end{pmatrix}, \quad \overrightarrow{OB} = \begin{pmatrix} 0 \\ 4 \\ 1 \end{pmatrix} \quad \text{and} \quad \overrightarrow{OC} = \begin{pmatrix} -3 \\ -2 \\ 2 \end{pmatrix}.$$

(a) The point D is such that $ABCD$ is a trapezium with $\overrightarrow{DC} = 3\overrightarrow{AB}$.

Find the position vector of D .

[2]

(b) The diagonals of the trapezium intersect at the point P .

Find the position vector of P .

[5]





1

DO NOT WRITE IN THIS MARGIN

(c) Using a scalar product, calculate angle ABC . [4]





10 A balloon in the shape of a sphere has volume V and radius r . Air is pumped into the balloon at a constant rate of 40π starting when time $t = 0$ and $r = 0$. At the same time, air begins to flow out of the balloon at a rate of $0.8\pi r$. The balloon remains a sphere at all times.

(a) Show that r and t satisfy the differential equation

$$\frac{dr}{dt} = \frac{50-r}{5r^2}. \quad [3]$$

(b) Find the quotient and remainder when $5r^2$ is divided by $50 - r$.

[3]





(c) Solve the differential equation in part (a), obtaining an expression for t in terms of r .

[6]

(d) Find the value of t when the radius of the balloon is 12.

[1]

.....
.....
.....





11 Let $f(x) = \frac{2e^{2x}}{e^{2x} - 3e^x + 2}$.

(a) Find $f'(x)$ and hence find the exact coordinates of the stationary point of the curve with equation $y = f(x)$. [5]





(b) Use the substitution $u = e^x$ and partial fractions to find the exact value of $\int_{\ln 3}^{\ln 5} f(x) dx$.

Give your answer in the form $\ln a$, where a is a rational number in its simplest form.

[9]





Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.



* 0000800000019 *



19

BLANK PAGE



DO NOT WRITE IN THIS MARGIN





Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

