



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS**9709/63**

Paper 6 Probability & Statistics 2

October/November 2024**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages. Any blank pages are indicated.





- 1 The heights of a certain species of deer are known to have standard deviation 0.35 m. A zoologist takes a random sample of 150 of these deer and finds that the mean height of the deer in the sample is 1.42 m.

(a) Calculate a 96% confidence interval for the population mean height. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Bubay says that 96% of deer of this species are likely to have heights that are within this confidence interval.

Explain briefly whether Bubay is correct. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



[illegible]



- 3 The times, T minutes, taken by a random sample of 75 students to complete a test were noted. The results were summarised by $\sum t = 230$ and $\sum t^2 = 930$.

(a) Calculate unbiased estimates of the population mean and variance of T . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

You should now assume that your estimates from part (a) are the true values of the population mean and variance of T .

- (b) The times taken by another random sample of 75 students were noted, and the sample mean, \bar{T} , was found.

Find the value of a such that $P(\bar{T} > a) = 0.234$. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



$$f(x) = \begin{cases} \frac{a}{x^2} - \frac{18}{x^3} & 2 \leq x \leq 3, \\ 0 & \text{otherwise,} \end{cases}$$

(a) Show that $a = \frac{27}{2}$. [3]

(b) Show that $E(X) = \frac{27}{2} \ln \frac{3}{2} - 3$. [3]



-
-
-

-
- This image shows a full page of white paper with horizontal blue or grey ruling lines. The lines are evenly spaced and run across the width of the page, providing a template for handwriting practice. There are no margins, text, or other markings on the page.



6 The numbers of customers arriving at service desks A and B during a 10-minute period have the independent distributions $Po(1.8)$ and $Po(2.1)$ respectively.

(a) Find the probability that during a randomly chosen 15-minute period more than 2 customers will arrive at desk A . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the probability that during a randomly chosen 5-minute period the total number of customers arriving at both desks is less than 4. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





- [illegible]



- 7 The number of accidents per year on a certain road has the distribution $Po(\lambda)$. In the past the value of λ was 3.3 . Recently, a new speed limit was imposed and the council wishes to test whether the value of λ has decreased. The council notes the total number, X , of accidents during **two** randomly chosen years after the speed limit was introduced and it carries out a test at the 5% significance level.

(a) Calculate the probability of a Type I error. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Given that $X = 2$, carry out the test. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....





- (c) The council decides to carry out another similar test at the 5% significance level using the same hypotheses and two different randomly chosen years.

Given that the true value of λ is 0.6, calculate the probability of a Type II error. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (d) Using $\lambda = 0.6$ and a suitable approximating distribution, find the probability that there will be more than 10 accidents in 30 years. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN