Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

6146662382

FURTHER MATHEMATICS **9231/12**

Paper 1 Further Pure Mathematics 1 **October/November 2022**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **20** pages. Any blank pages are indicated.

[Turn over

1 **(a)** Use the list of formulae (MF19) to find $\sum_{r=1}^{n} r(r+2)$ in terms of n, simplifying your answer. [2]

(b) Express $\frac{1}{r(r+2)}$ in partial fractions and hence find $\sum_{r=1}^{n} \frac{1}{r(r+2)}$ in terms of n. [4]

(c) Deduce the value of $\sum_{r=1}^{\infty} \frac{1}{r(r+2)}$. [1]

2 The equation $x^4 + 3x^2 + 2x + 6 = 0$ has roots $\alpha, \beta, \gamma, \delta$.

(a) Find a quartic equation whose roots are $\frac{1}{\alpha^2}, \frac{1}{\beta^2}, \frac{1}{\gamma^2}, \frac{1}{\delta^2}$ and state the value of $\frac{1}{\alpha^2} + \frac{1}{\beta^2} + \frac{1}{\gamma^2} + \frac{1}{\delta^2}$. [4]

(b) Find the value of $\beta^2\gamma^2\delta^2 + \alpha^2\gamma^2\delta^2 + \alpha^2\beta^2\delta^2 + \alpha^2\beta^2\gamma^2$. [3]

(c) Find the value of $\frac{1}{\alpha^4} + \frac{1}{\beta^4} + \frac{1}{\gamma^4} + \frac{1}{\delta^4}$. [2]

3 The matrix **M** is given by $\mathbf{M} = \begin{pmatrix} 1 & 0 \\ 0 & k \end{pmatrix}\begin{pmatrix} 1 & 0 \\ k & 1 \end{pmatrix}$, where k is a constant and $k \neq 0$ or 1.

(a) The matrix **M** represents a sequence of two geometrical transformations.

State the type of each transformation, and make clear the order in which they are applied. [2]

.....

.....

.....

(b) Write $\mathbf{M}^{-1}$ as the product of two matrices, neither of which is **I**. [2]

.....

.....

.....

.....

.....

(c) Show that the invariant points of the transformation represented by **M** lie on the line $y = \frac{k^2}{1-k}x$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) The triangle ABC in the x-y plane is transformed by **M** onto triangle DEF.

Find the value of k for which the area of triangle DEF is equal to the area of triangle ABC. [2]

4 The function f is such that $\mathrm{f}''(x) = \mathrm{f}(x)$.

Prove by mathematical induction that, for every positive integer n,

$$\frac{\mathrm{d}^{2n-1}}{\mathrm{d}x^{2n-1}}\big(x\,\mathrm{f}(x)\big) = x\,\mathrm{f}'(x) + (2n-1)\,\mathrm{f}(x). \qquad [7]$$

5 The curve C has polar equation $r = a\sec^2\theta$, where a is a positive constant and $0 \leqslant \theta \leqslant \frac{1}{4}\pi$.

(a) Sketch C, stating the polar coordinates of the point of intersection of C with the initial line and also with the half-line $\theta = \frac{1}{4}\pi$. [3]

.....

.....

(b) Find the maximum distance of a point of C from the initial line. [2]

.....

.....

.....

.....

(c) Find the area of the region enclosed by C, the initial line and the half-line $\theta = \frac{1}{4}\pi$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Find, in the form $y = \mathrm{f}(x)$, the Cartesian equation of C. [3]

6 The lines l_1 and l_2 have equations $\mathbf{r} = 2\mathbf{i} + \mathbf{k} + \lambda(\mathbf{i} - \mathbf{j} + 2\mathbf{k})$ and $\mathbf{r} = 2\mathbf{j} + 6\mathbf{k} + \mu(\mathbf{i} + 2\mathbf{j} - 2\mathbf{k})$ respectively.

The point P on l_1 and the point Q on l_2 are such that PQ is perpendicular to both l_1 and l_2.

(a) Find the length PQ. [5]

The plane Π_1 contains PQ and l_1.

The plane Π_2 contains PQ and l_2.

(b) (i) Write down an equation of Π_1, giving your answer in the form $\mathbf{r} = \mathbf{a} + s\mathbf{b} + t\mathbf{c}$. [1]

.....

.....

(ii) Find an equation of Π_2, giving your answer in the form $ax + by + cz = d$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find the acute angle between Π_1 and Π_2. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 The curve C has equation $y = \frac{x^2 - x}{x + 1}$.

(a) Find the equations of the asymptotes of C. [3]

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the exact coordinates of the stationary points on C. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Sketch C, stating the coordinates of any intersections with the axes. [3]

.....

(d) Sketch the curve with equation $y = \left|\frac{x^2 - x}{x + 1}\right|$ and find in exact form the set of values of x for which $\left|\frac{x^2 - x}{x + 1}\right| < 6$. [5]

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

BLANK PAGE

BLANK PAGE

BLANK PAGE