Cambridge Assessment International Education

Cambridge International AS & A Level

CANDIDATE NAME

CENTRE NUMBER | | | | | |

CANDIDATE NUMBER | | | | |

0510110257

FURTHER MATHEMATICS **9231/22**

Paper 2 Further Pure Mathematics 2 **October/November 2022**

2 hours

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages.

[Turn over

1 **(a)** Find the set of values of k for which the system of equations

$$\begin{aligned} x + 2y + 3z &= 1, \\ kx + 4y + 6z &= 0, \\ 7x + 8y + 9z &= 3, \end{aligned}$$

has a unique solution. [3]

(b) Interpret the situation geometrically in the case where the system of equations does not have a unique solution. [2]

2 A curve has equation

$$(x+1)y+y^2=2.$$

(a) Show that $\frac{dy}{dx}=-\frac{2}{3}$ at the point $(0,-2)$. [3]

(b) Find the value of $\frac{d^2y}{dx^2}$ at the point $(0,-2)$. [4]

3 **(a)** A curve has equation $y = e^x + \frac{1}{4}e^{-x}$, for $0 \leqslant x \leqslant 1$. Find, in terms of π and e, the area of the surface generated when the curve is rotated through 2π radians about the x-axis. [6]

(b) Using standard results from the list of formulae (MF19), or otherwise, find the Maclaurin's series for $e^{x} + \frac{1}{4}e^{-x}$ up to and including the term in x^2. [2]

4 Find the solution of the differential equation

$$(4t^2 - 1)\frac{\mathrm{d}x}{\mathrm{d}t} + 4x = 4t^2 - 1$$

for which $x = 3$ when $t = 1$. Give your answer in the form $x = \mathrm{f}(t)$. [9]

5 **(a)** Write down the fourth roots of unity. [1]

.....

.....

(b) Use de Moivre's theorem to show that

$$\cos 4\theta = 8\cos^4\theta - 8\cos^2\theta + 1.$$ [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Hence obtain the real roots of the equation

$$16(8x^4 - 8x^2 + 1)^4 - 9 = 0$$

in the form $\cos(q\pi)$, where q is a rational number. [5]

6



The diagram shows the curve $y = \dfrac{1}{\sqrt{x^2 + 2x}}$ for $x > 0$, together with a set of $(n-1)$ rectangles of unit width.

By considering the sum of the areas of these rectangles, show that

$$\sum_{r=1}^{n} \frac{1}{\sqrt{r^2 + 2r}} < \ln\left(n + 1 + \sqrt{n^2 + 2n}\right) + \tfrac{1}{3}\sqrt{3} - \ln\left(2 + \sqrt{3}\right). \qquad [10]$$

7 **(a)** It is given that λ is an eigenvalue of the non-singular square matrix **A**, with corresponding eigenvector **e**.

Show that λ^{-1} is an eigenvalue of $\mathbf{A}^{-1}$ for which **e** is a corresponding eigenvector. [2]

.....

.....

.....

.....

The matrix **A** is given by

$$\mathbf{A} = \begin{pmatrix} 2 & 0 & 3 \\ 15 & -4 & 3 \\ 3 & 0 & 2 \end{pmatrix}.$$

(b) Given that -1 is an eigenvalue of **A**, find a corresponding eigenvector. [2]

.....

.....

.....

.....

.....

(c) It is also given that $\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix}$ are eigenvectors of **A**. Find the corresponding eigenvalues. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Hence find a matrix $\mathbf{P}$ and a diagonal matrix $\mathbf{D}$ such that $\mathbf{A}^{-1} = \mathbf{PDP}^{-1}$. [2]

(e) Use the characteristic equation of $\mathbf{A}$ to show that $\mathbf{A}^{-1} = p\mathbf{A}^2 + q\mathbf{I}$, where p and q are rational numbers to be determined. [4]

8 It is given that $y = \cosh u$, where $u > 0$, and

$$\sqrt{\cosh^2 u - 1}\left(\frac{\mathrm{d}^2u}{\mathrm{d}x^2} + \frac{\mathrm{d}u}{\mathrm{d}x}\right) + \cosh u\left(\frac{\mathrm{d}u}{\mathrm{d}x}\right)^2 - 2\cosh u = 4\mathrm{e}^{-x}.$$

(a) Show that

$$\frac{\mathrm{d}^2y}{\mathrm{d}x^2} + \frac{\mathrm{d}y}{\mathrm{d}x} - 2y = 4\mathrm{e}^{-x}. \qquad [4]$$

(b) Find u in terms of x, given that, when $x = 0$, $u = \ln 3$ and $\frac{\mathrm{d}u}{\mathrm{d}x} = 3$. [10]

Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....