

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 3 3 6 6 6 8 4 6 1 9 \*



**CHEMISTRY**

**9701/34**

Paper 3 Advanced Practical Skills 2

**May/June 2018**

**2 hours**

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.  
Give details of the practical session and laboratory where appropriate, in the boxes provided.  
Write in dark blue or black pen.  
You may use an HB pencil for any diagrams or graphs.  
Do not use staples, paper clips, glue or correction fluid.  
**DO NOT WRITE IN ANY BARCODES.**

Answer **all** questions.  
Electronic calculators may be used.  
You may lose marks if you do not show your working or if you do not use appropriate units.  
Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11.  
A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.  
The number of marks is given in brackets [ ] at the end of each question or part question.

<b>Session</b>	
<b>Laboratory</b>	

<b>For Examiner's Use</b>	
<b>1</b>	
<b>2</b>	
<b>Total</b>	

This document consists of **12** printed pages.

## Quantitative Analysis

Read through the whole method before starting any practical work. Where appropriate, prepare a table for your results in the space provided.

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

- 1 Glucose,  $C_6H_{12}O_6$ , is a sugar that can act as a reducing agent. You will investigate how an increase in temperature affects the rate of the redox reaction between glucose and acidified potassium manganate(VII).

**FB 1** is  $0.010 \text{ mol dm}^{-3}$  acidified potassium manganate(VII),  $KMnO_4$ .

**FB 2** is  $1.0 \text{ mol dm}^{-3}$  sulfuric acid,  $H_2SO_4$ .

**FB 3** is an aqueous solution containing  $32.8 \text{ g dm}^{-3}$  glucose,  $C_6H_{12}O_6$ , distilled water

You will measure the time it takes for the purple colour to disappear. Your table of results on page 4 should include the rate of reaction for each experiment.

### (a) Method

#### Experiment 1

- Fill the burette with **FB 1**.
- Add  $10.00 \text{ cm}^3$  of **FB 1** into the  $250 \text{ cm}^3$  beaker.
- Use the  $50 \text{ cm}^3$  measuring cylinder to transfer  $50.0 \text{ cm}^3$  of **FB 2** into the beaker containing **FB 1**.
- Use the same measuring cylinder to transfer  $50.0 \text{ cm}^3$  of distilled water into the same beaker.
- Place the beaker on the tripod and heat its contents to between  $65^\circ\text{C}$  and  $70^\circ\text{C}$ .
- While the solution in the beaker is heating pour  $25.0 \text{ cm}^3$  of **FB 3** into the  $25 \text{ cm}^3$  measuring cylinder.
- When the temperature of the contents of the beaker has reached between  $65^\circ\text{C}$  and  $70^\circ\text{C}$ , remove the Bunsen burner and **carefully** place the hot beaker onto the white tile.
- Record the temperature of the solution in the beaker.
- Add the  $25.0 \text{ cm}^3$  of **FB 3** and **immediately** start timing.
- Stir the contents of the beaker once and stop timing as soon as the solution turns colourless. Record the time to the nearest second.
- Record the temperature of the solution as soon as it is colourless.
- Calculate and record the average temperature of the reaction mixture to one decimal place.
- Empty, rinse and dry the beaker so it is ready for use in **Experiment 2**.

**Experiment 2**

- Add 10.00 cm<sup>3</sup> of **FB 1** into the 250 cm<sup>3</sup> beaker.
- Use the 50 cm<sup>3</sup> measuring cylinder to transfer 50.0 cm<sup>3</sup> of **FB 2** into the beaker containing **FB 1**.
- Use the same measuring cylinder to transfer 50.0 cm<sup>3</sup> of distilled water into the same beaker.
- Place the beaker on the tripod and heat its contents to between 30 °C and 35 °C.
- While the solution in the beaker is heating pour 25.0 cm<sup>3</sup> of **FB 3** into the 25 cm<sup>3</sup> measuring cylinder.
- When the temperature of the contents of the beaker has reached between 30 °C and 35 °C, remove the Bunsen burner and **carefully** place the hot beaker onto the white tile.
- Record the temperature of the solution in the beaker.
- Add the 25.0 cm<sup>3</sup> of **FB 3** and **immediately** start timing.
- Stir the contents of the beaker once and stop timing as soon as the solution turns colourless. Record the time to the nearest second.
- Record the temperature of the solution as soon as it is colourless.
- Calculate and record the average temperature of the reaction mixture to one decimal place.
- Empty, rinse and dry the beaker so it is ready for use in **Experiment 3**.

**Experiments 3, 4 and 5**

- Repeat the method for **Experiment 2** at three different temperatures.
- Keep the temperature of the contents of the beaker between room temperature and 70 °C.
- Record all your results in your table.

## Results

The rate of reaction can be calculated as shown.

$$\text{rate} = \frac{1000}{\text{reaction time}}$$

Calculate the rate of reaction for each experiment and include this in your table.

I	
II	
III	
IV	
V	
VI	
VII	
VIII	

[8]

- (b) Plot a graph of rate ( $y$ -axis) against average temperature ( $x$ -axis) on the grid opposite. Select a scale on the  $x$ -axis to include an average temperature of  $15.0^{\circ}\text{C}$ . Label any points you consider anomalous.

Draw a line of best fit and extrapolate it to  $15.0^{\circ}\text{C}$ .


[4]

- (c) Use your graph to calculate the **time** to the nearest second that the reaction would have taken if the average temperature had been  $52.5^{\circ}\text{C}$ . Show **on the grid** how you obtained your answer.

time = ..... s [2]

- (d) Explain, by referring to your graph or your table of results, how the rate of reaction is affected by an increase in temperature.

.....  
 .....  
 ..... [2]



I	
II	
III	
IV	

- (e) (i) Calculate the concentration of glucose in **FB 3** in  $\text{mol dm}^{-3}$ .

concentration of glucose in **FB 3** = .....  $\text{mol dm}^{-3}$  [1]

- (ii) Under certain conditions, 1.0 mole of acidified potassium manganate(VII),  $\text{KMnO}_4$ , can oxidise 2.5 moles of glucose.

Calculate the volume of  $0.010 \text{ mol dm}^{-3}$  acidified  $\text{KMnO}_4$  that would react with **all** the glucose present in  $25.0 \text{ cm}^3$  of **FB 3**.

[3]

- (iii) The formula of glucose can be written as  $\text{CHO}(\text{CHOH})_4\text{CH}_2\text{OH}$ .

Suggest the formula of an organic product of the oxidation of glucose.

..... [1]

- (f) (i) Calculate the maximum percentage error in the reaction time recorded for **Experiment 1**. Assume the error of the timer is  $\pm 1 \text{ s}$ .

maximum percentage error in **Experiment 1** = ..... % [1]

- (ii) You have carried out experiments at five different temperatures.

Identify an experiment, if any, you should have repeated. Give a reason for your answer.

.....  
 .....  
 ..... [1]

- (g) Suggest **two** ways to improve the accuracy of the results for this investigation.

1 .....  
 .....  
 2 .....  
 ..... [2]

[Total: 25]

## Qualitative Analysis

Where reagents are selected for use in a test, the **name** or **correct formula** of the element or compound must be given.

At each stage of any test you are to record details of the following:

- colour changes seen;
- the formation of any precipitate and its solubility in an excess of the reagent added;
- the formation of any gas and its identification by a suitable test.

You should indicate clearly at what stage in a test a change occurs.

If any solution is warmed, a **boiling tube** must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

**No additional tests for ions present should be attempted.**

- 2 Sandell's solution reacts in a similar way to Fehling's reagent.  
You will need to heat Sandell's solution in a hot water bath when using it in tests.

Half fill the 250 cm<sup>3</sup> beaker with water and place it on the tripod and gauze. Heat the water until it is boiling then turn off the Bunsen burner. This will be your hot water bath.

(a) **FB 4**, **FB 5** and **FB 6** are all solutions of carbohydrates.

- Sugars and starch are carbohydrates.
- Some sugars contain an aldehyde group so act as reducing agents.
- Other sugars do not contain an aldehyde group.

- (i) For each test use a 1 cm depth of the solution in a test-tube. Record all your observations in the table.

test	observations		
	FB 4	FB 5	FB 6
Add 2 or 3 drops of aqueous iodine.			
Add 2 or 3 drops of acidified potassium manganate(VII) and <b>allow to stand for two minutes.</b>			
Add a 3 cm depth of Sandell's solution and place the tube in the hot water bath for two minutes.			

[3]

- (ii) Circle the carbohydrate that could be starch.

FB 4

FB 5

FB 6

Circle the carbohydrate that contains an aldehyde group.

FB 4

FB 5

FB 6

[1]

- (iii) Suggest a different test, other than using Fehling's reagent, that could be carried out to identify the presence of an aldehyde group. State the reagent(s) you would use and the expected observation if the result were positive.

**Do not carry out your test.**

reagent(s) .....

observation .....

[1]

- (b) (i) **FB 7** and **FB 8** are two of the components of Sandell's solution. Each contains one cation and one anion. Two of the ions are listed in the Qualitative Analysis Notes.

For each test use a 1 cm depth of solution in a test-tube. Record all your observations in the table.

<i>test</i>	<i>observations</i>	
	<b>FB 7</b>	<b>FB 8</b>
Add a few drops of aqueous silver nitrate.		
Add a few drops of aqueous barium nitrate or aqueous barium chloride, then		
add dilute nitric acid.		
Add a few drops of aqueous iodine.		
Add a 1 cm depth of aqueous iron(II) sulfate.		
Add a 1 cm depth of <b>FB 8</b> .		X

[6]



- (ii) Identify the ions in **FB 7** and **FB 8**. If you are unable to identify any of the ions, write 'unknown'.

**FB 7** cation ..... anion .....

**FB 8** cation ..... anion .....

[2]

- (iii) Write an ionic equation for any reaction in (i) that produced a precipitate. Include state symbols.

..... [2]

[Total: 15]

## Qualitative Analysis Notes

## 1 Reactions of aqueous cations

ion	reaction with	
	NaOH(aq)	NH <sub>3</sub> (aq)
aluminium, Al <sup>3+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH <sub>4</sub> <sup>+</sup> (aq)	no ppt. ammonia produced on heating	–
barium, Ba <sup>2+</sup> (aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.
calcium, Ca <sup>2+</sup> (aq)	white ppt. with high [Ca <sup>2+</sup> (aq)]	no ppt.
chromium(III), Cr <sup>3+</sup> (aq)	grey-green ppt. soluble in excess	grey-green ppt. insoluble in excess
copper(II), Cu <sup>2+</sup> (aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe <sup>2+</sup> (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe <sup>3+</sup> (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg <sup>2+</sup> (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn <sup>2+</sup> (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. soluble in excess

## 2 Reactions of anions

<i>ion</i>	<i>reaction</i>
carbonate, $\text{CO}_3^{2-}$	$\text{CO}_2$ liberated by dilute acids
chloride, $\text{Cl}^-(\text{aq})$	gives white ppt. with $\text{Ag}^+(\text{aq})$ (soluble in $\text{NH}_3(\text{aq})$ )
bromide, $\text{Br}^-(\text{aq})$	gives cream ppt. with $\text{Ag}^+(\text{aq})$ (partially soluble in $\text{NH}_3(\text{aq})$ )
iodide, $\text{I}^-(\text{aq})$	gives yellow ppt. with $\text{Ag}^+(\text{aq})$ (insoluble in $\text{NH}_3(\text{aq})$ )
nitrate, $\text{NO}_3^-(\text{aq})$	$\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and Al foil
nitrite, $\text{NO}_2^-(\text{aq})$	$\text{NH}_3$ liberated on heating with $\text{OH}^-(\text{aq})$ and Al foil
sulfate, $\text{SO}_4^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (insoluble in excess dilute strong acids)
sulfite, $\text{SO}_3^{2-}(\text{aq})$	gives white ppt. with $\text{Ba}^{2+}(\text{aq})$ (soluble in excess dilute strong acids)

## 3 Tests for gases

<i>gas</i>	<i>test and test result</i>
ammonia, $\text{NH}_3$	turns damp red litmus paper blue
carbon dioxide, $\text{CO}_2$	gives a white ppt. with limewater (ppt. dissolves with excess $\text{CO}_2$ )
chlorine, $\text{Cl}_2$	bleaches damp litmus paper
hydrogen, $\text{H}_2$	'pops' with a lighted splint
oxygen, $\text{O}_2$	relights a glowing splint

## The Periodic Table of Elements

		Group															
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">1 H hydrogen 1.0</div> <div style="border: 1px solid black; padding: 2px;">2 He helium 4.0</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">3 Li lithium 6.9</div> <div style="border: 1px solid black; padding: 2px;">4 Be beryllium 9.0</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">5 B boron 10.8</div> <div style="border: 1px solid black; padding: 2px;">6 C carbon 12.0</div> <div style="border: 1px solid black; padding: 2px;">7 N nitrogen 14.0</div> <div style="border: 1px solid black; padding: 2px;">8 O oxygen 16.0</div> <div style="border: 1px solid black; padding: 2px;">9 F fluorine 19.0</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">11 Na sodium 23.0</div> <div style="border: 1px solid black; padding: 2px;">12 Mg magnesium 24.3</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">13 Al aluminium 27.0</div> <div style="border: 1px solid black; padding: 2px;">14 Si silicon 28.1</div> <div style="border: 1px solid black; padding: 2px;">15 P phosphorus 31.0</div> <div style="border: 1px solid black; padding: 2px;">16 S sulfur 32.1</div> <div style="border: 1px solid black; padding: 2px;">17 Cl chlorine 35.5</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">19 K potassium 39.1</div> <div style="border: 1px solid black; padding: 2px;">20 Ca calcium 40.1</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">21 Sc scandium 45.0</div> <div style="border: 1px solid black; padding: 2px;">22 Ti titanium 47.9</div> <div style="border: 1px solid black; padding: 2px;">23 V vanadium 50.9</div> <div style="border: 1px solid black; padding: 2px;">24 Cr chromium 52.0</div> <div style="border: 1px solid black; padding: 2px;">25 Mn manganese 54.9</div> <div style="border: 1px solid black; padding: 2px;">26 Fe iron 55.8</div> <div style="border: 1px solid black; padding: 2px;">27 Co cobalt 58.9</div> <div style="border: 1px solid black; padding: 2px;">28 Ni nickel 58.7</div> <div style="border: 1px solid black; padding: 2px;">29 Cu copper 63.5</div> <div style="border: 1px solid black; padding: 2px;">30 Zn zinc 65.4</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">31 Ga gallium 69.7</div> <div style="border: 1px solid black; padding: 2px;">32 Ge germanium 72.6</div> <div style="border: 1px solid black; padding: 2px;">33 As arsenic 74.9</div> <div style="border: 1px solid black; padding: 2px;">34 Se selenium 79.0</div> <div style="border: 1px solid black; padding: 2px;">35 Br bromine 79.9</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">37 Rb rubidium 85.5</div> <div style="border: 1px solid black; padding: 2px;">38 Sr strontium 87.6</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">39 Y yttrium 88.9</div> <div style="border: 1px solid black; padding: 2px;">40 Zr zirconium 91.2</div> <div style="border: 1px solid black; padding: 2px;">41 Nb niobium 92.9</div> <div style="border: 1px solid black; padding: 2px;">42 Mo molybdenum 95.9</div> <div style="border: 1px solid black; padding: 2px;">43 Tc technetium —</div> <div style="border: 1px solid black; padding: 2px;">44 Ru ruthenium 101.1</div> <div style="border: 1px solid black; padding: 2px;">45 Rh rhodium 102.9</div> <div style="border: 1px solid black; padding: 2px;">46 Pd palladium 106.4</div> <div style="border: 1px solid black; padding: 2px;">47 Ag silver 107.9</div> <div style="border: 1px solid black; padding: 2px;">48 Cd cadmium 112.4</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">51 Sb antimony 121.8</div> <div style="border: 1px solid black; padding: 2px;">52 Te tellurium 127.6</div> <div style="border: 1px solid black; padding: 2px;">53 I iodine 126.9</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">55 Cs caesium 132.9</div> <div style="border: 1px solid black; padding: 2px;">56 Ba barium 137.3</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">57–71 lanthanoids</div> <div style="border: 1px solid black; padding: 2px;">72 Hf hafnium 178.5</div> <div style="border: 1px solid black; padding: 2px;">73 Ta tantalum 180.9</div> <div style="border: 1px solid black; padding: 2px;">74 W tungsten 183.8</div> <div style="border: 1px solid black; padding: 2px;">75 Re rhenium 186.2</div> <div style="border: 1px solid black; padding: 2px;">76 Os osmium 190.2</div> <div style="border: 1px solid black; padding: 2px;">77 Ir iridium 192.2</div> <div style="border: 1px solid black; padding: 2px;">78 Pt platinum 195.1</div> <div style="border: 1px solid black; padding: 2px;">79 Au gold 197.0</div> <div style="border: 1px solid black; padding: 2px;">80 Hg mercury 200.6</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">81 Tl thallium 204.4</div> <div style="border: 1px solid black; padding: 2px;">82 Pb lead 207.2</div> <div style="border: 1px solid black; padding: 2px;">83 Bi bismuth 209.0</div> <div style="border: 1px solid black; padding: 2px;">84 Po polonium —</div> <div style="border: 1px solid black; padding: 2px;">85 At astatine —</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">87 Fr francium —</div> <div style="border: 1px solid black; padding: 2px;">88 Ra radium —</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">89–103 actinoids</div> <div style="border: 1px solid black; padding: 2px;">104 Rf rutherfordium —</div> <div style="border: 1px solid black; padding: 2px;">105 Db dubnium —</div> <div style="border: 1px solid black; padding: 2px;">106 Sg seaborgium —</div> <div style="border: 1px solid black; padding: 2px;">107 Bh bohrium —</div> <div style="border: 1px solid black; padding: 2px;">108 Hs hassium —</div> <div style="border: 1px solid black; padding: 2px;">109 Mt meitnerium —</div> <div style="border: 1px solid black; padding: 2px;">110 Ds darmstadtium —</div> <div style="border: 1px solid black; padding: 2px;">111 Rg roentgenium —</div> <div style="border: 1px solid black; padding: 2px;">112 Cn copernicium —</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 2px;">113 Nh nihonium —</div> <div style="border: 1px solid black; padding: 2px;">114 Fl flerovium —</div> <div style="border: 1px solid black; padding: 2px;">115 Mc moscovium —</div> <div style="border: 1px solid black; padding: 2px;">116 Lv livermorium —</div> <div style="border: 1px solid black; padding: 2px;">117 Ts tennessine —</div> <div style="border: 1px solid black; padding: 2px;">118 Og oganeson —</div> </div>															

57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.4	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
89 Ac actinium	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —

lanthanoids

actinoids