



## Cambridge International AS & A Level

---

**CHEMISTRY**

**9701/23**

Paper 3 AS Level Structured Questions

**May/June 2020**

MARK SCHEME

Maximum Mark: 60

---

**Published**

Students did not sit exam papers in the June 2020 series due to the Covid-19 global pandemic.

This mark scheme is published to support teachers and students and should be read together with the question paper. It shows the requirements of the exam. The answer column of the mark scheme shows the proposed basis on which Examiners would award marks for this exam. Where appropriate, this column also provides the most likely acceptable alternative responses expected from students. Examiners usually review the mark scheme after they have seen student responses and update the mark scheme if appropriate. In the June series, Examiners were unable to consider the acceptability of alternative responses, as there were no student responses to consider.

Mark schemes should usually be read together with the Principal Examiner Report for Teachers. However, because students did not sit exam papers, there is no Principal Examiner Report for Teachers for the June 2020 series.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the June 2020 series for most Cambridge IGCSE™ and Cambridge International A & AS Level components, and some Cambridge O Level components.

---

This document consists of **11** printed pages.

**Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

**GENERIC MARKING PRINCIPLE 1:**

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

**GENERIC MARKING PRINCIPLE 2:**

Marks awarded are always **whole marks** (not half marks, or other fractions).

**GENERIC MARKING PRINCIPLE 3:**

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

**GENERIC MARKING PRINCIPLE 4:**

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

**GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

**GENERIC MARKING PRINCIPLE 6:**

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

**Science-Specific Marking Principles**

- |   |   |
|---|---|
| 1 | Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.   |
| 2 | The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.   |
| 3 | Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).   |
| 4 | The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.  |
| 5 | <p><u>'List rule' guidance</u> (see examples below)</p> <p>For questions that require <i>n</i> responses (e.g. State <b>two</b> reasons ...):</p> <ul style="list-style-type: none"><li>• The response should be read as continuous prose, even when numbered answer spaces are provided</li><li>• Any response marked <i>ignore</i> in the mark scheme should not count towards <i>n</i></li><li>• Incorrect responses should not be awarded credit but will still count towards <i>n</i></li><li>• Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should <b>not</b> be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response</li><li>• Non-contradictory responses after the first <i>n</i> responses may be ignored even if they include incorrect science.</li></ul> |

**6** Calculation specific guidance

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form, (e.g.  $a \times 10^n$ ) in which the convention of restricting the value of the coefficient ( $a$ ) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.


Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

**7** Guidance for chemical equations


Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.


Question	Answer	Marks						
1(a)(i)	<b>M1</b> green flame / white flame (with green tinge) <b>M2</b> white solid formed	<b>2</b>						
1(a)(ii)	$2\text{Ba(s)} + \text{O}_2\text{(g)} \rightarrow 2\text{BaO(s)}$	<b>1</b>						
1(b)(i)	<b>M1</b> heat (followed by) add water <b>M2</b> $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$ <b>M3</b> $\text{CaO} + \text{H}_2\text{O} \rightarrow \text{Ca(OH)}_2$	<b>3</b>						
1(b)(ii)	thermal decomposition.	<b>1</b>						
1(b)(iii)	neutralise acid soil / reduce the acidity in soil / increase the pH in soil	<b>1</b>						
1(c)(i)	$4\text{Ga} + 3\text{O}_2 \rightarrow 2\text{Ga}_2\text{O}_3$	<b>1</b>						
1(c)(ii)	(+) <b>3</b>	<b>1</b>						
1(c)(ii)	<table border="1"> <thead> <tr> <th>reagent and conditions</th> <th>Formula of gallium containing product</th> </tr> </thead> <tbody> <tr> <td><b>M1</b> gallium oxide+ hot HCl(aq)</td> <td><b>M1</b> <math>\text{GaCl}_3</math></td> </tr> <tr> <td><b>M2</b> gallium oxide+hot concentrated NaOH(aq)</td> <td><b>M2</b> <math>\text{NaGa(OH)}_4</math> OR <math>\text{NaGaO}_2</math></td> </tr> </tbody> </table>	reagent and conditions	Formula of gallium containing product	<b>M1</b> gallium oxide+ hot HCl(aq)	<b>M1</b> $\text{GaCl}_3$	<b>M2</b> gallium oxide+hot concentrated NaOH(aq)	<b>M2</b> $\text{NaGa(OH)}_4$ OR $\text{NaGaO}_2$	<b>2</b>
reagent and conditions	Formula of gallium containing product							
<b>M1</b> gallium oxide+ hot HCl(aq)	<b>M1</b> $\text{GaCl}_3$							
<b>M2</b> gallium oxide+hot concentrated NaOH(aq)	<b>M2</b> $\text{NaGa(OH)}_4$ OR $\text{NaGaO}_2$							

Question	Answer	Marks
2(a)	<p><b>EITHER</b></p> <p><b>M1</b> mass of an atom / isotope</p> <p><b>M2</b> relative / compared to 1/12 (the mass) of (an atom of) C-12 OR on a scale in which a C-12 (atom / isotope) has (a mass of exactly) 12 (units)</p> <p><b>OR</b></p> <p><b>M1</b> mass of one mol (of atoms) of an isotope</p> <p><b>M2</b> relative / compared to 1/12 (the mass) of 1 mol of C-12 OR in which one mol C-12 (atom / isotope) has a mass of (exactly) 12 g</p>	2
2(b)	<p>% abundance of <math>^{63}\text{Cu}</math> = 72.5%</p> <p>% abundance of <math>^{65}\text{Cu}</math> = 27.5%</p> <p><b>M1</b> correct algebraic expression AND correct calculation of <math>x</math> for one isotope</p> <p>% ab of <math>^{63}\text{Cu}</math> = <math>x</math> <math>(x/100 \times 63) + ((1-x)/100 \times 65) = 63.55</math> so <math>x = 72.5</math></p> <p>OR</p> <p>% ab of <math>^{65}\text{Cu}</math> = <math>x</math> <math>(1-x)/100 \times 63) + x/100 \times 65) = 63.55</math> so <math>x = 27.5</math></p> <p><b>M2</b> calculation of abundance of other isotope by <math>100 - x</math></p>	2
2(c)(i)	metallic	1
2(c)(ii)	<p>diagram showing the bonding in a sample of copper</p>  <p><b>M1</b> diagram shows regular arrangement of spheres labelled as positively charged ions / +2 or +1 / cations</p> <p><b>M2</b> diagram shows surrounded by electrons and clearly labelled as 'delocalised electrons'</p>	3
2(c)(iii)	$(1s^2) 2s^2 2p^6 3s^2 3p^6 3d^{10} 4s^1$ OR $(1s^2) 2s^2 2p^6 3s^2 3p^6 4s^1 3d^{10}$	1

Question	Answer	Marks						
2(d)(i)	<p><b>M1</b> calculate the number mol <math>S_2O_3^{2-}</math> added  <math>20/1000 \times 0.10 = 2 \times 10^{-3} = 0.002</math> (mol <math>S_2O_3^{2-}</math>)</p> <p><b>M2</b> calculate number mol <math>CuSO_4</math> in <math>250cm^3</math>            (1 mol <math>S_2O_3^{2-}</math> : 1 mol <math>CuSO_4</math>) = 0.002 mol <math>CuSO_4</math> in <math>25cm^3</math></p> <p>so 0.02 mol <math>CuSO_4</math> in <math>250cm^3</math></p>	2						
2(d)(ii)	<table border="1"> <tbody> <tr> <td><b>M1</b> amount of <math>CuSO_4</math> in 10.68 g of <math>CuSO_4 \cdot xH_2O</math></td> <td><math>7.98 / (159.6) = \underline{0.05}</math> (mol)</td> </tr> <tr> <td><b>M2</b> amount of <math>H_2O</math> in 10.68 g of <math>CuSO_4 \cdot xH_2O</math></td> <td><math>(10.68 - 7.98) / 18 = 2.7 / 18 = \underline{0.15}</math> (mol)</td> </tr> <tr> <td><b>M3</b> value of x</td> <td>(mol <math>H_2O \div</math> mol <math>CuSO_4 =</math>) 3</td> </tr> </tbody> </table>	<b>M1</b> amount of $CuSO_4$ in 10.68 g of $CuSO_4 \cdot xH_2O$	$7.98 / (159.6) = \underline{0.05}$ (mol)	<b>M2</b> amount of $H_2O$ in 10.68 g of $CuSO_4 \cdot xH_2O$	$(10.68 - 7.98) / 18 = 2.7 / 18 = \underline{0.15}$ (mol)	<b>M3</b> value of x	(mol $H_2O \div$ mol $CuSO_4 =$ ) 3	3
<b>M1</b> amount of $CuSO_4$ in 10.68 g of $CuSO_4 \cdot xH_2O$	$7.98 / (159.6) = \underline{0.05}$ (mol)							
<b>M2</b> amount of $H_2O$ in 10.68 g of $CuSO_4 \cdot xH_2O$	$(10.68 - 7.98) / 18 = 2.7 / 18 = \underline{0.15}$ (mol)							
<b>M3</b> value of x	(mol $H_2O \div$ mol $CuSO_4 =$ ) 3							

Question	Answer	Marks
3(a)(i)	hydrolysis	1
3(a)(ii)	 <p><b>M3</b> label / shading indicating greater area under curve of <math>E_a(\text{cat})</math> AND state greater number / proportion of sucrose molecules have energy greater than <math>E_a(\text{cat})</math> (so faster rate)</p> <p><b>M2</b> <math>E_a</math> cat</p> <p><b>M1</b> <math>E_a</math> (no cat)</p>	3



Question	Answer	Marks
3(b)(i)	an atom which is bonded to four different substituents / groups / atoms	1
3(b)(ii)	ALL three chiral carbons need to be shown by * 	1
3(b)(iii)	<i>empirical formula of fructose</i> CH <sub>2</sub> O	1
3(c)(i)	<b>M1</b> (enthalpy change) when 1 mole of a substance <b>M2</b> burns/combusts/reacts in excess air/oxygen <b>OR</b> completely burns/combusts/reacts in air/oxygen	2
3(c)(ii)	$C_{12}H_{22}O_{11} + 12 O_2 \rightarrow 12CO_2 + 11H_2O$	1
3(c)(iii)	<b>M1</b> $\Delta H = -5643 - (-2805 + -2810)$ <b>M2</b> = $-28 \text{ kJ mol}^{-1}$	2

Question	Answer	Marks
4(a)	<b>M1</b> (no reaction) not enough energy <b>M2</b> bromine (free) radicals are not produced OR homolytic fission of bromine does not occur.	2
4(b)(i)	free-radical substitution	1

Question	Answer	Marks
4(b)(ii)	$C_6H_{14} + (\cdot)Br \rightarrow (\cdot)C_6H_{13} + HBr$	1
4(b)(iii)	$(\cdot)C_6H_{13} + Br_2 \rightarrow C_6H_{13}Br + (\cdot)Br$	1
4(b)(iv)	$(\cdot)C_6H_{13} + (\cdot)Br \rightarrow C_6H_{13}Br$	1
4(c)(i)	$CH_3(CH_2)_3CH=CH_2$	1
4(c)(ii)	cold <b>AND</b> acidified <b>AND</b> dilute 2 marks for 3 correct conditions 1 mark for 2 correct conditions	2
4(d)(i)	addition <b>OR</b> reduction	1
4(d)(ii)	<b>M1</b> only sigma / $\sigma$ (bonds) in hexane / alkanes <b>M2 B</b> has sigma $\sigma$ (bonds) and a / one pi / $\pi$ (bond)	2

Question	Answer	Marks
5(a)(i)	dehydration	1
5(a)(ii)	<b>M1</b> correct identification of butan-2-ol <b>M2</b> correct displayed formula including correct connectivity of C–O–H $  \begin{array}{cccc}  H & H & H & H \\    &   &   &   \\  H-C & -C & -C & -C-H \\    &   &   &   \\  H & H & O & H \\  & &   & \\  & & H &   \end{array}  $	2

Question	Answer	Marks								
5(a)(iii)	<table border="1" data-bbox="338 213 952 478"> <tr> <td data-bbox="338 213 472 279">isomer</td> <td data-bbox="472 213 952 279">name</td> </tr> <tr> <td data-bbox="338 279 472 344">C</td> <td data-bbox="472 279 952 344">cis but-2-ene</td> </tr> <tr> <td data-bbox="338 344 472 410">D</td> <td data-bbox="472 344 952 410">trans but-2-ene</td> </tr> <tr> <td data-bbox="338 410 472 475">E</td> <td data-bbox="472 410 952 475">but -1-ene</td> </tr> </table> <p data-bbox="338 512 705 576">2 marks for 3 correct names 1 mark for 2 correct names</p>	isomer	name	C	cis but-2-ene	D	trans but-2-ene	E	but -1-ene	2
isomer	name									
C	cis but-2-ene									
D	trans but-2-ene									
E	but -1-ene									
5(b)(i)	<table border="1" data-bbox="338 608 952 906"> <tr> <td data-bbox="338 608 409 673"></td> <td data-bbox="409 608 952 673">Functional group</td> </tr> <tr> <td data-bbox="338 673 409 738">F</td> <td data-bbox="409 673 952 738"><b>M1</b> Ester / RCOOR<sup>(<sup>o</sup>)</sup> / RCO<sub>2</sub>R<sup>(<sup>o</sup>)</sup></td> </tr> <tr> <td data-bbox="338 738 409 836">G</td> <td data-bbox="409 738 952 836"><b>M2</b> Carboxyl / carboxylic acid / RCOOH</td> </tr> <tr> <td data-bbox="338 836 409 906">H</td> <td data-bbox="409 836 952 906"><b>M3</b> Alcohol / hydroxy / R-OH</td> </tr> </table>		Functional group	F	<b>M1</b> Ester / RCOOR <sup>(<sup>o</sup>)</sup> / RCO <sub>2</sub> R <sup>(<sup>o</sup>)</sup>	G	<b>M2</b> Carboxyl / carboxylic acid / RCOOH	H	<b>M3</b> Alcohol / hydroxy / R-OH	3
	Functional group									
F	<b>M1</b> Ester / RCOOR <sup>(<sup>o</sup>)</sup> / RCO <sub>2</sub> R <sup>(<sup>o</sup>)</sup>									
G	<b>M2</b> Carboxyl / carboxylic acid / RCOOH									
H	<b>M3</b> Alcohol / hydroxy / R-OH									
5(b)(ii)	triiodomethane	1								
5(b)(iii)	<b>M1 G</b> = HCOOH / HCO <sub>2</sub> H <b>M2 H</b> = C <sub>2</sub> H <sub>5</sub> OH	2								