



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



CHEMISTRY

9701/42

Paper 4 A Level Structured Questions

May/June 2023

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 100.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.
- Important values, constants and standards are printed in the question paper.

This document has **24** pages.

1 (a) Group 2 carbonates decompose when heated to form the metal oxide and carbon dioxide.

- (i) Suggest a mechanism for the decomposition of the carbonate ion by adding **two** curly arrows in Fig. 1.1.



Fig. 1.1

[1]

- (ii) Describe the variation in the thermal stability of Group 2 carbonates. Explain your answer.

.....

 [3]

- (b) (i) Define lattice energy.

.....

 [2]

- (ii) The lattice energy of the Group 2 carbonates, $\Delta H_{\text{latt}}^{\ominus}(\text{MCO}_3)$, becomes less exothermic down the group.

The lattice energy of the Group 2 oxides, $\Delta H_{\text{latt}}^{\ominus}(\text{MO})$, also becomes less exothermic down the group.

$\Delta H_{\text{latt}}^{\ominus}(\text{MCO}_3)$ and $\Delta H_{\text{latt}}^{\ominus}(\text{MO})$ change by **different** amounts going down the group.

Suggest how the standard enthalpy change of the decomposition reaction for Group 2 carbonates changes down the group.

Explain your reasoning in terms of the relative sizes of the anions and the relative changes in lattice energy down the group.

.....


 [2]

(c) Potassium sulfite, K_2SO_3 , is used as a food additive.

The concentration of sulfite ions, SO_3^{2-} , can be determined by titration using aqueous acidified manganate(VII) ions, MnO_4^- .

- A 250 cm^3 solution contains 3.40 g of impure K_2SO_3 .
- 25.0 cm^3 of this solution requires 22.40 cm^3 of $0.0250\text{ mol dm}^{-3}$ acidified MnO_4^- to reach the end-point. All the SO_3^{2-} ions are oxidised. None of the other species in the impure K_2SO_3 are oxidised.

The reaction occurs as shown by the two half-equations.



(i) Give the ionic equation for the reaction between SO_3^{2-} and acidified MnO_4^- .

..... [1]

(ii) Calculate the percentage purity of the sample of K_2SO_3 .
Show your working.

percentage purity of $\text{K}_2\text{SO}_3 = \dots\dots\dots$ [3]

(d) Potassium disulfite, $\text{K}_2\text{S}_2\text{O}_5$, is another food additive. The disulfite ion, $\text{S}_2\text{O}_5^{2-}$, has the displayed formula shown in Fig. 1.2.




Fig. 1.2

Deduce the geometry (shape) around the $\text{S}(\alpha)$ atom in $\text{S}_2\text{O}_5^{2-}$.

geometry around $\text{S}(\alpha) \dots\dots\dots$ [1]

2 (a) State **two** typical chemical properties of a transition element.

1

2

[1]

(b) Aqueous solutions of cobalt(II) salts contain the complex ion $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$.

(i) Define complex ion.

.....

..... [1]

(ii) Samples of $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ are reacted separately with an excess of aqueous ammonia, with an excess of concentrated HCl and with an excess of aqueous sodium hydroxide, as shown in Fig. 2.1.




Fig. 2.1

Complete Table 2.1 about the reactions shown by $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$.

Table 2.1

reagent added to $[\text{Co}(\text{H}_2\text{O})_6]^{2+}(\text{aq})$	formula of cobalt species formed	colour and state of cobalt species formed	type of reaction
an excess of $\text{NH}_3(\text{aq})$	A =		
an excess of concentrated HCl	B =		
an excess of $\text{NaOH}(\text{aq})$	C =		

[4]

(c) The ethanedioate ion, $\text{C}_2\text{O}_4^{2-}$, can act as a bidentate ligand.

(i) Explain what is meant by a bidentate ligand.

.....

 [2]

(ii) The complex $[\text{Co}(\text{H}_2\text{O})_2(\text{C}_2\text{O}_4)\text{BrCl}]^-$ exists as stereoisomers.

Complete the three-dimensional diagrams in Fig. 2.2 to show **four** stereoisomers of $[\text{Co}(\text{H}_2\text{O})_2(\text{C}_2\text{O}_4)\text{BrCl}]^-$.

The $\text{C}_2\text{O}_4^{2-}$ ligand is represented using $\text{O} \text{---} \text{O}$.




Fig. 2.2

[3]

(iii) State the oxidation state of cobalt in this complex and a type of stereoisomerism shown.

oxidation state of cobalt

type of stereoisomerism

[1]

[Total: 12]

- 3 (a) Complete Table 3.1 by placing **one** tick (✓) in each row to indicate the sign of each type of energy change under standard conditions.

Table 3.1

energy change	always positive	always negative	can be either negative or positive
bond energy			
enthalpy change of atomisation			
enthalpy change of formation			

[1]

- (b) Define standard enthalpy change of atomisation.

.....

.....

..... [2]

- (c) Table 3.2 shows some energy changes.

Table 3.2

energy change	value/kJ mol ⁻¹
standard enthalpy change of atomisation of silver	+285
first ionisation energy of silver	+731
second ionisation energy of silver	+2074
bond energy of O=O	+496
bond energy of O–O	+150
first electron affinity of oxygen	–141
second electron affinity of oxygen	+798
first ionisation energy of oxygen	+1314
standard enthalpy change of formation of silver oxide, Ag ₂ O(s)	–31

Calculate the lattice energy, $\Delta H_{\text{latt}}^{\ominus}$, of $\text{Ag}_2\text{O}(\text{s})$ using relevant data from Table 3.2.

It may be helpful to draw a labelled energy cycle.

Show your working.

$\Delta H_{\text{latt}}^{\ominus}$ of $\text{Ag}_2\text{O}(\text{s}) = \dots\dots\dots \text{kJ mol}^{-1}$ [3]

- (d) Suggest the trend in the magnitude of the lattice energies of the silver compounds Ag_2S , Ag_2O and Ag_2Se .

Explain your answer.

.....
 least exothermic most exothermic

[2]

(e) Silver sulfite, $\text{Ag}_2\text{SO}_3(\text{s})$, is sparingly soluble in water.

(i) Give an expression for the solubility product, K_{sp} , of Ag_2SO_3 .

$$K_{\text{sp}} =$$

[1]

(ii) Calculate the equilibrium concentration of Ag^+ in a saturated solution of Ag_2SO_3 at 298 K.

[K_{sp} : Ag_2SO_3 , $1.50 \times 10^{-14} \text{ mol}^3 \text{ dm}^{-9}$ at 298 K]

$$[\text{Ag}^+] = \dots\dots\dots \text{ mol dm}^{-3} \quad [1]$$

(f) The standard enthalpy change of solution, $\Delta H_{\text{sol}}^\ominus$, of $\text{AgNO}_3(\text{s})$ in water is $+22.6 \text{ kJ mol}^{-1}$.

Suggest how the feasibility of dissolving $\text{AgNO}_3(\text{s})$ in water changes with temperature.

Explain your answer.

.....


.....

.....

..... [2]

[Total: 12]

- 4 (a) In aqueous solution, iron(III) ions react with iodide ions, as shown.



A series of experiments is carried out using different concentrations of Fe^{3+} and I^{-} , as shown in Table 4.1.

Table 4.1

experiment	$[\text{Fe}^{3+}]/\text{mol dm}^{-3}$	$[\text{I}^{-}]/\text{mol dm}^{-3}$	initial rate/ $\text{mol dm}^{-3}\text{s}^{-1}$
1	0.0400	0.0200	2.64×10^{-4}
2	0.1200	0.0200	7.92×10^{-4}
3	0.0800	0.0400	2.11×10^{-3}

- (i) Explain what is meant by overall order of reaction.

.....

 [1]

- (ii) Use the data in Table 4.1 to deduce the order of reaction with respect to Fe^{3+} and with respect to I^{-} .

Explain your reasoning.

.....

 [2]

- (iii) Use your answer to (a)(ii) to construct the rate equation for this reaction.

rate = [1]


- (iv) Use your answer to (a)(iii) and the data from experiment 1 to calculate the rate constant, k , for this reaction. Include the units of k .

$k = \dots\dots\dots$ units [2]

- (v) Describe qualitatively the effect of an increase in temperature on the rate constant and on the rate of this reaction.

.....
 [1]

(b) In aqueous solution, iodide ions react with acidified hydrogen peroxide, as shown.



The initial rate of reaction is found to be first order with respect to I^- , first order with respect to H_2O_2 and zero order with respect to H^+ .

Fig. 4.1 shows a possible four-step mechanism for this reaction.




Fig. 4.1

(i) Suggest which of the steps, 1, 2, 3 or 4, in this mechanism is the rate-determining step.

Explain your answer.

.....

 [1]

(ii) Identify a step in Fig. 4.1 that involves a redox reaction.

Explain your answer in terms of oxidation numbers.

.....

 [1]

(iii) Suggest the role of HIO in this mechanism.

Explain your reasoning.

.....

 [1]

[Total: 10]

5 (a) Methylbenzene can undergo different reactions, as shown in Fig. 5.1.




Fig. 5.1

- (i) Draw structures in Fig. 5.1 for the possible organic products of the three reactions shown. [3]
- (ii) Complete Table 5.1.

Table 5.1

	type of reaction	mechanism
reaction 1		
reaction 2		

[2]

- (b) When methylbenzene reacts with an electrophile, a substitution reaction occurs. No addition reaction takes place under these conditions.

Explain why no addition reaction takes place.

.....
 [1]

- (c) The reaction of methylbenzene with thionyl bromide, SOBr_2 , in the presence of an iron(III) bromide catalyst, FeBr_3 , is shown in Fig. 5.2.





Fig. 5.2

The mechanism of this reaction is similar to that of the bromination of benzene.

The first step of the mechanism generates the SOBr^+ electrophile, as shown.



- (i) The reaction of methylbenzene with SOBr^+ ions is shown in Fig. 5.3. Complete the mechanism in Fig. 5.3.

Include all relevant curly arrows and charges.

Draw the structure of the organic intermediate.




Fig. 5.3

[3]

- 6 (a) Perindopril is a drug used to treat heart disease.

perindopril




Fig. 6.1

- (i) State the number of chiral carbon atoms present in one molecule of perindopril.

..... [1]

- (ii) Suggest **one** benefit and **one** disadvantage of producing a drug such as perindopril as a single pure optical isomer.

benefit

.....

disadvantage

.....

[2]




- (b) (i) Name all the functional groups in perindopril.

.....

..... [2]

- (ii) A sample of perindopril is hydrolysed with hot aqueous acid.

Draw the structures of the **three** organic products of the **complete** acid hydrolysis of perindopril.



[3]

[Total: 8]

- 7 (a) Explain why phenol is brominated much more easily than benzene is brominated.

.....

.....

.....

.....

..... [3]

- (b) Iodine monobromide, I-Br, reacts with benzene in the presence of an AlBr₃ catalyst.

Predict whether the organic product will be bromobenzene or iodobenzene.
Explain your answer.

.....

.....

..... [1]

- (c) Fig. 7.1 shows some reactions of phenol.




Fig. 7.1

- (i) Give an equation for the reaction of phenol with Na(s).

..... [1]

- (ii) Draw the structure of the organic product, **R**, formed when phenol reacts with an excess of $\text{Br}_2(\text{aq})$.

[1]

- (iii) State the reagents and conditions for reaction 1 and reaction 2 in Fig. 7.1.

reaction 1

reaction 2

[2]

[Total: 8]

- 8 (a) Describe the difference in reactivity between chloroethane and chlorobenzene with $\text{OH}^-(\text{aq})$.
Explain your answer.

.....

.....

.....

..... [2]

- (b) Compound **T**, $\text{C}_5\text{H}_9\text{O}_2\text{Cl}$, is a useful synthetic intermediate.

Fig. 8.1 shows some reactions of **T**.




Fig. 8.1

(i) Give the systematic name for **T**.

..... [1]

(ii) Draw the structures of **W**, **X**, **Y** and **Z** in Fig. 8.1. [4]

(iii) State the reagents and conditions for steps 1 and 2 in Fig. 8.1.

step 1

step 2

[2]

(c) The proton (^1H) NMR spectrum of compound **T**, $\text{C}_5\text{H}_9\text{O}_2\text{Cl}$, in CDCl_3 is shown in Fig. 8.2.




Fig. 8.2

Table 8.1

environment of proton	example	chemical shift range δ /ppm
alkane	$-\text{CH}_3$, $-\text{CH}_2-$, $>\text{CH}-$	0.9–1.7
alkyl next to $\text{C}=\text{O}$	$\text{CH}_3-\text{C}=\text{O}$, $-\text{CH}_2-\text{C}=\text{O}$, $>\text{CH}-\text{C}=\text{O}$	2.2–3.0
alkyl next to aromatic ring	CH_3-Ar , $-\text{CH}_2-\text{Ar}$, $>\text{CH}-\text{Ar}$	2.3–3.0
alkyl next to electronegative atom	CH_3-O , $-\text{CH}_2-\text{O}$, $-\text{CH}_2-\text{Cl}$	3.2–4.0
attached to alkene	$=\text{CHR}$	4.5–6.0
attached to aromatic ring	$\text{H}-\text{Ar}$	6.0–9.0
aldehyde	HCOR	9.3–10.5
alcohol	ROH	0.5–6.0
phenol	$\text{Ar}-\text{OH}$	4.5–7.0
carboxylic acid	RCOOH	9.0–13.0
alkyl amine	$\text{R}-\text{NH}-$	1.0–5.0
aryl amine	$\text{Ar}-\text{NH}_2$	3.0–6.0
amide	RCONHR	5.0–12.0

- (i) Suggest why CDCl_3 is used as a solvent instead of CHCl_3 for the proton (^1H) NMR spectrum.

.....
 [1]

- (ii) Complete Table 8.2 for the proton (^1H) NMR spectrum of **T**.

Table 8.2

chemical shift δ /ppm	environment of proton	splitting pattern	number of ^1H atoms responsible for the peak
1.2			
2.8			
3.7			
3.9			

[4]

- (iii) Explain the splitting pattern of the peak at δ 3.9 ppm.

.....
 [1]

[Total: 15]

- 9 (a) Define standard cell potential, $E_{\text{cell}}^{\ominus}$.

.....
 [1]

- (b) An electrochemical cell is set up to measure $E_{\text{cell}}^{\ominus}$ of a cell consisting of an $\text{Fe}^{3+}/\text{Fe}^{2+}$ half-cell and a Cl_2/Cl^- half-cell.

Draw a labelled diagram of this electrochemical cell.

Include all necessary substances. It is **not** necessary to state conditions used.

[3]

- (c) The cell reaction for the electrochemical cell in (b) is shown.



Calculate ΔG^{\ominus} , in kJ mol^{-1} , for this cell reaction.

$\Delta G^{\ominus} = \dots\dots\dots \text{kJ mol}^{-1}$ [2]

(d) Another experiment is set up using the same electrochemical cell.

In this experiment the Fe^{2+} concentration is 0.15 mol dm^{-3} . All other concentrations remain at their standard values.

The Nernst equation is shown.

$$E = E^\ominus + (0.059/z) \log \frac{[\text{oxidised species}]}{[\text{reduced species}]}$$

(i) Use the Nernst equation to calculate the electrode potential, E , for the $\text{Fe}^{3+}/\text{Fe}^{2+}$ half-cell in this experiment.

$$[E^\ominus: \text{Fe}^{3+}/\text{Fe}^{2+} = +0.77\text{V}]$$

$$E = \dots\dots\dots \text{ V [1]}$$

(ii) Use your answer to (d)(i) to calculate E_{cell} for this electrochemical cell.

$$E_{\text{cell}} = \dots\dots\dots \text{ V [1]}$$

[Total: 8]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.022 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)

The Periodic Table of Elements

Group																	
1	2											13	14	15	16	17	18
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">1 H hydrogen 1.0</div> <div style="border: 1px solid black; padding: 5px;">2 He helium 4.0</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">3 Li lithium 6.9</div> <div style="border: 1px solid black; padding: 5px;">4 Be beryllium 9.0</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">5 B boron 10.8</div> <div style="border: 1px solid black; padding: 5px;">6 C carbon 12.0</div> <div style="border: 1px solid black; padding: 5px;">7 N nitrogen 14.0</div> <div style="border: 1px solid black; padding: 5px;">8 O oxygen 16.0</div> <div style="border: 1px solid black; padding: 5px;">9 F fluorine 19.0</div> <div style="border: 1px solid black; padding: 5px;">10 Ne neon 20.2</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">11 Na sodium 23.0</div> <div style="border: 1px solid black; padding: 5px;">12 Mg magnesium 24.3</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">13 Al aluminium 27.0</div> <div style="border: 1px solid black; padding: 5px;">14 Si silicon 28.1</div> <div style="border: 1px solid black; padding: 5px;">15 P phosphorus 31.0</div> <div style="border: 1px solid black; padding: 5px;">16 S sulfur 32.1</div> <div style="border: 1px solid black; padding: 5px;">17 Cl chlorine 35.5</div> <div style="border: 1px solid black; padding: 5px;">18 Ar argon 39.9</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">19 K potassium 39.1</div> <div style="border: 1px solid black; padding: 5px;">20 Ca calcium 40.1</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">21 Sc scandium 45.0</div> <div style="border: 1px solid black; padding: 5px;">22 Ti titanium 47.9</div> <div style="border: 1px solid black; padding: 5px;">23 V vanadium 50.9</div> <div style="border: 1px solid black; padding: 5px;">24 Cr chromium 52.0</div> <div style="border: 1px solid black; padding: 5px;">25 Mn manganese 54.9</div> <div style="border: 1px solid black; padding: 5px;">26 Fe iron 55.8</div> <div style="border: 1px solid black; padding: 5px;">27 Co cobalt 58.9</div> <div style="border: 1px solid black; padding: 5px;">28 Ni nickel 58.7</div> <div style="border: 1px solid black; padding: 5px;">29 Cu copper 63.5</div> <div style="border: 1px solid black; padding: 5px;">30 Zn zinc 65.4</div> <div style="border: 1px solid black; padding: 5px;">31 Ga gallium 69.7</div> <div style="border: 1px solid black; padding: 5px;">32 Ge germanium 72.6</div> <div style="border: 1px solid black; padding: 5px;">33 As arsenic 74.9</div> <div style="border: 1px solid black; padding: 5px;">34 Se selenium 79.0</div> <div style="border: 1px solid black; padding: 5px;">35 Br bromine 83.8</div> <div style="border: 1px solid black; padding: 5px;">36 Kr krypton 83.8</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">37 Rb rubidium 85.5</div> <div style="border: 1px solid black; padding: 5px;">38 Sr strontium 87.6</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">39 Y yttrium 88.9</div> <div style="border: 1px solid black; padding: 5px;">40 Zr zirconium 91.2</div> <div style="border: 1px solid black; padding: 5px;">41 Nb niobium 92.9</div> <div style="border: 1px solid black; padding: 5px;">42 Mo molybdenum 95.9</div> <div style="border: 1px solid black; padding: 5px;">43 Tc technetium —</div> <div style="border: 1px solid black; padding: 5px;">44 Ru ruthenium 101.1</div> <div style="border: 1px solid black; padding: 5px;">45 Rh rhodium 102.9</div> <div style="border: 1px solid black; padding: 5px;">46 Pd palladium 106.4</div> <div style="border: 1px solid black; padding: 5px;">47 Ag silver 107.9</div> <div style="border: 1px solid black; padding: 5px;">48 Cd cadmium 112.4</div> <div style="border: 1px solid black; padding: 5px;">49 In indium 114.8</div> <div style="border: 1px solid black; padding: 5px;">50 Sn tin 118.7</div> <div style="border: 1px solid black; padding: 5px;">51 Sb antimony 121.8</div> <div style="border: 1px solid black; padding: 5px;">52 Te tellurium 127.6</div> <div style="border: 1px solid black; padding: 5px;">53 I iodine 126.9</div> <div style="border: 1px solid black; padding: 5px;">54 Xe xenon 131.3</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">55 Cs caesium 132.9</div> <div style="border: 1px solid black; padding: 5px;">56 Ba barium 137.3</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">57–71 lanthanoids</div> <div style="border: 1px solid black; padding: 5px;">72 Hf hafnium 178.5</div> <div style="border: 1px solid black; padding: 5px;">73 Ta tantalum 180.9</div> <div style="border: 1px solid black; padding: 5px;">74 W tungsten 183.8</div> <div style="border: 1px solid black; padding: 5px;">75 Re rhenium 186.2</div> <div style="border: 1px solid black; padding: 5px;">76 Os osmium 190.2</div> <div style="border: 1px solid black; padding: 5px;">77 Ir iridium 192.2</div> <div style="border: 1px solid black; padding: 5px;">78 Pt platinum 195.1</div> <div style="border: 1px solid black; padding: 5px;">79 Au gold 197.0</div> <div style="border: 1px solid black; padding: 5px;">80 Hg mercury 200.6</div> <div style="border: 1px solid black; padding: 5px;">81 Tl thallium 204.4</div> <div style="border: 1px solid black; padding: 5px;">82 Pb lead 207.2</div> <div style="border: 1px solid black; padding: 5px;">83 Bi bismuth 209.0</div> <div style="border: 1px solid black; padding: 5px;">84 Po polonium —</div> <div style="border: 1px solid black; padding: 5px;">85 At astatine —</div> <div style="border: 1px solid black; padding: 5px;">86 Rn radon —</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">87 Fr francium —</div> <div style="border: 1px solid black; padding: 5px;">88 Ra radium —</div> </div>															
		<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="border: 1px solid black; padding: 5px;">89–103 actinoids</div> <div style="border: 1px solid black; padding: 5px;">104 Rf rutherfordium —</div> <div style="border: 1px solid black; padding: 5px;">105 Db dubnium —</div> <div style="border: 1px solid black; padding: 5px;">106 Sg seaborgium —</div> <div style="border: 1px solid black; padding: 5px;">107 Bh bohrium —</div> <div style="border: 1px solid black; padding: 5px;">108 Hs hassium —</div> <div style="border: 1px solid black; padding: 5px;">109 Mt meitnerium —</div> <div style="border: 1px solid black; padding: 5px;">110 Ds darmstadtium —</div> <div style="border: 1px solid black; padding: 5px;">111 Rg roentgenium —</div> <div style="border: 1px solid black; padding: 5px;">112 Cn copernicium —</div> <div style="border: 1px solid black; padding: 5px;">113 Nh nihonium —</div> <div style="border: 1px solid black; padding: 5px;">114 Fl flerovium —</div> <div style="border: 1px solid black; padding: 5px;">115 Mc moscovium —</div> <div style="border: 1px solid black; padding: 5px;">116 Lv livermorium —</div> <div style="border: 1px solid black; padding: 5px;">117 Ts tennessine —</div> <div style="border: 1px solid black; padding: 5px;">118 Og oganeson —</div> </div>															

lanthanoids	57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.4	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
actinoids	89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.