

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the October/November 2007 question paper

9701 CHEMISTRY

9701/02

Paper 2 (Theory 1), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.


CIE is publishing the mark schemes for the October/November 2007 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



UNIVERSITY of CAMBRIDGE
International Examinations


Page 2	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2007	9701	02

1 (a)




1s

spherical (1)



2s

larger spherical (1)



2p_x

double lobes along the x-axis (1)

[3]

(b) (i) attraction between bonding electrons and nuclei (1)

attraction is electrostatic (1)

(ii) H₂ s-s **overlap** clearly shown

must **not** be normal dot/cross diagram (1)

HCl s-p **overlap** clearly shown

overlap must involve s and p orbitals (1)

[4]

(c) (i) bonding electrons are unequally shared **or**


the molecule has a dipole/ δ^+ and δ^- ends to molecule (1)

(ii) the H and Cl atoms have different electronegativities

or chlorine is more electronegative than hydrogen (1)

[2]

Page 3	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2007	9701	02



allow two 'sausages' above **and** below the C-C axis

or two p orbitals **overlapping** sideways

to form one (localised) π bond over two carbon atoms

(1) [1]

(e) $\Delta H_f^\ominus = 2(-393.7) + 2(-285.9) - (-1411)$


$= + 51.8 \text{ kJ mol}^{-1}$ (units given in qu.)

(3)

penalise errors: no 2 for -393.7
no 2 for -285.9
wrong sign for $-(-1411)$

[3]

[Total: 13]



equation

(1)

state symbols

(1) [2]

(b) (i) giant ionic lattice (may be in diag.)

(1)

strong ionic bonds

(1)

(ii) simple molecular **or** discrete molecules

(may be shown in a diagram)




(1)

with **weak** intermolecular forces **or**

weak van der Waals' forces

between them

(1) [4]



(1) [1]

Page 4	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2007	9701	02


- (d) NaCl pH is 7 allow neutral (1)
- PCl_5 pH is between 1 and 4
- do **not** allow acidic (1) [2]

- (e) (i) 460 K Al_2Cl_6 (1)
- 1150 K AlCl_3 (1)

(ii) correct **dot-and-cross** diagram for AlCl_3 (1)

(iii) correct displayed structure for Al_2Cl_6 (1)

two correct co-ordinate bonds (1)



[5]

[Total: 14]

- 3 (a) P_4 (1)
- S_8 (1)
- Cl_2 (1) [3]

(b) (i) highest S_8 P_4 Cl_2 lowest (1)

allow S ... P ... Cl **or** names (1)

(ii) from S_8 to P_4 to Cl_2

there are fewer electrons in each molecule (1)

hence weaker van der Waals' forces (1) [3]


Page 5	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2007	9701	02

(c) (i) $S_2Cl_2 = (2 \times 32.1) + (2 \times 35.5) = 135.2$

$$n(S_2Cl_2) = \frac{2.7}{135.2} = 0.0199 = 0.02 \quad (1)$$

$$0.02 \text{ mol } S_2Cl_2 \rightarrow \frac{0.96}{32.1} = 0.03 \text{ mol S}$$

$$1.0 \text{ mol } S_2Cl_2 \rightarrow \frac{0.03 \times 1.0}{0.02} = 1.5 \text{ mol S} \quad (1)$$



correct products (1)


balanced equation (1) [4]

(d) oxidation product is H_2SO_3 (1)

reduction product is S (1) [2]


[Total: 12]

4 (a)



H atoms must be shown.

Structure must not contain any CH_3 groups (1) [1]





(1) [2]

(c) $CH_3CH(OH)CH_2CH_2CH_3$ (1)

$CH_3CH_2CH(OH)CH_2CH_3$ (1) [2]


Page 6	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2007	9701	02

- (d)  correct compound (1)
- correct mirror object/mirror image relationship in 3D (1) [2]

- (e)  e.g. cyclopentane structure

allow methylcyclobutane **or** dimethylcyclopropane (1) [1]

- (f) e.g.



two repeat units must be shown
relative positions of $-\text{CH}_3$ and $-\text{C}_2\text{H}_5$ may differ from those shown above (1) [1]

[Total: 9]

- 5 (a) (i) $\text{Cr}_2\text{O}_7^{2-}/\text{H}^+$ allow $\text{MnO}_4^-/\text{H}^+$ (1)
- (ii) from orange to **or** purple to colourless
green **or** green/blue (1) [2]
- (b) (i) to ensure complete oxidation of $-\text{CH}_2\text{OH}$
or to keep reactants in the reaction flask (1)
- (ii) CH_3CHO /ethanal (1) [2]
- (c) (i) CH_3I /iodomethane (1)
- (ii) nucleophilic substitution **or** hydrolysis (1) [2]

Page 7	Mark Scheme	Syllabus	Paper
	GCE A/AS LEVEL – October/November 2007	9701	02

(d) step I

red P + I₂ **or** HI(aq) **or** KBr/conc H₃PO₄ **or** PI₃ (1)

heat **but** room temperature for PI₃ (1)

step II

KCN in aqueous ethanol (1)

in aqueous ethanol, heat under reflux (1)

allow aqueous ethanol in either place

step III

aqueous mineral acid (**not** nitric acid)

or NaOH(aq) then aqueous mineral acid (1)

heat (1) [6]

[Total: 12]