

# Cambridge International AS & A Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 9 2 2 8 1 7 8 8 4 8 \*

**MATHEMATICS****9709/62**

Paper 6 Probability &amp; Statistics 2

**February/March 2021****1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

**INSTRUCTIONS**

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

**INFORMATION**

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

---

This document has **12** pages. Any blank pages are indicated.









3 An architect wishes to investigate whether the buildings in a certain city are higher, on average, than buildings in other cities. He takes a large random sample of buildings from the city and finds the mean height of the buildings in the sample. He calculates the value of the test statistic,  $z$ , and finds that  $z = 2.41$ .

(a) Explain briefly whether he should use a one-tail test or a two-tail test. [1]

.....  
.....  
.....

(b) Carry out the test at the 1% significance level. [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

4 On average, 1 in 400 microchips made at a certain factory are faulty. The number of faulty microchips in a random sample of 1000 is denoted by  $X$ .

(a) State the distribution of  $X$ , giving the values of any parameters. [1]

.....  
.....  
.....

(b) State an approximating distribution for  $X$ , giving the values of any parameters. [2]

.....  
.....  
.....

(c) Use this approximating distribution to find each of the following.

(i)  $P(X = 4)$ . [2]

.....  
.....  
.....  
.....

(ii)  $P(2 \leq X \leq 4)$ . [2]

.....  
.....  
.....  
.....  
.....  
.....

(d) Use a suitable approximating distribution to find the probability that, in a random sample of 700 microchips, there will be at least 1 faulty one. [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....







6 It is known that 8% of adults in a certain town own a Chantor car. After an advertising campaign, a car dealer wishes to investigate whether this proportion has increased. He chooses a random sample of 25 adults from the town and notes how many of them own a Chantor car.

(a) He finds that 4 of the 25 adults own a Chantor car.

Carry out a hypothesis test at the 5% significance level. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Explain which of the errors, Type I or Type II, might have been made in carrying out the test in part (a). [2]

.....

.....

.....

.....

.....

.....

.....

Later, the car dealer takes another random sample of 25 adults from the town and carries out a similar hypothesis test at the 5% significance level.

- (c) Find the probability of a Type I error. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

