



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



MATHEMATICS

9709/32

Paper 3 Pure Mathematics 3

October/November 2021

1 hour 50 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 75.
- The number of marks for each question or part question is shown in brackets [].

This document has **20** pages. Any blank pages are indicated.

- 1 Find the value of x for which $3(2^{1-x}) = 7^x$. Give your answer in the form $\frac{\ln a}{\ln b}$, where a and b are integers. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 3 (a) Given the complex numbers $u = a + ib$ and $w = c + id$, where a , b , c and d are real, prove that $(u + w)^* = u^* + w^*$. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Solve the equation $(z + 2 + i)^* + (2 + i)z = 0$, giving your answer in the form $x + iy$ where x and y are real. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 5 (a) On a sketch of an Argand diagram, shade the region whose points represent complex numbers z satisfying the inequalities $|z - 3 - 2i| \leq 1$ and $\text{Im } z \geq 2$. [4]

- (b) Find the greatest value of $\arg z$ for points in the shaded region, giving your answer in degrees. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6 (a) Using the expansions of $\sin(3x + 2x)$ and $\sin(3x - 2x)$, show that

$$\frac{1}{2}(\sin 5x + \sin x) \equiv \sin 3x \cos 2x. \quad [3]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

10 With respect to the origin O , the position vectors of the points A and B are given by $\vec{OA} = \begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$ and $\vec{OB} = \begin{pmatrix} 0 \\ 3 \\ 1 \end{pmatrix}$.

(a) Find a vector equation for the line l through A and B . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) The point C lies on l and is such that $\vec{AC} = 3\vec{AB}$.
Find the position vector of C . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

11 The equation of a curve is $y = \sqrt{\tan x}$, for $0 \leq x < \frac{1}{2}\pi$.

(a) Express $\frac{dy}{dx}$ in terms of $\tan x$, and verify that $\frac{dy}{dx} = 1$ when $x = \frac{1}{4}\pi$. [4]

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

The value of $\frac{dy}{dx}$ is also 1 at another point on the curve where $x = a$, as shown in the diagram.



(b) Show that $t^3 + t^2 + 3t - 1 = 0$, where $t = \tan a$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Use the iterative formula

$$a_{n+1} = \tan^{-1} \left(\frac{1}{3}(1 - \tan^2 a_n - \tan^3 a_n) \right)$$

to determine a correct to 2 decimal places, giving the result of each iteration to 4 decimal places. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

