

CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Level

MARK SCHEME for the October/November 2012 series

9231 FURTHER MATHEMATICS

9231/13

Paper 1, maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol ∇ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through" marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

Qu No	Commentary	Solution	Marks	Part Marks	Total
1	Use of: Use of: Obtains result.	$\sum_{n+1}^{2n} = \sum_1^{2n} - \sum_1^n$ $\sum_1^n r^2 = \frac{n(n+1)(2n+1)}{6}$ $\frac{2n(2n+1)(4n+1)}{6} - \frac{n(n+1)(2n+1)}{6}$ $= \frac{1}{6}n(2n+1)(8n+2-n-1) = \frac{1}{6}n(2n+1)(7n+1) \text{ (AG)}$	M1 M1 A1 A1	4	[4]
2	Sets determinant $\neq 0$. Factorises, or completes square. States result.	$\begin{vmatrix} a & 1 & 2 \\ 3 & -2 & 0 \\ 3 & -4 & -6a \end{vmatrix} \neq 0 \Rightarrow 12a^2 + 18a - 12 \neq 0$ $\Rightarrow 6(2a-1)(a+2) \neq 0$ $a \neq \frac{1}{2} \text{ or } -2 \text{ (Or by row operations.)}$	M1A1 M1 A1	4	[4]
3	Proposition. Proves base case. States inductive hypothesis. Proves inductive step. States conclusion.	$H_N : S_N = 1 - \frac{1}{(N+1)!}$ $S_1 = \frac{1}{2!} = \frac{1}{2} = 1 - \frac{1}{2!} \Rightarrow H_1 \text{ is true.}$ $H_k : \text{ Assume } S_k = 1 - \frac{1}{(k+1)!} \text{ is true.}$ $\Rightarrow S_{k+1} = 1 - \frac{1}{(k+1)!} + \frac{k+1}{(k+2)!} = \frac{(k+2)! - (k+2) + (k+1)}{(k+2)!}$ $\Rightarrow S_{k+1} = 1 - \frac{1}{(k+2)!} \therefore H_k \Rightarrow H_{k+1}.$ $\therefore \text{ (By PMI } H_n \text{ is) true for all positive integers } N.$	B1 B1 M1 A1 A1	5	[5]
4	Finds vector product. Finds area of triangle. Finds length of perpendicular.	$\vec{AB} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \quad \vec{AC} = \begin{pmatrix} 1 \\ 1 \\ 2 \end{pmatrix}$ $\vec{AB} \times \vec{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 3 \\ 1 & 1 & 2 \end{vmatrix} = \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix}$ $\text{Area of triangle } ABC = \frac{1}{2} \left \begin{pmatrix} 1 \\ 1 \\ -1 \end{pmatrix} \right = \frac{1}{2} \sqrt{3}$ $\frac{1}{2} \sqrt{1^2 + 2^2 + 3^2} d = \frac{1}{2} \sqrt{3} \Rightarrow d = \sqrt{\frac{3}{14}}$	B1 M1A1 M1A1 A1	3	[6]

Page 5	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

Qu No	Commentary	Solution	Marks	Part Marks	Total
5	<p>Sketches graph.</p> <p>Uses area of sector formula.</p> <p>Uses double angle formula Integrates.</p> <p>Obtains result.</p>	<p>Correct shape and orientation. Passing through (0,0) and (3,0)</p> $\text{Area} = 2 \times \frac{1}{2} \int_0^{\frac{\pi}{3}} (1 + 2 \cos \theta)^2 d\theta$ $= \int_0^{\frac{\pi}{3}} (1 + 4 \cos \theta + 4 \cos^2 \theta) d\theta$ $= \int_0^{\frac{\pi}{3}} (3 + 4 \cos \theta + 2 \cos 2\theta) d\theta$ $= [3\theta + 4 \sin \theta + \sin 2\theta]_0^{\frac{\pi}{3}}$ $= \left[\pi + \frac{5\sqrt{3}}{2} \right]$	<p>B1 B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>A1</p>	<p>2</p> <p>4</p>	<p>[6]</p>
6	<p>Differentiates.</p> <p>Obtains $\left(\frac{ds}{dt}\right)^2$</p> <p>Uses surface area formula about <u>y-axis</u>.</p>	$\dot{x} = 2t \quad \dot{y} = t^3 - \frac{1}{t}$ $(\dot{x})^2 + (\dot{y})^2 = 4t^2 + t^6 - 2t^2 + \frac{1}{t^2} = \left(t^3 + \frac{1}{t}\right)^2$ $S = \int 2\pi x ds = 2\pi \int_1^2 t^2 \left(t^3 + \frac{1}{t}\right) dt = 2\pi \int_1^2 (t^5 + t) dt$ $= 2\pi \left[\frac{1}{6} t^6 + \frac{1}{2} t^2 \right]_1^2 = 2\pi \left\{ \left[\frac{32}{3} + 2 \right] - \left[\frac{1}{6} + \frac{1}{2} \right] \right\} = 24\pi$	<p>B1</p> <p>M1A1</p> <p>M1A1</p> <p>M1A1</p>	<p>7</p>	<p>[7]</p>
7		$(\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha) = \alpha^2 + \beta^2 + \gamma^2 \Rightarrow \sum \alpha\beta = 1$ <p>Either</p> <p>Required equation is $x^3 - 4x^2 + x + c = 0$</p> $\Rightarrow \sum \alpha^3 - 4 \sum \alpha^2 + 4 + 3c = 0$ $\Rightarrow 3c = 56 - 34 - 4 = 18 \Rightarrow c = 6 \quad (\text{AG})$ <p>Or</p> $\alpha^3 + \beta^3 + \gamma^3 - 3\alpha\beta\gamma = (\alpha + \beta + \gamma)(\alpha^2 + \beta^2 + \gamma^2 - \alpha\beta - \beta\gamma - \gamma\alpha)$ <p>(or some other appropriate identity, e.g.</p> $(\alpha + \beta + \gamma)^3 = \alpha^3 + \beta^3 + \gamma^3 + 3(\alpha + \beta + \gamma)(\alpha\beta + \beta\gamma + \gamma\alpha) - 3\alpha\beta\gamma$ $\Rightarrow \dots \Rightarrow \alpha\beta\gamma = -6$ $\Rightarrow x^3 - 4x^2 + x + 6 = 0 \quad (\text{AG})$ $\Rightarrow (x+1)(x-2)(x-3) = 0 \Rightarrow x = -1, 2, 3.$	<p>M1A1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>(M1)</p> <p>(M1A1)</p> <p>A1</p> <p>M1A1</p>	<p>2</p> <p>6</p>	<p>[8]</p>

Page 6	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

Qu No	Commentary	Solution	Marks	Part Marks	Total
8	<p>Re-write.</p> <p>Obtains result.</p> <p>Use of Bin. Thm.</p> <p>Takes imaginary part and uses de M.</p> <p>Applies initial result.</p> <p>Equates imaginary parts to obtain result.</p>	$1 + z = 2 \cos^2 \frac{1}{2} \theta + 2i \sin \frac{1}{2} \theta \cos \frac{1}{2} \theta$ $= 2 \cos \frac{1}{2} \theta \left(\cos \frac{1}{2} \theta + i \sin \frac{1}{2} \theta \right) \quad (\text{AG})$ $(1 + z)^n = 1 + \binom{n}{1} z + \binom{n}{2} z^2 + \dots + \binom{n}{n} z^n$ $\therefore \text{Im}(1 + z)^n = \binom{n}{1} \sin \theta + \binom{n}{2} \sin 2\theta + \dots + \binom{n}{n} \sin n\theta$ <p>But $(1 + z)^n = 2^n \cos^n \frac{1}{2} \theta e^{i \frac{n}{2} \theta}$</p> $\therefore \binom{n}{1} \sin \theta + \binom{n}{2} \sin 2\theta + \dots + \binom{n}{n} \sin n\theta$ $= 2^n \cos^n \frac{1}{2} \theta \sin \frac{n}{2} \theta$	<p>M1A1</p> <p>A1</p> <p>M1</p> <p>M1</p> <p>M1A1</p> <p>B1</p> <p>A1</p>	<p>3</p> <p>6</p>	<p>[9]</p>
9	<p>States vertical asymptote.</p> <p>Divides and states oblique asymptote.</p> <p>Rearranges as quadratic in x.</p> <p>Uses discriminant to obtain condition stated.</p> <p>Differentiates, puts $= 0$ and obtains x-values.</p> <p>States stationary points.</p> <p>Sketch.</p>	<p>Vertical asymptote is $x = 2$.</p> <p>\Rightarrow oblique asymptote is $y = x - 1$.</p> $xy - 2y = x^2 - 3x + 3 \Rightarrow x^2 - (y + 3)x + (3 + 2y) = 0$ <p>For real x, $B^2 - 4AC \geq 0$</p> $\therefore (y + 3)^2 - 4(3 + 2y) \geq 0$ $\Rightarrow \dots \Rightarrow (y - 3)(y + 1) \geq 0$ $\Rightarrow y \leq -1 \text{ or } y \geq 3$ <p>\therefore no points for $-1 < y < 3$ (AG)</p> $y' = 1 - (x - 2)^{-2} = 0 \Rightarrow x = 1 \text{ or } 3$ <p>(Or uses $y = -1$ and 3 to obtain x-values.)</p> <p>Stationary points are $(1, -1)$ and $(3, 3)$</p> <p>One mark for each branch correctly placed.</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>M1</p> <p>A1A1</p> <p>B1B1</p>	<p>3</p> <p>4</p> <p>3</p> <p>2</p>	<p>[12]</p>

Page 7	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

Qu No	Commentary	Solution	Marks	Part Marks	Total
10	Differentiates implicitly.	$3x^2 + 3y^2 y' = 3y + 3xy'$	B1B1	4	
	Equates $\frac{dy}{dx}$ to zero.	$\Rightarrow \frac{dy}{dx} = \frac{y - x^2}{y^2 - 1} = 0$	M1		
	Obtains relationship.	$\Rightarrow y = x^2$	A1		
	Substitutes for y.	$\Rightarrow xy + y^3 = 3xy \Rightarrow y^3 = 2xy \Rightarrow y^2 = 2x \quad (y \neq 0)$	M1		
	Obtains x and y.	$\Rightarrow x^4 = 2x \Rightarrow x^3 = 2 \Rightarrow x = 2^{\frac{1}{3}}$ and $\Rightarrow y = 2^{\frac{2}{3}} \quad (x \neq 0)$	A1A1		
	Differentiates	$6x + 3y^2 y'' + 6y(y')^2 = 3y' + 3xy'' + 3y'$	B1	8	
	Uses $y' = 0$.	$\Rightarrow 6x = y''(3x - 3y^2) \Rightarrow y'' = \frac{2x}{x(1 - x^3)}$	B1B1 M1		
	Identifies maximum.	$\Rightarrow y'' = \frac{2}{1 - 2} = -2 \Rightarrow \text{max}$	A1		
	(Other watertight methods for showing a maximum accepted.)				[12]

Page 8	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

Qu No	Commentary	Solution	Marks	Part Marks	Total
11	Verifies result.	$\frac{d}{dx} \left(-\frac{1}{3}(1-x^2)^{\frac{3}{2}} + c \right) = -\frac{1}{3} \times \frac{3}{2} (1-x^2)^{\frac{1}{2}} \times (-2x) = x(1-x^2)^{\frac{1}{2}}$	B1	1	
	Correct parts.		M1		
	Integrates by parts.	$I_n = \int_0^1 x^n (1-x^2)^{\frac{1}{2}} dx = \int_0^1 x^{n-1} \cdot x(1-x^2)^{\frac{1}{2}} dx$			
	Substitutes limits.	$= \left[-x^{n-1} \cdot \frac{1}{3} (1-x^2)^{\frac{3}{2}} \right]_0^1 + \int_0^1 (n-1)x^{n-2} \cdot \frac{1}{3} (1-x^2)^{\frac{3}{2}} dx$	M1A1		
	Obtains reduction formula.	$= \frac{n-1}{3} \int_0^1 x^{n-2} (1-x^2)(1-x^2)^{\frac{1}{2}} dx$ $= \frac{n-1}{3} I_{n-2} - \frac{n-1}{3} I_n$ $\Rightarrow (n+2)I_n = (n-1)I_{n-2} \quad (\text{AG})$	M1	5	
	Uses substitution correctly.	$x = \sin u \Rightarrow dx = \cos u du$ Limits: $x = 0 \Rightarrow u = 0 \quad x = 1 \Rightarrow u = \frac{\pi}{2}$	M1		
	Uses double angle formula.	$\int_0^1 (1-x^2)^{\frac{1}{2}} dx = \int_0^{\frac{\pi}{2}} \cos^2 u du$	A1		
	Integrates correctly.	$= \int_0^{\frac{\pi}{2}} \frac{1}{2} (\cos 2u + 1) du$	M1		
	Uses reduction formula correctly.	$= \frac{1}{2} \left[\frac{\sin 2u}{2} + u \right]_0^{\frac{\pi}{2}} = \frac{\pi}{4} \quad (\text{AG})$	M1A1	5	
		$\Rightarrow I_2 = \frac{1}{4} \times \frac{\pi}{4} = \frac{\pi}{16} \Rightarrow I_4 = \frac{1}{2} \times \frac{\pi}{16} = \frac{\pi}{32}$	M1A1		

Page 9	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

Qu No	Commentary	Solution	Marks	Part Marks	Total
12	EITHER				
	Use of these results.	$\mathbf{Ae} = \lambda \mathbf{e}$ and $\mathbf{Be} = \mu \mathbf{e}$ $\mathbf{ABe} = \mathbf{A}\mu \mathbf{e} = \mu \mathbf{Ae} = \mu \lambda \mathbf{e} = \lambda \mu \mathbf{e}$	M1 A1	2	
	Finds missing eigenvalues of A .	$\begin{pmatrix} 3 & 2 & 2 \\ -2 & -2 & -2 \\ 1 & 2 & 2 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \Rightarrow \lambda = 0$	B1		
		$\begin{pmatrix} 3 & 2 & 2 \\ -2 & -2 & -2 \\ 1 & 2 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} \Rightarrow \lambda = 1$	B1	2	
	Finds missing eigenvector of A .	$\lambda = 2 \Rightarrow \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 2 \\ -2 & -4 & -2 \end{vmatrix} = \begin{pmatrix} 4 \\ -2 \\ 0 \end{pmatrix} \sim \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix}$	M1A1	2	
	Calculates eigenvectors of B	$\begin{pmatrix} -1 & 2 & 2 \\ 2 & 2 & 2 \\ -3 & -6 & -6 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \Rightarrow \mu = 0$	B1		
		$\begin{pmatrix} -1 & 2 & 2 \\ 2 & 2 & 2 \\ -3 & -6 & -6 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix} = \begin{pmatrix} -3 \\ 0 \\ 3 \end{pmatrix} \Rightarrow \mu = -3$	B1		
		$\begin{pmatrix} -1 & 2 & 2 \\ 2 & 2 & 2 \\ -3 & -6 & -6 \end{pmatrix} \begin{pmatrix} 2 \\ -1 \\ 0 \end{pmatrix} = \begin{pmatrix} -4 \\ 2 \\ 0 \end{pmatrix} \Rightarrow \mu = -2$	B1		
	Uses initial result to find eigenvalues of C . (1 mark for one correct value, 2 marks for all three.)	$\therefore \mathbf{C}$ has eigenvalues: $0 \times 0 = 0 \quad 1 \times (-3) = -3 \quad 2 \times (-2) = -4$	B2,1,0		
	Finds P and D .	$\mathbf{P} = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 0 & -1 \\ -1 & -1 & 0 \end{pmatrix} \text{ (OE) } \mathbf{D} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 16 \end{pmatrix}$	B1 M1A1	8	[14]

Page 10	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – October/November 2012	9231	13

Qu No	Commentary	Solution	Marks	Part Marks	Total
12	OR				
	Finds complementary function.	$m^2 + 6m + 13 = 0 \Rightarrow m = -3 \pm 2i$ $x = e^{-3t}(A \cos 2t + B \sin 2t)$	M1 A1		
	Finds particular integral.	$x = p \cos 2t + q \sin 2t$ $\dot{x} = -2p \sin 2t + 2q \cos 2t$ $\ddot{x} = -4p \cos 2t - 4q \sin 2t$	M1		
		$(9p + 12q) \cos 2t + (9q - 12p) \sin 2t = 75 \cos 2t$ $\Rightarrow p = 3 \quad q = 4$	M1A1 A1		
	Finds general solution.	$x = e^{-3t}(A \cos 2t + B \sin 2t) + 3 \cos 2t + 4 \sin 2t$	A1 B1	7	
	Uses initial conditions..	$x = 5$ when $t = 0 \Rightarrow 5 = A + 3 \Rightarrow A = 2$ $\dot{x} = -3e^{-3t}(A \cos 2t + B \sin 2t)$ $+ e^{-3t}(-2A \sin 2t + 2B \cos 2t) - 6 \sin 2t + 8 \cos 2t$ $\dot{x} = 0$ when $t = 0 \Rightarrow 0 = -6 + 8 + 2B \Rightarrow B = -1$	M1 A1		
	Obtains solution.	$x = e^{-3t}(2 \cos 2t - \sin 2t) + 3 \cos 2t + 4 \sin 2t$	A1	4	
	Obtains limit.	As $t \rightarrow \infty$, $e^{-3t} \rightarrow 0$ $\therefore x \approx 3 \cos 2t + 4 \sin 2t$ $\therefore x \approx 5 \left(\frac{3}{5} \cos 2t + \frac{4}{5} \sin 2t \right) = 5 \cos \left(2t - \tan^{-1} \frac{4}{3} \right)$ (AG)	M1 M1A1	3	[14]