
BIOLOGY

0610/51

Paper 5 Practical Test

October/November 2017

MARK SCHEME

Maximum Mark: 40

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is a registered trademark.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **9** printed pages.

Mark schemes will use these abbreviations

- ; separates marking points
- / alternatives
- I ignore
- R reject
- A accept (for answers correctly cued by the question, or guidance for examiners)
- AW alternative wording (where responses vary more than usual)
- AVP any valid point
- **ecf** credit a correct statement / calculation that follows a previous wrong response
- **ora** or reverse argument
- () the word / phrase in brackets is not required, but sets the context
- underline actual word given must be used by candidate (grammatical variants excepted)
- max indicates the maximum number of marks that can be given

Question	Answer	Marks	Guidance
1(a)(i)	one table drawn with appropriate number of columns ; correct column and row headings with appropriate units (pH and intensity / minutes) ; pH recorded for each solution ; colour of apple slices recorded by intensity for 0 and 10 minutes (numerical values not words) ; colour of apple slices recorded by intensity for 20 minutes (numerical values not words) ; suitable trend ;	6	I control if added to table R if units in data cell I units in data cells A if the correct data is not linked to a time
1(a)(ii)	2 and/or 3, 7, then 8 and/or 9 ;	1	A B and C in either order, A, D and E in either order
1(a)(iii)	to compare (what happens to the apple) with the solution and with no solution / at different pH values and no pH / with different pH values and the air / with the solution and with the air ;	1	I for comparison unqualified A to see what happens to the crushed apple when it is not exposed to the solution A to observe what would happen without the effect of pH A idea that the control shows that the liquid (or change in pH) is causing the results (and not the air)
1(a)(iv)	lemon juice ; has low pH/is acidic/(lemon juice has a pH of 2 and) previous experiment showed that apple won't go brown at pH2 ;	2	A olive oil A (olive oil) keeps out air/oxygen A enzyme doesn't work / denatured at pH2

Question	Answer	Marks	Guidance												
1(b)(i)	<table border="1"> <thead> <tr> <th data-bbox="309 213 741 277"><i>variable</i></th> <th data-bbox="748 213 1173 277"><i>controlled by</i></th> </tr> </thead> <tbody> <tr> <td data-bbox="309 282 741 416">volume / amount / mass, of solution / named solution / liquid</td> <td data-bbox="748 282 1173 416">adding 20 cm³</td> </tr> <tr> <td data-bbox="309 421 741 517">(same / one) apple</td> <td data-bbox="748 421 1173 517">all slices cut from the same apple</td> </tr> <tr> <td data-bbox="309 521 741 585">time soaking in solution</td> <td data-bbox="748 521 1173 585">all soaked for 2 mins</td> </tr> <tr> <td data-bbox="309 590 741 718">observation time / time intervals</td> <td data-bbox="748 590 1173 718">left for 20 min / results checked every 10 minutes</td> </tr> <tr> <td data-bbox="309 722 741 754">;</td> <td data-bbox="748 722 1173 754">;</td> </tr> </tbody> </table>	<i>variable</i>	<i>controlled by</i>	volume / amount / mass, of solution / named solution / liquid	adding 20 cm ³	(same / one) apple	all slices cut from the same apple	time soaking in solution	all soaked for 2 mins	observation time / time intervals	left for 20 min / results checked every 10 minutes	;	;	2	<p>1 mark for the variable, 1 mark for method of controlling which must related</p> <p>I temperature</p> <p>I time unqualified unless explanation clarifies</p> <p>A time apple was left in the Petri dish A oxygen / air exposure time</p>
<i>variable</i>	<i>controlled by</i>														
volume / amount / mass, of solution / named solution / liquid	adding 20 cm ³														
(same / one) apple	all slices cut from the same apple														
time soaking in solution	all soaked for 2 mins														
observation time / time intervals	left for 20 min / results checked every 10 minutes														
;	;														
1(b)(ii)	oxygen is needed (from the air) for the reaction ;	1	<p>A to expose the apple to oxygen I air</p>												

Question	Answer	Marks	Guidance												
1(c)	<i>any 1 from:</i> idea that it is a qualitative / subjective, method / judged by eye / similar browning looks the same / AW ;	1	A there are more than three shades of brown												
1(d)	<p><i>any pair from:</i></p> <table border="1" data-bbox="309 416 1171 1082"> <thead> <tr> <th data-bbox="309 416 741 480"><i>error</i></th> <th data-bbox="741 416 1171 480"><i>improvement</i></th> </tr> </thead> <tbody> <tr> <td data-bbox="309 480 741 647">amount of apple not the same / cutting inaccurate</td> <td data-bbox="741 480 1171 647">use same mass / weight / use a set volume of crushed apple</td> </tr> <tr> <td data-bbox="309 647 741 783">crushing uneven</td> <td data-bbox="741 647 1171 783">use a blender / mortar and pestle</td> </tr> <tr> <td data-bbox="309 783 741 879">idea some may have more time in solution than others</td> <td data-bbox="741 783 1171 879">test each separately</td> </tr> <tr> <td data-bbox="309 879 741 983">some apple exposed to air while others being crushed</td> <td data-bbox="741 879 1171 983">test each separately</td> </tr> <tr> <td data-bbox="309 983 741 1082">only one slice of apple per solution</td> <td data-bbox="741 983 1171 1082">have at least two more replicates</td> </tr> </tbody> </table> <p style="text-align: center;">; ;</p>	<i>error</i>	<i>improvement</i>	amount of apple not the same / cutting inaccurate	use same mass / weight / use a set volume of crushed apple	crushing uneven	use a blender / mortar and pestle	idea some may have more time in solution than others	test each separately	some apple exposed to air while others being crushed	test each separately	only one slice of apple per solution	have at least two more replicates	2	A (cut to the same size) by using a cutter / ruler, slicer, cutting instrument / AW
<i>error</i>	<i>improvement</i>														
amount of apple not the same / cutting inaccurate	use same mass / weight / use a set volume of crushed apple														
crushing uneven	use a blender / mortar and pestle														
idea some may have more time in solution than others	test each separately														
some apple exposed to air while others being crushed	test each separately														
only one slice of apple per solution	have at least two more replicates														

Question	Answer	Marks	Guidance
1(e)	<p>1 ref to using at least three temperatures ;</p> <p>2 stated temperatures or a description ;</p> <p>3 method described to maintain temperature(s) ;</p> <p>4 methodology described e.g. ref to leaving the enzyme and substrate separately to reach a set temperature ;</p> <p>5 ref to constant pH/ experiment carried out at the optimum pH;</p> <p>6 and 7 <i>controlled variables any two from:</i> same concentration of enzyme / same concentration of substrate / same volume of enzyme / same volume of substrate ;;</p> <p>8 leaving for 20 minutes / leaving for a set time / check at stated time intervals ;</p> <p>9 ref to substrate at optimum temperature turning brown first or having the highest colour intensity value ;</p> <p>10 repeat (at least) twice ;</p> <p>11 AVP e.g. repeat the investigation at temperatures near the optimum to obtain a more accurate optimum temperature ;</p>	6	<p>A ref. to measuring time for brown colour to appear</p> <p>A optimum temperature shows the darkest colour</p>

Question	Answer	Marks	Guidance
1(f)(i)	<p>A(xes) – labelled with units (time / minutes and percentage of enzyme activity remaining) ;</p> <p>S(cale) – suitable, even scale and data occupies at least half the grid ;</p> <p>P(lot) – all points plotted accurately $\pm 1/2$ square ;</p> <p>L(ines) – each line drawn with a ruler point to point or smoothed line ;</p> <p>Lines labelled or a key shown ;</p>	5	
1(f)(ii)	<p>for both fruits the enzyme activity decreased as time went on ;</p> <p>the apricot enzyme shows the greatest reduction after 15 minutes ;</p> <p>avocado retains enzyme activity for longer after heating than the apricot ;</p>	1	A apricot enzymes are the most easily destroyed by heat / denatured faster / ora ;

Question	Answer	Marks	Guidance
2(a)(i)	<p>red blood cells do not have a nucleus / white blood cells have a nucleus ;</p> <p>red blood cells, have a light area in the centre / are biconcave ;</p> <p>there are more red blood cells / fewer white blood cells ;</p> <p>red blood cells contents not granular AW / white blood cells contents granular ;</p> <p>red blood cells smaller than phagocytes / ora ;</p> <p>red blood cells are overlapping / white blood cells do not overlap ;</p>	2	
2(a)(ii)	<p>O single clear lines on both cell membranes <u>and</u> no shading in the nucleus ;</p> <p>S monocyte larger than 2.5 cm and neutrophil larger than 2 cm provided ;</p> <p>D1 monocyte is larger than the neutrophil ;</p> <p>D2 neutrophil nucleus has two distinct parts joined by a narrow section, larger part at least twice the size of the smaller part ;</p>	4	

Question	Answer			Marks	Guidance												
2(b)(i)	<table border="1"> <thead> <tr> <th data-bbox="327 218 602 317">type of blood cell</th> <th data-bbox="609 218 880 317">diameters / mm</th> <th data-bbox="887 218 1158 317">average diameter / mm</th> </tr> </thead> <tbody> <tr> <td data-bbox="327 322 602 421">red blood cell</td> <td data-bbox="609 322 880 421">13±1 12±1</td> <td data-bbox="887 322 1158 421">12.5±1</td> </tr> <tr> <td data-bbox="327 426 602 525">lymphocyte</td> <td data-bbox="609 426 880 525">12±1 11±1</td> <td data-bbox="887 426 1158 525">11.5±1</td> </tr> <tr> <td data-bbox="327 529 602 628">phagocyte</td> <td data-bbox="609 529 880 628">22±1 18±1</td> <td data-bbox="887 529 1158 628">20.0±1</td> </tr> </tbody> </table> <p style="text-align: right;">;;;</p>			type of blood cell	diameters / mm	average diameter / mm	red blood cell	13±1 12±1	12.5±1	lymphocyte	12±1 11±1	11.5±1	phagocyte	22±1 18±1	20.0±1	3	mp1 for header units mp2 for six measurements mp3 for three correct averages from candidates results
type of blood cell	diameters / mm	average diameter / mm															
red blood cell	13±1 12±1	12.5±1															
lymphocyte	12±1 11±1	11.5±1															
phagocyte	22±1 18±1	20.0±1															
2(b)(ii)	8 or 9 (∞m) ;;;			3	A ecf from 2(b)(i)												