



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
International General Certificate of Secondary Education

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 8 7 7 2 0 7 8 0 2 1 \*

**CHEMISTRY**

**0620/31**

Paper 3 (Extended)

**October/November 2013**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 16.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of **14** printed pages and **2** blank pages.



1 For each of the following, name an element which matches the description.

(a) It is used as a fuel in nuclear reactors.

..... [1]

(b) It is the only non-metal which is a good conductor of electricity.

..... [1]

(c) Inert electrodes are made from this metal.

..... [1]

(d) This gaseous element is used to fill balloons in preference to hydrogen.

..... [1]

(e) An element which can form an ion of the type  $X^{3-}$ .

..... [1]

(f) It has the same electron distribution as the calcium ion,  $Ca^{2+}$ .

..... [1]

(g) The element is in Period 5 and Group VI.

..... [1]

[Total: 7]

2 (a) Give **three** differences in physical properties between the Group I metal, potassium, and the transition element, iron.

- 1. ....
- 2. ....
- 3. .... [3]

(b) The following metals are in order of reactivity.


potassium  
zinc  
copper

For those metals which react with water or steam, name the products of the reaction, otherwise write 'no reaction'.

- potassium .....
- .....
- zinc .....
- .....
- copper .....
- ..... [5]

[Total: 8]

- 3 Ammonia is manufactured by the Haber process.



The forward reaction is exothermic.

- (a) Describe how the reactants are obtained.

- (i) Nitrogen


.....  
 ..... [2]

- (ii) Hydrogen

.....  
 .....  
 ..... [3]

- (b) The percentage of ammonia in the equilibrium mixture varies with temperature and pressure.

- (i) Which pair of graphs, **A**, **B** or **C**, shows correctly how the percentage of ammonia at equilibrium varies with temperature and pressure?



The pair with **both graphs correct** is ..... [1]

(ii) Give a full explanation of why the pair of graphs you have chosen in (i) is correct.

.....

.....

.....

.....

.....

.....

.....

.....

..... [6]

(iii) Catalysts do not alter the position of equilibrium. Explain why a catalyst is used in this process.

.....


.....

.....



..... [2]

[Total: 14]

- 4 20.0 g of small lumps of calcium carbonate and 40 cm<sup>3</sup> of hydrochloric acid, concentration 2.0 mol/dm<sup>3</sup>, were placed in a flask on a top pan balance. The mass of the flask and contents was recorded every minute.



The mass of carbon dioxide given off was plotted against time.



In all the experiments mentioned in this question, the calcium carbonate was in excess.

- (a) (i) Explain how you could determine the mass of carbon dioxide given off in the first five minutes.

..... [1]

- (ii) Label the graph **F** where the reaction rate is the fastest, **S** where it is slowing down and **0** where the rate is zero. [2]

- (iii) Explain how the shape of the graph shows where the rate is fastest, where it is slowing down and where the rate is zero.

.....  
 .....  
 ..... [2]

- (b) Sketch on the same graph, the line which would have been obtained if 20.0 g of small lumps of calcium carbonate and 80 cm<sup>3</sup> of hydrochloric acid, concentration 1.0 mol/dm<sup>3</sup>, had been used. [2]

(c) Explain in terms of collisions between reacting particles each of the following.


- (i) The reaction rate would be slower if 20.0 g of larger lumps of calcium carbonate and 40 cm<sup>3</sup> of hydrochloric acid, concentration 2.0 mol/dm<sup>3</sup>, were used.

.....  
 .....  
 ..... [2]

- (ii) The reaction rate would be faster if the experiment was carried out at a higher temperature.

.....  
 .....  
 ..... [2]

(d) Calculate the maximum mass of carbon dioxide given off when 20.0 g of small lumps of calcium carbonate react with 40 cm<sup>3</sup> of hydrochloric acid, concentration 2.0 mol/dm<sup>3</sup>.





number of moles of HCl used =

mass of carbon dioxide = ..... g [4]

[Total: 15]

- 5 The alkenes are unsaturated hydrocarbons. They form a homologous series, the members of which have the same chemical properties. They undergo addition reactions and are easily oxidised.

(a) The following hydrocarbons are isomers.



- (i) Explain why these two hydrocarbons are isomers.

.....  
..... [2]

- (ii) Give the structural formula of another hydrocarbon which is isomeric with the above.

[1]

- (b) Give the structural formula and name of each of the products of the following addition reactions.

- (i) ethene and bromine

structural formula of product

name of product ..... [2]

- (ii) propene and hydrogen

structural formula of product

name of product ..... [2]

- (iii) but-1-ene and water

structural formula of product

name of product ..... [2]



(c) Alkenes can be oxidised to carboxylic acids.

- (i) For example, propene,  $\text{CH}_3-\text{CH}=\text{CH}_2$ , would produce ethanoic acid,  $\text{CH}_3-\text{COOH}$ , and methanoic acid,  $\text{H}-\text{COOH}$ . Deduce the formulae of the alkenes which would form the following carboxylic acids when oxidised.

ethanoic acid and propanoic acid

only ethanoic acid


[2]

- (ii) Describe the colour change you would observe when an alkene is oxidised with acidified potassium manganate(VII).

..... [2]

(d) Alkenes polymerise to form addition polymers.

Draw the structural formula of poly(cyanoethene), include at least **two** monomer units. The structural formula of the monomer, cyanoethene, is given below.



[3]

[Total: 16]

6 Lead is an excellent roofing material. It is malleable and resistant to corrosion. Lead rapidly becomes coated with basic lead carbonate which protects it from further corrosion.

(a) Lead has a typical metallic structure which is a lattice of lead ions surrounded by a 'sea' of mobile electrons. This structure is held together by attractive forces called a metallic bond.


(i) Explain why there are attractive forces in a metallic structure.

.....  
..... [2]

(ii) Explain why a metal, such as lead, is malleable.

.....  
..... [2]

(b) Basic lead(II) carbonate is heated in the apparatus shown below. Water and carbon dioxide are produced.



(i) Silica gel absorbs water. Silica gel often contains anhydrous cobalt(II) chloride. When this absorbs water it changes from blue to pink. Suggest a reason.

..... [1]

(ii) Soda lime is a mixture of sodium hydroxide and calcium oxide. Why do these two substances react with carbon dioxide?


.....  
..... [2]

(iii) Name **two** substances formed when soda lime reacts with carbon dioxide.

..... [2]

- (c) Basic lead(II) carbonate has a formula of the type  $x\text{PbCO}_3 \cdot y\text{Pb(OH)}_2$  where x and y are whole numbers.

Determine x and y from the following information.



When heated, the basic lead(II) carbonate gave 2.112 g of carbon dioxide and 0.432 g of water.

Mass of one mole of  $\text{CO}_2 = 44 \text{ g}$

Mass of one mole of  $\text{H}_2\text{O} = 18 \text{ g}$

Number of moles of  $\text{CO}_2$  formed = ..... [1]



Number of moles of  $\text{H}_2\text{O}$  formed = ..... [1]

x = ..... and y = .....

Formula of basic lead(II) carbonate is ..... [1]

[Total: 12]

- 7 (a) The following are two examples of substitution reactions. Only the reaction involving chlorine is a photochemical reaction.



- (i) Explain the phrase *substitution reaction*.

.....  
..... [1]


- (ii) How do photochemical reactions differ from other reactions?

.....  
..... [1]

- (b) Bond forming is exothermic, bond breaking is endothermic. Explain the difference between an exothermic reaction and an endothermic reaction.

.....  
..... [2]

- (c) Use the bond energies to show that the following reaction is exothermic. Bond energy is the amount of energy (kJ/mol) which must be supplied to break one mole of the bond.



Bond energies in kJ/mol

Cl-Cl +242

C-Cl +338

C-H +412

H-Cl +431

bonds broken                      energy in kJ/mol

.....

.....

total energy = .....

bonds formed                      energy in kJ/mol

.....

.....

total energy = .....

.....

..... [4]

[Total: 8]





**DATA SHEET**  
**The Periodic Table of the Elements**

		Group																																																																																																																																																									
I	II	III	IV	V	VI	VII	0																																																																																																																																																				
		1 <b>H</b> Hydrogen 1					4 <b>He</b> Helium 2																																																																																																																																																				
7 <b>Li</b> Lithium 3	9 <b>Be</b> Beryllium 4		11 <b>B</b> Boron 5	12 <b>C</b> Carbon 6	13 <b>Al</b> Aluminium 13	14 <b>N</b> Nitrogen 7	15 <b>O</b> Oxygen 8	16 <b>F</b> Fluorine 9	17 <b>Ne</b> Neon 10	18 <b>Ar</b> Argon 18	19 <b>K</b> Potassium 19	20 <b>Ca</b> Calcium 20	21 <b>Sc</b> Scandium 21	22 <b>Ti</b> Titanium 22	23 <b>V</b> Vanadium 23	24 <b>Cr</b> Chromium 24	25 <b>Mn</b> Manganese 25	26 <b>Fe</b> Iron 26	27 <b>Co</b> Cobalt 27	28 <b>Ni</b> Nickel 28	29 <b>Cu</b> Copper 29	30 <b>Zn</b> Zinc 30	31 <b>Ga</b> Gallium 31	32 <b>Ge</b> Germanium 32	33 <b>As</b> Arsenic 33	34 <b>Se</b> Selenium 34	35 <b>Br</b> Bromine 35	36 <b>Kr</b> Krypton 36	37 <b>Rb</b> Rubidium 37	38 <b>Sr</b> Strontium 38	39 <b>Y</b> Yttrium 39	40 <b>Zr</b> Zirconium 40	41 <b>Nb</b> Niobium 41	42 <b>Mo</b> Molybdenum 42	43 <b>Tc</b> Technetium 43	44 <b>Ru</b> Ruthenium 44	45 <b>Rh</b> Rhodium 45	46 <b>Pd</b> Palladium 46	47 <b>Ag</b> Silver 47	48 <b>Cd</b> Cadmium 48	49 <b>In</b> Indium 49	50 <b>Sn</b> Tin 50	51 <b>Sb</b> Antimony 51	52 <b>Te</b> Tellurium 52	53 <b>I</b> Iodine 53	54 <b>Xe</b> Xenon 54	55 <b>Cs</b> Caesium 55	56 <b>Ba</b> Barium 56	57 <b>La</b> Lanthanum 57	58-71 <b>Lanthanoid series</b>	72 <b>Hf</b> Hafnium 72	73 <b>Ta</b> Tantalum 73	74 <b>W</b> Tungsten 74	75 <b>Re</b> Rhenium 75	76 <b>Os</b> Osmium 76	77 <b>Ir</b> Iridium 77	78 <b>Pt</b> Platinum 78	79 <b>Au</b> Gold 79	80 <b>Hg</b> Mercury 80	81 <b>Tl</b> Thallium 81	82 <b>Pb</b> Lead 82	83 <b>Bi</b> Bismuth 83	84 <b>Po</b> Polonium 84	85 <b>At</b> Astatine 85	86 <b>Rn</b> Radon 86	87 <b>Fr</b> Francium 87	88 <b>Ra</b> Radium 88	89 <b>Ac</b> Actinium 89	†90-103 <b>Actinoid series</b>	90 <b>Th</b> Thorium 90	91 <b>Pa</b> Protactinium 91	92 <b>U</b> Uranium 92	93 <b>Np</b> Neptunium 93	94 <b>Pu</b> Plutonium 94	95 <b>Am</b> Americium 95	96 <b>Cm</b> Curium 96	97 <b>Bk</b> Berkelium 97	98 <b>Cf</b> Californium 98	99 <b>Es</b> Einsteinium 99	100 <b>Fm</b> Fermium 100	101 <b>Md</b> Mendelevium 101	102 <b>No</b> Nobelium 102	103 <b>Lr</b> Lawrencium 103	104 <b>Rf</b> Rutherfordium 104	105 <b>Db</b> Dubnium 105	106 <b>Sg</b> Seaborgium 106	107 <b>Bh</b> Bohrium 107	108 <b>Hs</b> Hassium 108	109 <b>Mt</b> Meitnerium 109	110 <b>Ds</b> Darmstadtium 110	111 <b>Rg</b> Roentgenium 111	112 <b>Cn</b> Copernicium 112	113 <b>Nh</b> Nihonium 113	114 <b>Fl</b> Flerovium 114	115 <b>Lv</b> Livermorium 115	116 <b>Ts</b> Tennessine 116	117 <b>Og</b> Oganesson 117	118 <b>Uu</b> Ununseptium 118	119 <b>Uub</b> Ununseptium 119	120 <b>Uuq</b> Ununseptium 120	121 <b>Uub</b> Ununseptium 121	122 <b>Uuq</b> Ununseptium 122	123 <b>Uub</b> Ununseptium 123	124 <b>Uuq</b> Ununseptium 124	125 <b>Uub</b> Ununseptium 125	126 <b>Uuq</b> Ununseptium 126	127 <b>Uub</b> Ununseptium 127	128 <b>Uuq</b> Ununseptium 128	129 <b>Uub</b> Ununseptium 129	130 <b>Uuq</b> Ununseptium 130	131 <b>Uub</b> Ununseptium 131	132 <b>Uuq</b> Ununseptium 132	133 <b>Uub</b> Ununseptium 133	134 <b>Uuq</b> Ununseptium 134	135 <b>Uub</b> Ununseptium 135	136 <b>Uuq</b> Ununseptium 136	137 <b>Uub</b> Ununseptium 137	138 <b>Uuq</b> Ununseptium 138	139 <b>Uub</b> Ununseptium 139	140 <b>Uuq</b> Ununseptium 140	141 <b>Uub</b> Ununseptium 141	142 <b>Uuq</b> Ununseptium 142	143 <b>Uub</b> Ununseptium 143	144 <b>Uuq</b> Ununseptium 144	145 <b>Uub</b> Ununseptium 145	146 <b>Uuq</b> Ununseptium 146	147 <b>Uub</b> Ununseptium 147	148 <b>Uuq</b> Ununseptium 148	149 <b>Uub</b> Ununseptium 149	150 <b>Uuq</b> Ununseptium 150	151 <b>Uub</b> Ununseptium 151	152 <b>Uuq</b> Ununseptium 152	153 <b>Uub</b> Ununseptium 153	154 <b>Uuq</b> Ununseptium 154	155 <b>Uub</b> Ununseptium 155	156 <b>Uuq</b> Ununseptium 156	157 <b>Uub</b> Ununseptium 157	158 <b>Uuq</b> Ununseptium 158	159 <b>Uub</b> Ununseptium 159	160 <b>Uuq</b> Ununseptium 160	161 <b>Uub</b> Ununseptium 161	162 <b>Uuq</b> Ununseptium 162	163 <b>Uub</b> Ununseptium 163	164 <b>Uuq</b> Ununseptium 164	165 <b>Uub</b> Ununseptium 165	166 <b>Uuq</b> Ununseptium 166	167 <b>Uub</b> Ununseptium 167	168 <b>Uuq</b> Ununseptium 168	169 <b>Uub</b> Ununseptium 169	170 <b>Uuq</b> Ununseptium 170	171 <b>Uub</b> Ununseptium 171	172 <b>Uuq</b> Ununseptium 172	173 <b>Uub</b> Ununseptium 173	174 <b>Uuq</b> Ununseptium 174	175 <b>Uub</b> Ununseptium 175

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).

	<b>a</b>	a = relative atomic mass
<b>Key</b>	<b>X</b>	X = atomic symbol
	<b>b</b>	b = proton (atomic) number

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.