

Cambridge IGCSE™

MATHEMATICS**0580/21**

Paper 2 (Extended)

May/June 2025

MARK SCHEME

Maximum Mark: 100

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2025 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

This document consists of **10** printed pages.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics-Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- 5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

Annotations guidance for centres

Examiners use a system of annotations as a shorthand for communicating their marking decisions to one another. Examiners are trained during the standardisation process on how and when to use annotations. The purpose of annotations is to inform the standardisation and monitoring processes and guide the supervising examiners when they are checking the work of examiners within their team. The meaning of annotations and how they are used is specific to each component and is understood by all examiners who mark the component.

We publish annotations in our mark schemes to help centres understand the annotations they may see on copies of scripts. Note that there may not be a direct correlation between the number of annotations on a script and the mark awarded. Similarly, the use of an annotation may not be an indication of the quality of the response.

The annotations listed below were available to examiners marking this component in this series.

Annotations

| Annotation | Meaning |
|-------------|--------------------------------------|
| A | More information required |
| A0 | Accuracy mark awarded zero |
| A1 | Accuracy mark awarded one |
| A2 | Accuracy mark awarded two |
| A3 | Accuracy mark awarded three |
| B0 | Independent mark awarded zero |
| B1 | Independent mark awarded one |
| B2 | Independent mark awarded two |
| B3 | Independent mark awarded three |
| BOD | Benefit of the doubt |
| C | Communication mark |
| X | Incorrect |
| FT | Follow through |
| Highlighter | Highlight a key point in the working |
| ISW | Ignore subsequent work |
| M0 | Method mark awarded zero |
| M1 | Method mark awarded one |
| M2 | Method mark awarded two |
| M3 | Method mark awarded three |

| Annotation | Meaning |
|---|--|
| MR | Misread |
| O | Omission |
| Off-page comment | Allows comments to be entered at the bottom of the RM marking window and then displayed when the associated question item is navigated to. |
| On-page comment | Allows comments to be entered in speech bubbles on the candidate response. |
| Pre | Premature rounding/approximation |
| SC | Special case |
| SEEN | Indicates that work/page has been seen |
| TE | Transcription error |
|  | Correct |
| XP | Correct answer from incorrect working |

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M Method marks, awarded for a valid method applied to the problem.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation ‘dep’ is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

| | |
|------|----------------------------|
| awrt | answers which round to |
| cao | correct answer only |
| dep | dependent |
| FT | follow through after error |
| isw | ignore subsequent working |
| nfww | not from wrong working |
| oe | or equivalent |
| rot | rounded or truncated |
| SC | Special Case |
| soi | seen or implied |

| Question | Answer | Marks | Partial Marks |
|----------|--|-------|---|
| 1 | $8c - 2d$ final answer | 2 | B1 for answer $8c - kd$ or $kc - 2d$ or for correct answer seen and spoilt |
| 2 | $[w =] 158$ $[x =] 76$ $[y =] 82$ | 4 | B1 for w correct B1 for x correct B2FT for $y = 158 - \text{their } x$ correctly evaluated or B1 for 22 (identified) or 82 in position vertically opposite to y or for $y = 158 - \text{their } x$ |
| 3 | 1770 | 3 | B2 for 270 or M2 for $1500 + \frac{1500 \times 3 \times 6}{100}$ or M1 for $\frac{1500 \times 3 [\times 6]}{100}$ |
| 4 | $\frac{7}{12}$ oe | 3 | M1 for $\frac{6}{24}$ oe M1 for correct use of common denominator in subtraction $\frac{5}{6} - \text{their } \frac{6}{24}$, e.g. $\frac{20}{24}$ and $\frac{6}{24}$ oe If 0 scored, SC1 for answer $\frac{1}{16}$ oe |
| 5 | 12 | 2 | M1 for $\frac{360}{180-150}$ or for $\frac{180(n-2)}{n} = 150$ |
| 6(a) | Correct line drawn | 2 | M1 for a line with gradient 2 or for a line with positive gradient and intercept at $y = 1$ |
| 6(b) | $[x =] 2$, $[y =] 5$ | 1 | FT intersection of <i>their (a)</i> with the given line |
| 7 | $\frac{4}{15}$ | 3 | B2 for $\frac{24}{90}$ oe or M1 for 26.66... – 2.66... oe or for $90x = 24$ oe or for $\frac{2}{10} + \frac{6}{90}$ oe |
| 8(a) | $\begin{pmatrix} 14 \\ 13 \end{pmatrix}$ | 2 | B1 for $\begin{pmatrix} 14 \\ k \end{pmatrix}$ or for $\begin{pmatrix} k \\ 13 \end{pmatrix}$ or for $\begin{pmatrix} 22 \\ 10 \end{pmatrix}$ |
| 8(b) | 24 | 2 | M1 for $y + 5^2 = 7^2$ or better |
| 9 | 4 nfww | 3 | M2 for $8 + 20 + 8n = 6(2 + 4 + n)$ or better or M1 for $\frac{2 \times 4 + 4 \times 5 + n \times 8}{2 + 4 + n} [= 6]$ oe or $8 + 20 + 8n$ or $6(2 + 4 + n)$ |

| Question | Answer | Marks | Partial Marks |
|----------|---|-------|--|
| 10(a) | 110 | 1 | |
| 10(b) | 55 | 1 | FT <i>their</i> $110 \div 2$ |
| 10(c) | 55 | 1 | FT <i>their</i> (b) Provided <i>their</i> (b) < 90 |
| 10(d) | 15 | 1 | FT $70 - \text{their (b)}$ Provided <i>their</i> (b) < 70 |
| 11(a) | For correct ruled tangent and $[a =] 1, [b =] 2$ | 2 | B1 for correct ruled tangent or both values correct without a correct tangent |
| 11(b) | $[y =] 5 - 3x$ | 3 | 3FT <i>their</i> (a) provided $m < 0, c \neq 0$ B1 for $(\text{their } -3)x + c$ or M1 for correct $\frac{\text{rise}}{\text{run}}$ for <i>their</i> line B1 for $mx + c$ where c is the correct intercept for <i>their</i> graph, $m \neq 0$ |
| 12(a) | 1.4 | 2 | B1 for $[\text{UQ}] = 5.6$ or for $[\text{LQ}] = 4.2$ or SC1 for 0.7 |
| 12(b) | 5.4 | 2 | B1 for 84 seen |
| 13(a) | 12.5 oe | 5 | M4 for $8[\pi]h = 100[\pi]$ OR M3 for $\pi \times 6 \times 10 + 2 \times \pi \times 6^2 = 2 \times \pi \times 4^2 + 2 \times \pi \times 4 \times h$ oe OR M1 for $\pi \times 6 \times 10 + 2 \times \pi \times 6^2$ oe M1 $2 \times \pi \times 4^2 + 2 \times \pi \times 4 \times h$ oe OR SC2 for answer 16.5 |
| 13(b) | 14 | 3 | M2 for $\sqrt{10^2 - 6^2}$ or M1 for $6^2 + x^2 = 10^2$ |
| 14(a) | -10 | 1 | |

| Question | Answer | Marks | Partial Marks |
|-----------|---|-------|--|
| 14(b) | $\frac{x+4}{3}$ oe | 2 | M1 for correct first step $y + 4 = 3x$ or $x = 3y - 4$ or $\frac{y}{3} = x - \frac{4}{3}$ |
| 14(c) | $a = 12, b = -1$ | 2 | B1 for either a or b correct or M1 for $3(4x + 1) - 4$ |
| 14(d) | $\frac{22 - 7x}{(3x - 4)(4x + 1)}$ or $\frac{22 - 7x}{12x^2 - 13x - 4}$ final answer | 3 | B1 for $2(4x + 1) - 5(3x - 4)$ oe or better isw B1 for common denominator $(3x - 4)(4x + 1)$ oe isw |
| 15(a) | $17 - 7\sqrt{5}$ final answer | 2 | B1 for 3 correct terms from $2 - 6\sqrt{5} - \sqrt{5} + 3 \times \sqrt{5} \times \sqrt{5}$ oe |
| 15(b) | $\frac{3\sqrt{10}}{5}$ cao | 2 | M1 for $\frac{6}{\sqrt{10}} \times \frac{\sqrt{10}}{\sqrt{10}}$ oe |
| 16 | $3x^3 + 5x^2 - 34x - 24$ final answer | 3 | B2 for correct expansion unsimplified or simplified four-term expression of correct form with 3 terms correct or B1 for one pair of brackets expanded with at least 3 terms out of 4 correct |
| 17(a) | $\frac{9}{100}$ oe | 2 | M1 for $\frac{3}{10} \times \frac{3}{10}$ |
| 17(b)(i) | $\frac{2}{5}$ $\frac{4}{5}$ $\frac{1}{5}$ | 2 | B1 for $\frac{2}{5}$ |
| 17(b)(ii) | $\frac{16}{30}$ oe | 3 | 3FT their tree diagram dep on probabilities < 1 M2FT for $\frac{4}{6} \times \frac{2}{5} + \frac{2}{6} \times \frac{4}{5}$ or M1FT for $\frac{4}{6} \times \frac{2}{5}$ or for $\frac{2}{6} \times \frac{4}{5}$ |
| 18(a) | $\frac{12}{x}$ | 1 | |
| 18(b) | $\frac{10}{x-4}$ | 1 | |

| Question | Answer | Marks | Partial Marks |
|-----------|--|-------|---|
| 18(c) | $their \frac{10}{x-4} - their \frac{12}{x} = 1 \text{ oe}$ | M1 | |
| | $10x - 12x + 48 = x^2 - 4x$ | M2 | Correctly multiplying <i>their</i> brackets and clearing algebraic fractions e.g. $(x-4)\left(\frac{12}{x} + 1\right) = 10$ leading to $12 - \frac{48}{x} + x - 4 = 10$ and then $12x - 48 + x^2 - 4x = 10x$ or M1 for correctly clearing, or correctly collecting into a single fraction, two fractions both with different algebraic denominators e.g. $10x - 12(x-4) = x(x-4)$ or $\frac{10x - 12(x-4)}{x(x-4)} [= 1]$ |
| | Leading to $0 = x^2 - 2x - 48$ | A1 | With no errors or omissions seen, dep on M3 |
| 18(d) | $(x+6)(x-8)$ | M2 | M1 for $x(x-8) + 6(x-8)$ or $x(x+6) - 8(x+6)$ or $(x+a)(x+b)$ where $ab = -48$ or $a + b = -2$ |
| | -6, 8 | B1 | |
| 18(e) | 1.5 or $1\frac{1}{2}$ | 1 | FT $\frac{12}{their 8}$ |
| 19 | $\frac{1}{9}$ | 2 | M1 for 9^{-1} or for $\frac{1}{3^2}$ or $\frac{1}{\sqrt[3]{729}}$ |
| 20 | $6\sqrt{3} \text{ oe}$ | 4 | B1 for $\tan 30 = \frac{1}{\sqrt{3}}$ or $\frac{\sqrt{3}}{3}$ or for $\sin 60 = \frac{\sqrt{3}}{2}$ and $\sin 30 = \frac{1}{2}$ M2 for $\frac{6}{\tan 30}$ or $\frac{6\sin(60)}{\sin(30)}$ or M1 for $\frac{6}{x} = \tan 30$ or $\frac{\sin 60}{x} = \frac{\sin 30}{6}$ |
| 21(a)(i) | $2\mathbf{m} - \mathbf{n}$ | 1 | |
| 21(a)(ii) | $2\mathbf{m} + \mathbf{n}$ | 1 | |

| Question | Answer | Marks | Partial Marks |
|----------|--|-------|--|
| 21(b) | $\overrightarrow{CD} = 10\mathbf{m} - 5\mathbf{n}$ | M2 | Allow M2 for equivalents $\overrightarrow{CD} = -2\mathbf{n} + 10\mathbf{m} - 3\mathbf{n}$ or $\overrightarrow{CD} = -(2\mathbf{m} + \mathbf{n}) + 2\mathbf{m} - \mathbf{n} + 10\mathbf{m} - 3\mathbf{n}$ For M2, FT <i>their (a)</i> e.g. $\overrightarrow{CD} = \text{their } \overrightarrow{CO} + \text{their } \overrightarrow{OA} + 10\mathbf{m} - 3\mathbf{n}$ or M1 for correct route for \overrightarrow{CD} using the lines of the diagram with \overrightarrow{AD} e.g. $\overrightarrow{CD} = \overrightarrow{CA} + \overrightarrow{AD}$ oe |
| | $\overrightarrow{CD} = 5\overrightarrow{OA}$ leading to CD is parallel to OA [$\therefore OACD$ is a trapezium] | A1 | Dependent on M2 |
| 22(a) | $[n =] 3, [q =] -6$ | 2 | B1 for each correct value |
| 22(b) | (1, 4) and (3, 0) | 4 | B3 for (1, 4) or (3, 0) or for two correct values of x or M2 for $[3](x-1)(x-3) [= 0]$ oe $\text{or } x = \frac{-(-12) \pm \sqrt{(-12)^2 - 4 \times 3 \times 9}}{2 \times 3} \text{ oe}$ or M1 for writing $\frac{dy}{dx} = 0$ or for $3x^2 - 12x + 9 = 0$ |
| 23 | $\frac{2x}{x-5}$ final answer | 3 | B1 for $2x(x+5)$ B1 for $(x-5)(x+5)$ |