



Cambridge IGCSE™

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

ADDITIONAL MATHEMATICS

0606/22

Paper 2

May/June 2026

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- You should use a scientific calculator where appropriate.
- You must show all necessary working clearly.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- For π , use either your calculator value or 3.142.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages.

List of formulas

Equation of a circle with centre (a, b) and radius r .

$$(x - a)^2 + (y - b)^2 = r^2$$

Curved surface area, A , of cone of radius r , sloping edge l .

$$A = \pi r l$$

Surface area, A , of sphere of radius r .

$$A = 4\pi r^2$$

Volume, V , of pyramid or cone, base area A , height h .

$$V = \frac{1}{3} Ah$$

Volume, V , of sphere of radius r .

$$V = \frac{4}{3} \pi r^3$$

Quadratic equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series

$$u_n = a + (n-1)d$$

$$S_n = \frac{1}{2} n(a + l) = \frac{1}{2} n\{2a + (n-1)d\}$$

Geometric series

$$u_n = ar^{n-1}$$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

Identities

$$\sin^2 A + \cos^2 A = 1$$

$$\sec^2 A = 1 + \tan^2 A$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

Formulas for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2} ab \sin C$$



1 The point A is the minimum point on the curve $y = x^2 - 6x + 10$.

(a) Using the method of completing the square, find the coordinates of A .

[2]

(b) The point B has coordinates $(6, 7)$.
The curve cuts the y -axis at the point C .

Find the equation of the line through B parallel to the line AC .

[2]

- 2 Show that there is no real value of k for which the equation
- $$(k-2)x^2 + (2k+3)x + 3 = 0$$
- has two equal roots.

[3]

- 3 (a) Differentiate $y = \sqrt{x+1} + \cos(5x+3)$ with respect to x .

[3]


- (b) Find $\int \sec^2 \frac{x}{3} dx$.

[3]



- 4 (a) On the axes, sketch the graph of $y = |x^2 - 5|$.
State the intercepts with the coordinate axes.

[3]



- (b) It is given that $f(x) = ax + b$ where a and b are non-zero integers.
The solutions of the equation $|f(x)| = 4$ are $x = 2.5$ and $x = -1.5$.

Find the possible expressions for $f(x)$.

[4]

- 5 (a) In this question, 6-digit numbers do not start with 0.

Find how many 6-digit numbers have 6 different digits and are divisible by 2.

[3]

- (b) A committee of 10 people is to be selected from 9 students and 7 teachers.

Find the number of different committees that can be selected if the committee must have at least 4 students and at least 4 teachers.

[3]



- 6 In this question, a , b and c are integers.

When the expansion of $(a+x)^4(1+ax)^5$ is written in ascending powers of x , the first three terms are $16+bx+cx^2$.

Find all the possible values of a , b and c .

[8]



- 7 In this question, $\mathbf{i}$ is a unit vector due east and $\mathbf{j}$ is a unit vector due north. Distances are measured in kilometres and time is measured in hours.

Point P has position vector $2\mathbf{i} + 5\mathbf{j}$ relative to an origin O .

At 13 00, boat A sails from point P with velocity vector $8\mathbf{i} + 8\mathbf{j}$.

- (a) Find the position vector of A at the end of 2 hours of sailing. [1]

- (b) Write down the position vector of A at the end of t hours of sailing. [1]

- (c) Point Q has position vector $20\mathbf{i} + 8\mathbf{j}$ relative to the origin O .
At 15 00, boat B sails from point Q on a bearing of 300° with a constant speed of $5\sqrt{3}$.

Find the position vector of B at 17 00. [3]



(d) Find the distance between the two boats at 17 00.

[3]



8 The function f is defined, for $x \geq -1$, by $f(x) = 1 + 2^x$.

(a) Find the range of f .

[1]

(b) (i) Write down the domain of f^{-1} .


[1]

(ii) Find an expression for $f^{-1}(x)$.

[2]

(iii) On the axes, sketch the graph of $y = f(x)$ and hence the graph of $y = f^{-1}(x)$.
State any intercepts with the coordinate axes.

[4]



(c) The function g is defined, for $x \geq -1$, by $g(x) = x(x+k)$ where k is a constant.

(i) Explain why g exists.

[1]

(ii) Find an expression for $g'(x)$.

[1]

(iii) It is given that $g(3) = 126$.

Find the value of k .

[2]



- 9 A particle P moves in a straight line such that t seconds after passing through a fixed point O , its displacement from O , s metres, is given by $s = e^{4t} - 10e^{2t} + 9$.

(a) The velocity of P is $v \text{ ms}^{-1}$.

Find the value of t for which v has a stationary value.

[5]

(b) Find the acceleration of P when $t = \ln 2$.

[2]



10 Solve the equation $\tan(3x+1) = 5 \sin(3x+1)$ for $-\frac{\pi}{3} < x < \frac{\pi}{3}$.

[6]



11 Solutions to this question by accurate drawing will not be accepted.

A circle C_1 has centre $(3, 5)$.

The point $P(1, 4)$ lies on circle C_1 .

A circle C_2 has centre $(12, 8)$.

The line with equation $y = 19 - 3x$ is the common chord of circles C_1 and C_2 .

Find the radius of circle C_2 .

[9]





Continuation of working space for Question 11.

Question 12 is on the back page.



- 12 A sphere has volume $V \text{ cm}^3$ and radius $r \text{ cm}$.

Given that r varies with time t , find V at the instant when $\frac{dV}{dt} = 9\frac{dr}{dt}$. [4]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.

