



Cambridge IGCSE™

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



ADDITIONAL MATHEMATICS

0606/23

Paper 2

October/November 2023

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY

Identities

$$\begin{aligned} \sin^2 A + \cos^2 A &= 1 \\ \sec^2 A &= 1 + \tan^2 A \\ \operatorname{cosec}^2 A &= 1 + \cot^2 A \end{aligned}$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}bc \sin A$$

- 1 The functions f and g are defined as follows, for all real values of x .

$$f(x) = 2 \sin x + 3 \cos x$$

$$g(x) = e^{3x} - 1$$

- (a) Find $fg(0)$.

[2]

- (b) Find $gg(x)$.

[1]

- (c) Solve the equation $g^{-1}(x) = \frac{1}{3} \ln 5$.

[3]

- 2 Find the values of k for which the curve $y = x^2 + kx + (4k - 15)$ is completely above the x -axis. [4]

3 (a) Solve the following simultaneous equations.

$$3 \log_2 x + 2 \log_2 y = 24$$

$$5 \log_2 x - 3 \log_2 y = 2$$

[5]

(b) Solve the equation $\frac{2^{t+4}}{2^{1-2t}} = 512$. [4]

- 4 Find the exact value of $\int_3^5 \frac{(x-1)^2}{x^3} dx$. [6]

- 5 The curved surface area of a cylinder with radius r and height h is $2\pi rh$.

A closed cylinder has radius r cm and volume 1000 cm 3 .

- (a) Show that the total surface area of the cylinder is $2\pi r^2 + \frac{2000}{r}$ cm 2 . [3]

- (b) Find the value of r which makes this area a minimum. You should show that your value of r gives a minimum for this area. [5]


- 6 A particle travels in a straight line. Its displacement, s metres, from the origin, at time t seconds, where $t > 2$, is given by $s = \ln(4t^2 - 5) - t$.

(a) Find expressions for the velocity, $v \text{ ms}^{-1}$, and acceleration, $a \text{ ms}^{-2}$, of the particle. [4]

(b) Find the time when the particle is at rest. [3]

(c) Find the acceleration at this time. [2]

7 DO NOT USE A CALCULATOR IN THIS QUESTION.



You may use the following trigonometrical ratios.

$$\sin 60^\circ = \frac{\sqrt{3}}{2}, \sin 45^\circ = \frac{\sqrt{2}}{2}$$

$$\cos 60^\circ = \frac{1}{2}, \cos 45^\circ = \frac{\sqrt{2}}{2}$$

$$\tan 60^\circ = \sqrt{3}, \tan 45^\circ = 1$$

- (a) Given that the area of triangle ABC is $\frac{3 + \sqrt{3}}{4}$, show that $\sin 75^\circ = \frac{\sqrt{6} + \sqrt{2}}{4}$. [5]

- (b) Hence find the exact length of AC . [2]

- 8 (a) Show that $\frac{\sin x}{\tan x - 1} - \frac{\cos x}{\tan x + 1} = \frac{\cos x}{\sin^2 x - \cos^2 x}$. [5]

- (b) Hence solve the equation $\frac{\sin x}{\tan x - 1} - \frac{\cos x}{\tan x + 1} = 1$ for $0^\circ < x < 360^\circ$. [5]

9 A curve has equation $y = xe^{2x}$.

(a) Find $\frac{dy}{dx}$.

[2]

(b) Find the equation of the normal to the curve at $x = 1$.

[4]

- (c) Use your answer to **part (a)** to find the exact value of $\int_0^2 2xe^{2x} dx$. [5]

- 10 (a) In an arithmetic progression the 5th term is 11. The 7th term is three times the 2nd term. Find the 1st term and the common difference.

[4]

(b) A different arithmetic progression (AP) and a geometric progression (GP) have the following properties.

- The 1st terms of the AP and GP are both 3.
- The 2nd term of the AP is the same as the 3rd term of the GP.
- The 6th term of the AP is the same as the 5th term of the GP.
- The common ratio of the GP is greater than 1.

Find the common difference of the AP and the common ratio of the GP.

[6]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.