
ADDITIONAL MATHEMATICS**0606/11**

Paper 1

October/November 2018

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2018 series for most Cambridge IGCSE™, Cambridge International A and AS Level components and some Cambridge O Level components.

This document consists of **9** printed pages.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.


Types of mark


- M Method marks, awarded for a valid method applied to the problem.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more ‘method’ steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation ‘**dep**’ is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

awrt	answers which round to
cao	correct answer only
dep	dependent
FT	follow through after error
isw	ignore subsequent working
nfww	not from wrong working
oe	or equivalent
rot	rounded or truncated
SC	Special Case
soi	seen or implied

Question	Answer	Marks	Guidance
1(a)		B3	B1 for 2 cycles, one max and 2 min points in the correct places and up to a max at each end B1 for going between 2 and -4 B1 for starting at (0,2) and finishing at (360,2)
1(b)(i)	4	B1	
1(b)(ii)	60° or $\frac{\pi}{3}$	B1	
2(i)	$\left(p\left(-\frac{1}{2}\right) = \right) -\frac{1}{4} + \frac{5}{4} - 2 + a = 2$ $(q(-2) =) 16 - 6a + b = 0$	M1	For either $p\left(-\frac{1}{2}\right) = 2$ or $q(-2) = 0$
	$a = 3$	A1	
	$b = 2$	A1	
2(ii)	$r(x) = 2x^3 + x^2 - 5x + 1$	M1	For $r(x)$ using <i>their</i> $p(x)$ and $q(x)$
	$r\left(\frac{2}{3}\right) = \frac{16}{27} + \frac{4}{9} - \frac{10}{3} + 1$	M1	For $r\left(\frac{2}{3}\right)$
	$= -\frac{35}{27}$	A1	Must be exact

Question	Answer	Marks	Guidance
3	$(3+ kx)^6 =$ $729 + 1458kx + 1215k^2x^2$	B2	B1 for $1458kx$ or $1215k^2x^2$
	Terms in x^2 for $(2-x)(3+kx)^6$ $= -1458k + 2430k^2$ $2430k^2 - 1458k = 972$	M1	For attempt at further expansion to obtain 2 terms in x^2 and equating to 972
	$5k^2 - 3k - 2 = 0$ $(5k+2)(k-1) = 0$	M1	Dep for solution of resulting 3 term quadratic
	$k = -\frac{2}{5}$	A1	
	$k = 1$	A1	
4(i)	$\left(x - \frac{9}{2}\right)^2 - \frac{49}{4}$	B2	B1 for $\frac{9}{2}$ or $\frac{49}{4}$
4(ii)	$\left(\frac{9}{2}, -\frac{49}{4}\right)$	B1	FT <i>their p</i> and <i>q</i>
4(iii)		B3	B1 for shape B1 for cusps at $(1, 0)$ and $(8, 0)$ B1 for all correct, passing through $(0, 8)$ with maximum in correct position
4(iv)	$\frac{49}{4}$	B1	FT <i>their q</i>
5(i)	$\frac{1}{2}r^2\theta = 48$	B1	
	$\theta = \frac{96}{r^2}$	B1	Dep Must have previous B1
5(ii)	$r\theta = 12$	B1	For statement of arc length
	$r \times \frac{96}{r^2} = 12$	M1	For attempt to use part (i) to find either r or θ
	$r = 8$ and $\theta = 1.5$	A1	For both

Question	Answer	Marks	Guidance
5(iii)	$\text{Area} = 48 - \left(\frac{1}{2} r^2 \sin \theta \right)$	M1	For a complete method including a correct attempt for the area of the triangle using their r and θ
	$=16.1$	A1	
6(i)	For $\frac{4x}{2x^2 + 3}$	B1	
		M1	For attempt to differentiate a quotient or appropriate product
	$\frac{dy}{dx} = \frac{(5x+2) \frac{4x}{2x^2+3} - 5 \ln(2x^2+3)}{(5x+2)^2}$	A1	All other terms correct
	When $x=0$ $\frac{dy}{dx} = \frac{-5 \ln 3}{4}$	A1	For given answer
6(ii)	$y = \frac{1}{2} \ln 3$ or 0.549	B1	May be implied by tangent equation, allow 0.55
	Equation of tangent $y = \left(-\frac{5}{4} \ln 3 \right) x + \frac{1}{2} \ln 3$ or $y = -1.37x + 0.549$	B1	
7(a)	$\lg 100 = 2$	B1	
	$3 \lg x = \lg x^3$	B1	
	$\lg \frac{100x^3}{y}$	B1	
7(b)(i)	$6x^2 + 7x - 3 = 0$ $(2x+3)(3x-1) = 0$	M1	For obtaining in suitable quadratic form and attempt to solve
	$x = -\frac{3}{2}$ $x = \frac{1}{3}$	A1	For both

Question	Answer	Marks	Guidance
7(b)(ii)	$x = \log_a 3$ $\frac{1}{3} = \log_a 3$	M1	For realising connection with part (i) and attempt to solve $\frac{1}{3} = \log_a 3$ or $-\frac{3}{2} = \log_a 3$
	$a = 27$	A1	
	$-\frac{3}{2} = \log_a 3$	M1	For realising connection with part (i) and attempt to solve $-\frac{3}{2} = \log_a 3$ or $\frac{1}{3} = \log_a 3$
	$a = \left(\frac{1}{3}\right)^{\frac{2}{3}}$ or 0.481 or $\left(\frac{1}{9}\right)^{\frac{1}{3}}$ oe	A1	
8(i)		M1	For attempt to use chain rule to obtain $kx(5x^2 + 4)^{\frac{1}{2}}$ where k is a constant
	$\frac{3}{2}(10x)(5x^2 + 4)^{\frac{1}{2}}$	A1	Allow unsimplified
8(ii)		M1	For attempt to use part (i) if in correct form of $m(5x^2 + 4)^{\frac{3}{2}}$
	$\frac{1}{15}(5x^2 + 4)^{\frac{3}{2}} (+c)$	A1	FT on <i>their</i> $\frac{1}{k}(5x^2 + 4)^{\frac{3}{2}}$
8(iii)		M1	For use of limits if <i>their</i> (ii) Must be in the form $m(5x^2 + 4)^{\frac{3}{2}}$
	$\frac{1}{15}\left((5a^2 + 4)^{\frac{3}{2}} - 4^{\frac{3}{2}}\right) \left[= \frac{19}{15} \right]$	A1	
	$(5a^2 + 4)^{\frac{3}{2}} = 27$	M1	Dep For complete and correct method to deal with the power of $\frac{3}{2}$
	leading to $a = 1$	A1	
9(i)	3	B1	

Question	Answer	Marks	Guidance
9(ii)		M1	For attempt to differentiate to obtain $a + be^{-t}$
	$\frac{ds}{dt} = 4 - 3e^{-t}$	A1	All correct
	$2 = 4 - 3e^{-t}$	M1	Dep for correct attempt to solve equation involving exponential where $e^{-t} > 0$
	leading to $t = \ln \frac{3}{2}$ or $-\ln \frac{2}{3}$	A1	Must be an exact form
9(iii)	When $t = \ln 5$, $\frac{ds}{dt} = \frac{17}{5}$	M1	For attempt to find value of $\frac{ds}{dt}$ when $t = \ln 2$
		M1	Dep for attempt to use method of small changes
	$\partial s = \frac{17h}{5}$	A1	
10(i)	Velocity of A $\begin{pmatrix} 6 \\ 8 \end{pmatrix}$	B1	For velocity, may be implied by later work
	When $t = 6$, $\mathbf{r}_A = \begin{pmatrix} 2 \\ -5 \end{pmatrix} + 6 \begin{pmatrix} 6 \\ 8 \end{pmatrix}$	M1	For a complete and correct method
	$= \begin{pmatrix} 38 \\ 43 \end{pmatrix}$	A1	For 43
10(ii)	$\mathbf{r}_B = \begin{pmatrix} 16 \\ 37 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} t$	B1	
10(iii)	$\begin{pmatrix} 16 \\ 37 \end{pmatrix} + \begin{pmatrix} 4 \\ 2 \end{pmatrix} t = \begin{pmatrix} 2 \\ -5 \end{pmatrix} + \begin{pmatrix} 6 \\ 8 \end{pmatrix} t$	M1	For equating position vectors at a time t
	$16 + 4t = 2 + 6t$ or $37 + 2t = -5 + 8t$	M1	Dep for equating like vectors at least once
	$t = 7$	A1	Allow from one correct equation
	Both equations lead to $t = 7$	B1	For showing that $t = 7$ satisfies both equations thus verifying collision, or equivalent
10(iv)	$\begin{pmatrix} 44 \\ 51 \end{pmatrix}$	B1	

Question	Answer	Marks	Guidance
11(a)(i)		B1	For critical values
	$2 \leq f \leq 4$	B1	Dep For correct inequality and notation
11(a)(ii)	$x = 3 \cos y$ $\cos 2y = 0.5$	M1	For attempt to find f^{-1} and equate to 0.5
	$2y = \frac{\pi}{3}$	M1	Dep For correct attempt to solve, dealing with the double angle
	$y = \frac{\pi}{6}$	A1	
11(b)	$g^2(x) = g(3 - x^2)$ $= 3 - (3 - x^2)^2$	M1	For correct attempt at g^2 , allow unsimplified
	$-6 + 6x^2 - x^4 = -6$ $6x^2 - x^4 = 0$	M1	Dep for equating to -6 and attempt to solve to obtain a non-zero root
	$x = 0$	B1	
	$x = \pm\sqrt{6}$	A1	