## MARK SCHEME for the May/June 2013 series

# 0625 PHYSICS

0625/31

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.



Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0625	31

### NOTES ABOUT MARK SCHEME SYMBOLS & OTHER MATTERS

- M marks are method marks upon which further marks depend. For an M mark to be scored, the point to which it refers **must** be seen in a candidate's answer. If a candidate fails to score a particular M mark, then none of the dependent marks can be scored.
- B marks are independent marks, which do not depend on other marks. For a B mark to be scored, the point to which it refers must be seen specifically in the candidate's answers.
- A marks In general A marks are awarded for final answers to numerical questions.
  If a final numerical answer, eligible for A marks, is correct, with the correct unit and an acceptable number of significant figures, all the marks for that question are normally awarded.
  It is very occasionally possible to arrive at a correct answer by an entirely wrong approach. In these rare circumstances, do not award the A marks, but award C marks on their merits. However, correct numerical answers with no working shown gain all the marks available.
- C marks are compensatory marks in general applicable to numerical questions. These can be scored even if the point to which they refer are not written down by the candidate, **provided subsequent working gives evidence that they must have known it.** For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct substitution or working which shows he knew the equation, then the C mark is scored. A C mark is not awarded if a candidate makes two points which contradict each other. Points which are wrong but irrelevant are ignored.
- brackets () around words or units in the mark scheme are intended to indicate wording used to clarify the mark scheme, but the marks do not depend on seeing the words or units in brackets, e.g. 10 (J) means that the mark is scored for 10, regardless of the unit given.
- <u>underlining</u> indicates that this <u>must</u> be seen in the answer offered, or something very similar.
- OR / or indicates alternative answers, any one of which is satisfactory for scoring the marks.
- e.e.o.o. means 'each error or omission'.
- o.w.t.t.e. means 'or words to that effect'.
- Spelling Be generous about spelling and use of English. If an answer can be understood to mean what we want, give credit. However, beware of and do not allow ambiguities, accidental or deliberate: e.g. spelling which suggests confusion between reflection / refraction / diffraction / thermistor / transistor / transformer.
- Not/NOT Indicates that an incorrect answer is not to be disregarded, but cancels another otherwise correct alternative offered by the candidate i.e. right plus wrong penalty applies.
- Ignore Indicates that something which is not correct or irrelevant is to be disregarded and does not cause a right plus wrong penalty.

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – May/June 2013	0625	31

e.c.f. meaning 'error carried forward' is mainly applicable to numerical questions, but may in particular circumstances be applied in non-numerical questions. This indicates that if a candidate has made an earlier mistake and has carried an incorrect value forward to subsequent stages of working, marks indicated by ecf may be awarded, provided the subsequent working is correct, bearing in mind the earlier mistake. This prevents a candidate being penalised more than once for a particular mistake, but **only** applies to marks annotated e.c.f.

#### Significant Figures

Answers are normally acceptable to any number of significant figures  $\dot{u}$  2. Accept answers that round to give the correct answer to 2 s.f. Any exceptions to this general rule will be specified in the mark scheme.

Units Deduct one mark for each incorrect or missing unit from a final answer that would otherwise gain all the marks available for that answer: maximum 1 per question.

#### Arithmetic errors

Deduct one mark if the **only** error in arriving at a final answer is clearly an arithmetic one.

#### Transcription errors

Deduct one mark if the only error in arriving at a final answer is because given or previously calculated data has clearly been misread but used correctly.

Fractions e.g. <sup>1</sup>/<sub>2</sub>, <sup>1</sup>/<sub>4</sub>, 1/10 etc. are only acceptable where specified.

	Page 4					Syllabus	Paper	,			
					IGCS	E – May/	June 2013		0625	31	
1	(a)				volume O pols expla		er unit volur	ne			B1
	(b)	(i)	(vol = = 22.	=) mass/d 48 cm³	lensity O to 2 or r	R 60.7/2.3 nore sig. f	70 ïgs				C1 A1
		(ii)	OR 2	2.48 / (5	0 × 30)		thickness = figs. e.c.f. <b>(b</b>				C1 A1
	(c)	c) (i) micrometer/screw gauge / (vernier/digital) callipers						B1			
		(ii)		k zero of æ / fold s		sed / cut s	sheet into se	everal piece	es / detail of how	to use	B1
			measure thickness of sheet <u>in different places</u> OR measure thickness of several pieces together calculate/obtain average thickness OR divide answer by number of measuremen					asurements/	B1		
				es/places		<u> </u>			,		B1
										[Tot	tal 9]
2	(a)	a) underline or circle force underline or circle velocity					B1 B1				
	(b)	(i)	4.07	– 4.1 (s)							B1
		(ii)	OR o	other corr	ect value	words Of s from gra nd 10 m/s		÷ (ans. to <b>(</b>	b)(i))		C1 A1
		(iii) area under graph OR $\frac{1}{2}(u + v)t$ OR $\frac{1}{2} \times 40 \times (ans. to (b)(i))$ OR $s = ut + \frac{1}{2}at^2$ OR $v^2 = u^2 + 2as$ OR numbers substituted 82 m						C1 A1			
	(c)	c) graph continues in straight line to 6 s					B1				
										[Tot	tal 8]

	Page 5			Mark Scheme	Syllabus	Paper		
				IGCSE – May/June 2013	0625	31		
3	(a)	(i)		(loss of P.E. =) mgh OR 92 × 10 × 1500 1.38 × 10 <sup>6</sup> J correct use of mgh with $h = 500$ or 2000 gains 1 ma	irk only	C1 A1		
		. ,		(K.E. =) $\frac{1}{2} mv^2$ OR $\frac{1}{2} \times 92 \times 52^2$ 1.244 × 10 <sup>5</sup> J at least 2 sig. figs		C1 A1		
	(a)	<ul> <li>(ii) difference is due to: (work done in overcoming) air resistance/drag OR energy converted to/lost as heat (by air resistance/drag)</li> </ul>						
	(b)	(i)	incre	eases		B1		
		(ii)	920	Ν		B1		
						[Total 7]		
4	(a)	(i)		ntion of vacuum OR glass is a poor conductor vacuum/gap between walls has no molecules/atoms	s/particles	B1		
		(ii)		ace/silver (of walls) is good reflector/poor absorber ( ace/silver (of walls) is poor emitter (of radiation)	of radiation)	B1 B1		
	(b)			opper/lid/bung/cover/top to reduce/prevent (loss of h ion/radiation/evaporation OR to prevent steam/hot va		M1 B1		
			made of insulator OR example of insulator to reduce/prevent (loss of heat by) convection/radiation/evaporation OR to prevent steam/hot air leaving					
						[Total 6]		
5	(a)	(i) a (i)		<ul> <li>ii) marked together to maximum of 3 marks</li> <li>ecules escape/leave the liquid/form gas or vapour</li> </ul>		B1		
		(ii)		poration OR heat/(thermal) energy needed for evapo (er) molecules/high(er) energy molecules escape	pration leaves sweat	cooler B1		
			OR	slow(er) molecules left behind t flows from body to warm the sweat (so body cools)		B1 B1		
	(b)	(i)		=) <i>mc∆θ</i> OR <i>mcT</i> OR 60 × 4000 × 0.50 × 10 <sup>5</sup> J / 120 kJ		C1 A1		
		(ii)		<i>mL</i> in any form OR (m =) $Q/L$ OR either with numbe = $1.2 \times 10^5 / 2.4 \times 10^6$ =) 0.05 kg e.c.f from <b>(b)(i)</b>	ers	C1 A1		
						[Total 7]		

	Page 6			Mark Scheme	Syllabus	Paper		
				IGCSE – May/June 2013	0625	31		
6	(a)	(i) (ii)	expla mole	ssure =) force/area OR force per unit area OR ( <i>P</i> =) ained ecules collide with/hit walls/surface (of box)	<i>F/A</i> with symbols	B1 B1 B1		
		molecule(s) exert force on wall pressure is total force / force of all molecules divided by (total) area of wall						
	(b)	(i) $(P =) h\rho g$ OR in words OR $0.25 \times 13600 \times 10$ 34 000 Pa OR N/m <sup>2</sup> allow 1 mark for <i>h</i> = 250 used and $3.4 \times 10^7$ Pa obtained						
		<ul> <li>(ii) (P = 1.02 × 10<sup>5</sup> − 34 000)</li> <li>68 000 Pa or N/m<sup>2</sup></li> <li>e.c.f. from (b)(i) only if (b)(i) is less than 1.02 × 10<sup>5</sup></li> </ul>						
						[Total 7]		
7	(a)	two of: ray <u>through</u> centre of lens undeviated ray parallel to axis refracted to right hand focus rays through left hand focus refracted parallel to axis			B2			
		rays extrapolated to a point						
		accuracy marks: image 6 cm from lens image 6 cm high						
	(b)	) image is virtual/not real <u>AND</u> cannot be seen on screen OR no rays come from (position of) image				B1		

	Page 7		Mark Scheme	Syllabus	Paper
			IGCSE – May/June 2013	0625	31
8	<b>(a)</b> 15	–25 H	z to 15 000–25000 Hz / 15–25 kHz		B1
	(b) (i)	clos	ion) where air layers/molecules/particles are pushed er (than normal) (region) where (air) pressure reject/(air (more) comp	-	-
		UR	(region) where (air) pressure raised/air (more) comp	oressed/more dense	e di
	(ii)	· •	ion) where air layers/molecules are pushed apart/fa (region) where (air) pressure reduced/air expanded	r(ther) apart (than n	ormal) B1
	(c) (i)	(sou	nd is) loud(er) OR volume (of sound is) increased		B1
	(ii)	sour	nd has a higher frequency/pitch OR higher note (hea	ard)	B1
			0 OR 1.6 (s) seen OR <i>v</i> = 2 <i>d</i> /1.9 OR 500 (m) seen OR <i>v</i> = (2 <i>d</i> + 500)/3.5		C1 C1
			= 500 / 1.6 = 312.5  m/s at least 2 sig. figs		A1
					[Total 8]
9	(a) (i)	all la	amps off		
	(ii)	12 🖸	2 lamps (only) on		B1
	(iii)	4 Ω	lamps (only) on		
	(b) (i)	12 \	/		B1
	(ii)		//R in any form OR V/R OR 12/12		C1
			A OR 1 A . from <b>(b)(i)</b>		A1
	<b>(c)</b> cu	rrent iı	n 4 Ω lamp = 3 (A) (current in 12 Ω lamp is in <b>(b)(ii)</b> )	1	C1
	•	,	OR $I^2 R$ W for 4 $\Omega$ lamp; $P$ = 12 W for 12 $\Omega$ lamp		C1 A1
	è.	c.f. froi	m <b>(b)(ii)</b>		AI
	оғ ( <i>Р</i>	≺ '=) V²/	R		(C1)
			$r^{2}/4$ = 36 W for 4 Ω lamp OR 12 <sup>2</sup> /12 = 12 W for 12 Ω $r^{2}/4$ = 36 W for 4 Ω lamp AND 12 <sup>2</sup> /12 = 12 W for 12 Ω		(C1) (A1)
	Ò	ר <sup>(</sup> ד			. ,
	•	' =) <i>V</i> ²/ ame <i>V</i>	R for all lamps		(B1) (M1)
	4 9	$\Omega$ lamp	) has higher power / 12 $\Omega$ has lower power		(A1)
					[Total 7]

	Ра	ge 8		Mark Scheme	Syllabus	Paper
				IGCSE – May/June 2013	0625	31
10	(a)	arro	ows c	3 concentric circles centred on wire lockwise on each circle / at least one circle of circles increasing as radius increases		B1 B1 B1
	(b)	(i)	arro	w pointing down on side AB, up on side CD		B1
		(ii)	line OR I	es on AB and CD are opposite OR up and down and (so cause rotation) have moments in same sense / direction cause couple / torque	d separated / not i	n same B1
	C Vi C			everse current in loop or keep current in AB or CD in keep current on side near a pole in the same direction ical every half turn when AB and CD swap sides		) coil is
			so th rotat	when AB and CD swap sides nat: tion continues (in same direction) so that rotation doesn't reverse its direction		B1
				to maintain sense/direction of moments/couple coil turns more than half a revolution		B1
						[Total 7]
11	(a)	(i)	•	otons eutrons		B1 B1
		(ii)	a (fa	ast moving) electron		B1
	(b) eleo		ctron/	electrons removed from/gained by the molecule		B1
	(c)	OF		e because particle is charged the force on the particles is perpendicular to their pa direction of force changes as direction of motion cha		B1
				article <u>curve</u> up the page in at least half of width of f	ield	B1
				article $\underline{curve}$ opposite to $\alpha$ -particle curve OR down p ature anywhere	age if $\alpha$ line has r	no B1
			sma	ller radius of $\beta$ path clear		B1
						[Total 8]