

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

	CANDIDATE NAME		
	CENTRE CANDIDA NUMBER	TE	
* 0 5 0 1 6 1 0 4 3	PHYSICS Paper 2 Core Candidates answer on the Question Paper. No Additional Materials are required.	October/Nove 1 hour	0625/23 ember 2012 15 minutes
4	READ THESE INSTRUCTIONS FIRST	For Exami	ner's Use
	Write your Centre number, candidate number and name on all the work you hand in.	1	
	Write in dark blue or black pen. You may use a pencil for any diagrams or graphs. Do not use staples, paper clips, highlighters, glue or correction fluid.	2	
	DO NOT WRITE IN ANY BARCODES.	3	
	Answer all questions. You may lose marks if you do not show your working or if you do not use appropriate units.	4	
	Take the weight of 1 kg to be 10 N (i.e. acceleration of free fall = 10 m/s^2).	5	
	At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.	6	
		7	
		8	
		9	
		10	
		11	
		12	
		Total	

This document consists of **19** printed pages and **1** blank page.



1 Two boys, X and Y, decide to measure the speed of some of the vehicles travelling along a road. The two boys stand 405 m apart beside the road, as shown in Fig. 1.1.

For Examiner's Use





Fig. 1.1 (not to scale)

Boy X has a stopwatch which he sets to zero. As a vehicle passes boy Y, boy Y drops his hand as a signal to boy X to start his stopwatch. Boy X then stops the stopwatch as the vehicle goes past him.

The appearance of the stopwatch is then as shown in Fig. 1.2.





(a) How long did it take for the vehicle to travel from Y to X?

time =s [1]

(b) Calculate the average speed of the vehicle as it travels from Y to X.

average speed =[4]

(c)	The vehicle in (a)) and (b) is acceleratin	g as it travels from Y to X.
-----	----------------------------	---------------------------------	------------------------------

(i) How does its speed at X compare with that at Y?

Tick one box.

greater than at Y	
same as that at Y	
less than that at Y	

(ii) How does its speed at X compare with the average speed calculated in (b)?

Tick one box.

greater than average speed

same as average speed

less than average speed

[2]

For Examiner's Use

[Total: 7]

2 Solids, liquids and gases have different properties. The list below gives some of them.

	completely fills the container
shape	<pre>{ completely fills the container fills the container from the bottom fixed shape</pre>
	fixed shape
	move around, close together
molecules	<pre>{ move around, close together move around, far apart vibrate about a fixed position</pre>
	vibrate about a fixed position

Use descriptions from the list to complete the table. Any description may be used more than once if appropriate. Two spaces have been filled in to help you.

	shape	molecules	
(a) solid			[2]
(b) liquid		move around, close together	[1]
(c) gas	completely fills the container		[1]

[Total: 4]

3 Here is a list of energy resources available to the world. Some of these are renewable and some are non-renewable.

In the first blank column, put a tick by any **two** resources that are renewable.

In the second blank column, put a tick by any **two** resources that are non-renewable.

	renewable	non-renewable
coal		
hydroelectricity		
nuclear energy		
oil		
solar energy		
tidal energy		
wind energy		

[4]

For

Examiner's Use

[Total: 4]

4 An aluminium rod is cut into a longer section and a shorter section, as shown in Fig. 4.1.




Fig. 4.1

(a) The shorter section of the rod is placed into a measuring cylinder containing water.

Fig. 4.2 shows the appearance of the measuring cylinder before and after this is done.




Fig. 4.2 (not full size)

(i) Calculate the volume of the shorter section of aluminium rod.

volume = cm³ [3]

https://xtremepape.rs/

	(ii)	The mass of this shorter section is measured as 21.2 g.	For Examiner's
		1. Name a laboratory instrument that might have been used to measure this mass.	Use
		[1]	
		2. Calculate the density of aluminium.	
		density =[4]	
(b)	(i)	Name an instrument that could be used to measure the length of the longer section of aluminium rod.	
		[1]	
	(ii)	Suggest a method, different from that in Fig. 4.2, that could be used to determine the volume of this longer section.	
		[2]	

5 (a) In Fig. 5.1, A and B are two parallel plane mirrors. A ray of light strikes mirror A at an angle of incidence of 45°. The ray then reflects, to strike mirror B.




Fig. 5.1

- (i) State the name given to the angle *x* shown on Fig. 5.1.
 (ii) State the value of

 angle *x*,

 (iii) 2. Fig. 5.4 and the state of the stat
- (iii) On Fig. 5.1, use your ruler to draw the path of the ray after it leaves the surface of B. [1]

https://xtremepape.rs/

(b) The mirror B is now rotated so that it reflects the ray of light back along its original path. On Fig. 5.2, draw mirror B in the correct position to do this.




Fig. 5.2

[1]

[Total: 5]

Some water in a glass beaker is heated from below, as shown in Fig. 6.1. 6






	Fig. 6.1		
(a)	Nar	me the process by which therma	l energy is transferred
	(i)	through the glass,	
	(ii)	throughout the water	[2]
(b)	the		e temperature of the water begins to rise. Although tant, eventually the temperature becomes steady at
	Sug	ggest why this happens.	
			[1]
(c)	aga		ased. The temperature of the water begins to rise dy at a higher temperature. This time many bubbles
	(i)	State what is now happening to	o the water.
			[1]
	(ii)	What gas do the bubbles conta	ain? Tick one box.
		air	
		hydrogen	
		oxygen	
		steam	[1]
			[Total: 5]

https://xtremepape.rs/

			••	
7	(a)	Sta	te what is meant by the <i>echo</i> of a sound.	For
				Examiner's Use
			[2]	
	(b)		scribe how the echo of a sound may be demonstrated. Include a diagram that shows proximate sizes and distances.	
		dia	gram	
		des	cription of method	
			[3]	
	(c)	The	e demonstration in (b) is used to find the speed of sound in air.	
		(i)	Which two measurements should be made?	
		()		
			1	
			2[2]	
		<i>(</i> ;;)		
		(ii)	State how you would calculate the speed of sound from these measurements.	
			[2]	
			[Total: 9]	
				1

https://xtremepape.rs/

8 Fig. 8.1 represents the circuit that operates two of the lamps on a car.




Fig. 8.1

(a) In the space below, draw the circuit diagram for this circuit, using conventional symbols.

[3]

(b) The car battery has an e.m.f. of 12V and, when the lamps are switched on, there is a current of 1.6A in each lamp.

Calculate the resistance of one of the lamps.

resistance =[4]

https://xtremepape.rs/

(c) When the switch is turned on, both lamps should light up. On one occasion when the driver operates the switch, lamp L₂ fails to light up. Examiner's Suggest a reason for this.[1] (d) An amateur workman connects a length of wire across lamp L_2 and shorts it out. When the switch is closed for the first time after this, what happens, if anything, to (i) the fuse, (ii) lamp L₁, lamp L₂? (iii) [3]

13

[Total: 11]

For

Use

9 (a) Magnets A and B, shown in Fig. 9.1, attract each	h other
--	---------



Fig. 9.1

The S pole of magnet A has been marked.

On Fig. 9.1, mark the polarities of the other poles, using the letters N or S. [1]

(b) A soft-iron rod and a steel rod each have coils around them. Both rods are initially unmagnetised. The coils are attached to circuits, as shown in Fig. 9.2.




Fig. 9.2

(i) Use the following statements to complete the table referring to the soft-iron rod and the steel rod shown in Fig. 9.2.

magnetised

loses its magnetism keeps its magnetism

	switch closed	switch open
soft iron		
steel		

[2]


For Examiner's Use

(ii) Which words apply to the force between the rods when the switches are closed?


Tick one box.

no force
attractive force
repulsive force

[1]



(d)	The positively-charged plastic rod is removed and replaced by a plastic rod with a negative charge.	For Examiner's Use
	Describe the position that the suspended sphere now takes.	036
	[1]	
	[Total: 6]	



- 1000 count rate counts/s 800 600 400 200-40 20 50 10 30 60 time/s Fig. 12.1 (a) From Fig. 12.1, find the time taken for the count rate to decrease from 1000 counts/s to 125 counts/s. time = s [1]
- **12** Fig. 12.1 shows the graph of the count rate from a radioactive source over a period of time. The readings have already had the background count rate subtracted.

For Examiner's Use

(b) How many half-lives of the radioactive material were there during the time interval in (a)?

number of half lives =[1]

(c) From your answers to (a) and (b), calculate the half-life of the material.

half-life = s [2]

(d) On Fig. 12.1, sketch the curve that might have been plotted if the background count rate had not been subtracted. [1]

[Total: 5]

BLANK PAGE

20

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.