

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		



COMBINED SCIENCE

0653/52

Paper 5 Practical Test

May/June 2023

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- Notes for use in qualitative analysis are provided in the question paper.

For Examiner's Use		
1		
2		
3		
4		
Total		

This document has 12 pages. Any blank pages are indicated.

DC (DE/SG) 316016/2 © UCLES 2023

[Turn over

You are going to investigate okra, a fruit which contains many seeds.

	You are provided with a section of okra on a white tile. (a) In the box provided, draw a large, clear pencil drawing of the cut surface of the okra.				

[3]

© UCLES 2023 0653/52/M/J/23

1

(b)	(i)	Measure the diameter of the section of okra on the white tile.
		diameter of okra on white tile = mm [1]
	(ii)	Suggest why it is difficult to measure the diameter of the okra accurately.
		[1]
	(iii)	Measure the diameter of your drawing in (a) . Draw a line on your drawing to show where you have measured.
		diameter of your drawing = mm [1]
	(iv)	Calculate the magnification of your drawing.
		Use the equation shown.
		$magnification = \frac{\text{diameter of your drawing}}{\text{diameter of okra on white tile}}$
		magnification =[1]
		[Total: 7]

2 The enzyme amylase breaks down starch to form a reducing sugar.

Plan an investigation to determine the relationship between temperature and the time taken to completely break down starch by amylase. Iodine is a brown solution that turns blue/black in the presence of starch.

You are provided with:

- 1% amylase solution
- 1% starch solution
- iodine solution

You may also use any other common laboratory apparatus.

You are not required to do this investigation.

In your plan include:

- the additional apparatus needed
- a brief description of the method and an explanation of any safety precautions you will take
- what you will measure
- which variables you will keep constant
- how you will process your results to draw a conclusion.

You may include a labelled diagram if you wish.

You may include a table that can be used to record the results if you wish.

You do **not** need to include any results in your results table.

[7]
17

3 You are going to investigate a white solid **H**.

(a) (i) Procedure

- Measure the mass of the clean dry test-tube labelled H.
- Record this mass in Table 3.1.
- Place two spatula loads of solid H into the test-tube.
- Measure the mass of the test-tube and solid H.
- Record this mass in Table 3.1.
- Using the test-tube holder, heat solid **H** safely for one minute using a blue Bunsen burner flame.
- Observe solid H during heating.
- Lay the test-tube on the laboratory mat and allow the test-tube to cool down.

Table 3.1

	mass /g
empty test-tube	
test-tube and solid H before heating	
test-tube and the solid after heating	

		[2]
(ii)	Describe your observation of solid H during heating.	
		[1]
	While you are waiting for the test-tube to cool down do part (b).	
(iii)	When the test-tube is cool, measure the mass of the test-tube and the solid after heat	ing.
	Record this mass in Table 3.1.	[1]
(iv)	Describe your observation of the solid after cooling.	
		[1]
(v)	Calculate the mass of solid H in the test-tube before heating.	
	Use the equation shown.	
	mass of solid H before heating = mass of test-tube and solid H before heating - mass of empty test-tube	

mass of solid **H** before heating = g [1]

	(vi)	Calculate the mass of the solid in the test-tube after heating.
		Use the equation shown.
		mass of the solid after heating = mass of test-tube and the solid after heating - mass of empty test-tube
		mass of the solid after heating = g [1]
	(vii)	There is a loss in mass when solid H is heated.
		Suggest one reason for this loss in mass.
		[1]
((viii)	Calculate the percentage loss in mass.
		Use the equation shown.
ре	ercent	age loss in mass = $\frac{\text{mass of solid } \mathbf{H} \text{ before heating} - \text{mass of the solid after heating}}{\text{mass of solid } \mathbf{H} \text{ before heating}} \times 100$
		mass of solid if before fleating
		Give your answer to two significant figures.
		percentage loss in mass =[2]
	(ix)	Explain why it is a good idea to heat solid H for at least five minutes rather than one minute.
		[1]
	(x)	State one reason why the test-tube must be heated with a blue Bunsen burner flame rather than a yellow Bunsen burner flame.
		[1]
(b)	Put	about 3 cm depth of dilute hydrochloric acid in a clean test-tube.
	Add	I one spatula load of solid H .
	Des	scribe one observation.
		[1]
	Go	back to (a)(iii) to finish question (a).
		[Total: 13]
		Tiolai. To

4 You are going to measure the focal length *F* of a convex lens.

Arrange the equipment as shown in Fig. 4.1.




Fig. 4.1

(a) Procedure

- Switch on the lamp.
- Place the illuminated object (a triangle) at the 0 cm mark on the metre rule.
- Place the lens at a distance $u = 10.0 \,\mathrm{cm}$ from the illuminated object.
- Place the screen at a distance $D = 95.0 \,\mathrm{cm}$ from the illuminated object.

An out of focus, fuzzy image appears on the screen.

- Move the lens slowly towards the screen until the image formed is in focus, and as sharp as possible.
- (i) Measure the distances u and v to the nearest 0.1 cm.

Record *u* and *v* in the first row of Table 4.1.

Table 4.1

D /cm	u /cm	v /cm	uv /
95.0			
85.0			
75.0			
70.0			
65.0			

[1]

(ii) Repeat the measurements for the four other values of *D* shown in Table 4.1.

Record the distances *u* and *v* in Table 4.1 against the correct values of *D*.

[2]

(iii) Calculate the product *uv* and record it for each value of *D* in the final column of Table 4.1. Use the equation shown.

$$uv = u \times v$$


[1]

(iv) Add the unit to the column heading for uv in Table 4.1.

[1]

(b) (i) On the grid, plot a graph of *uv* (vertical axis) against *D*.

You do **not** need to start your graph from the origin (0, 0).



[3]

(ii) Draw the best-fit straight line.

[1]

(c) The focal length F of the lens is equal to the gradient of your line.

Indicate on your graph the values you choose to calculate the gradient.

Calculate the gradient of your line.

	F =[2]
(d) (i)	F can also be calculated without plotting a graph, by using the results for one value of D.
	Suggest why plotting a graph and calculating a gradient to find the value of F gives a more accurate answer than calculating F for one value of D .
	[1]
(ii)	State one precaution that you take when doing the experiment to make your readings as accurate as possible.
	[1]
	[Total: 13]

BLANK PAGE

NOTES FOR USE IN QUALITATIVE ANALYSIS

Tests for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide, then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify, then add aqueous barium nitrate	white ppt.

Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
ammonium (NH ₄ ⁺)	ammonia produced on warming	-
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt. or very slight white ppt.
copper(II) (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess, giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess, giving a colourless solution	white ppt., soluble in excess, giving a colourless solution

Tests for gases

gas	test and test result
ammonia (NH ₃)	turns damp red litmus paper blue
carbon dioxide (CO ₂)	turns limewater milky
chlorine (Cl ₂)	bleaches damp litmus paper
hydrogen (H ₂)	'pops' with a lighted splint
oxygen (O ₂)	relights a glowing splint

Flame tests for metal ions

metal ion	flame colour
lithium (Li ⁺)	red
sodium (Na ⁺)	yellow
potassium (K ⁺)	lilac
copper(II) (Cu ²⁺)	blue-green

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.