

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge International General Certificate of Secondary Education

MARK SCHEME for the October/November 2015 series

0654 CO-ORDINATED SCIENCES

0654/31

Paper 3 (Extended Theory), maximum raw mark 120

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0654	31

- 1 (a) (i) 8 ; [1]
- (ii) neutron ; [1]
- (iii) 15 electrons ;
arranged 2.8.5 ; [2]
- (b) 3 shared pairs ;
1 lone pair on central atom and no extra electrons ;
(max 1 if symbols missing or incorrect) [2]
- (c) (i) Haber (process) ; [1]
- (ii) $\text{CH}_4 + \text{H}_2\text{O} \rightarrow 3\text{H}_2 + \text{CO}$
1 mark for H_2 ; 1 mark for CO ; 1 mark for fully correct ; [3]
- (iii) catalyst / to speed up the reaction / to facilitate reaction ; [1]
- [Total: 11]**
- 2 (a) chloroplast ; [1]
- (b) light ;
chemical ; [2]
- (c) (i) (oxygen) from photosynthesis ;
(carbon dioxide) from respiration ;
(nothing) because rate of photosynthesis equals rate of respiration ; [3]
- (ii) dead / no chloroplasts ; [1]
- [Total: 7]**
- 3 (a) **B** (no mark)
particles are touching and randomly arranged ; [1]
- (b) (i) warmer ;
larger surface area ;
faster air flow ; [max 1]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0654	31

- (ii) evaporation can occur at any temperature (above melting point)/boiling only happens at the boiling point ;
 evaporation happens only at the surface/boiling happens throughout the liquid ;
 boiling takes energy in (endothermic) to occur/evaporation lets only the molecules with the highest kinetic energy out ;
 evaporation can occur using the internal energy of the system/boiling requires an external source of heat ;
 evaporation produces cooling/boiling does not ;
 evaporation is a slow process/boiling is a rapid process ; [max 2]

(c) (i) (energy =) power \times time ;
 $= 18\,000 \times 3600 = 64\,800\,000 \text{ J}$ or $18 \times 3600 = 64\,800 \text{ kJ}$; [2]

- (ii) when voltage is high, current is lower ;
 less energy is transferred as thermal energy ; [2]

- (iii) lowers the voltage/has less turns on secondary coil than primary ; [1]

[Total: 9]

- 4 (a) a change in a gene or a chromosome ; [1]

- (b) (i) mutation in the parents ;
passed on to offspring in reproduction ; [2]

- (ii) ionising radiation/ γ /X-rays/ultraviolet rays ; [1]

- (iii) less able to find food/find a mate/escape predators ; [1]

- (c) adapted ;
 survive ;
 alleles ;
 selection ; [4]

[Total: 9]

- 5 (a) (i) (with propane) no change/no reaction ;
 (with propene) bromine solution decolourised ; [2]

- (ii) propene molecules contain double bond propane all single bonds/propene contains fewer hydrogen atoms/correct formulae given and assigned ; [1]

- (b) (i) goes milky (cloudy)/goes milky then clears ;
 it is reacting with carbon dioxide/the reaction gives off carbon dioxide ; [2]

- (ii) $(12 \times 6) + (1 \times 12) + (16 \times 6) = 180$; [1]

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0654	31

- (iii) idea that moles dissolved = volume \times concentration / so may see
 moles = $5.0 \times 3.5 = 17.5$ moles ;
 then required mass = moles \times molar mass / so may see
 mass = $17.5 \times 180 = 3150$ (g) **or** 3.15 kg ;
 ($5.0 \times 3.5 \times 180 = 3150$ (g) *award 2 marks*)

OR

- mass in $1 \text{ dm}^3 = 3.5 \times 180 = 630$ g ;
 mass in $5 \text{ dm}^3 = 630 \times 5 = 3150$ (g) ;

[max 2]

(c) (i) nitrogen ; [1]

(ii) protein / polypeptide ; [1]

[Total: 10]

- 6 (a) rays hit wall at angle greater than critical angle ;
 only reflection / no refraction / no light exiting side of fibre ;
 rays undergo total internal reflection at walls of fibre ;

[max 2]

(b) (i) can pass through tissue ;
 less ionising so less damage caused ;

[max 1]

(ii) 13 (hours) ; [1]

(iii) 4 half-lives ;
 50 (counts per minute) ; [2]

[Total: 6]

7 (a) any part of the nervous system except brain / spinal cord ; [1]

(b) (i) response to a stimulus / response to protect body ;
 immediate / automatic / without conscious thought ; [2]

(ii) carry impulses / AW from receptors to CNS ;
 carry impulses / AW from CNS to effectors / muscle ;
 reference to sensory neurons / motor neurons ; [max 2]

(c) (i) (nervous system is) shorter lasting ; [1]

(ii) nervous system has electrical impulses ;
 hormones are chemicals carried in blood ; [2]

[Total: 8]

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0654	31

- 8 (a) (i) less attraction / filler not magnetic but steel is / owtte ; [1]
- (ii) no – aluminium is not magnetic ; [1]
- (b) (i) $I = \frac{V}{R}$;
 $= \frac{12}{2.5} = 4.8 \text{ (A)}$;
amps/A ; [3]
- (ii) (charge =) current \times time ;
 $= 4.8 \times 2 \times 60 = 576 \text{ (C)}$; [2]
- (iii) use of $\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2}$;
 $R_T = 1.25 \text{ (}\Omega\text{)}$; [2]
- (c) (energy =) SHC \times mass \times change in temperature ;
 $= 4200 \times 4 \times 80 = 1\,344\,000 \text{ (J)}$; [2]

[Total: 11]


- 9 (a) electrolysis ; [1]
- (b) (i) Al ions are positive / opposite charges attract ; [1]
- (ii) each Al ion gains electrons ;
ions are discharged ;
(each ion gains 3 electrons, award 2 marks) [2]
- (c) (i) Fe^{3+} ;
reference to charge balance / $3 \times 2-$ balanced by $2 \times 3+$ / owtte ; [2]
- (ii) iron more reactive than copper / aluminium more reactive than copper
(from own knowledge of reactivity series) ;
since Al more reactive than iron it must be more reactive than copper
(from information in question) ;
(so Al does displace Cu) [2]

[Total: 8]

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0654	31

10 (a) mayfly larvae / caddis flies / freshwater shrimps / water lice / bloodworms ; [1]

(b) (i) arrow anywhere in the shaded area ; [1]



(ii) microorganisms ;
 respiration deoxygenates water ;
 which prevents respiration ;
 toxic ;
 heavy metals bioaccumulation ; [max 3]

(c) (i) rain of low pH / pH less than 7 / polluted with (named) acid ; [1]

(ii) reduced use of fossil fuels ;
 public transport ;
 alternative energy sources ;
 (chemical) absorbers / filters on (factory) chimneys ;
 education / taxation / public awareness measures ; [max 2]

[Total: 8]

11 (a) (KE =) $\frac{1}{2} mv^2$;
 $= \frac{1}{2} \times 4000 \times 0.4 \times 0.4 = 320$ (J) ; [2]

(b) (work done =) force \times distance ;
 $= 3000 \times 2 = 6000$ (J) ; [2]

(c) (i) (pressure =) $\frac{\text{force}}{\text{area}}$;
 $\frac{40\,000}{1600} = 25$ (N/cm²) ; [2]

(ii) 250 000 (Pa) ; [1]

Page 7	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0654	31

(d) (i) (*higher than 30 Hz – no mark*)
lowest frequency detected is 10–30 Hz ; [1]

(ii) particles vibrate ;
(particles vibrate) parallel to direction of sound travel/energy transfer ;
compressions and rarefactions ;
description of compressions/rarefactions ; [max 2]

(e) (time =) $\frac{\text{distance}}{\text{speed}}$;
 $\frac{6000}{330} = 18.(18)$ (s) ; [2]

(f) eureka can / displacement method ;
volume of water displaced is the volume of the object ; [2]

[Total: 14]

12 (a) magnesium + sulfuric acid ;
zinc carbonate + sulfuric acid → (*zinc sulfate + carbon dioxide +*) water ; [2]

(b) (i) thermal energy → chemical (potential) energy ; [1]

(ii) reaction is endothermic/temperature decreases ; [1]

(c) (i) no gas produced / gas stops after 75 s ;
because reaction is complete / all the calcium carbonate has reacted ; [2]

(ii) generally similar shape ;
everywhere below original curve ;
maximum volume of gas at 45 to 50 cm³ ; [3]

(iii) (kinetic) energy / speed of (acid) particles increases ;
increases the frequency of collision / more successful collisions ; [2]

[Total: 11]

13 (a) anther correctly labelled (at the top) ; [1]

(b) pollen ;
male gamete ; [2]

Page 8	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – October/November 2015	0654	31

(c) large/bright petals ;
scent ;
nectar ;
flower parts / anthers / stigmas inside the flower ;
sticky pollen ;

[max 2]

(d) (i) by animals ;
hook to attach to fur / eaten and egested ;

[2]

(ii) seed / embryo ;

[1]

[Total: 8]