

CANDIDATE  
NAME


--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**CHEMISTRY**

Paper 2 Theory

**5070/21**

**May/June 2018**

**1 hour 30 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

**Section A**

Answer **all** questions.

Write your answers in the spaces provided in the Question Paper.

**Section B**

Answer any **three** questions.

Write your answers in the spaces provided in the Question Paper.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of **19** printed pages and **1** blank page.



**Section A**

Answer **all** the questions in this section in the spaces provided.

The total mark for this section is 45.

1 Choose from the gases to answer the questions.

- argon
- carbon dioxide
- carbon monoxide
- ethane
- hydrogen
- methane
- neon
- ozone
- sulfur dioxide
- sulfur trioxide

Each gas can be used once, more than once or not at all.

Which gas:

- (a) is used to manufacture margarine  
.....[1]
- (b) is used as a food preservative  
.....[1]
- (c) has a molecule that contains only eight atoms  
.....[1]
- (d) is formed from bacterial decay of vegetable matter  
.....[1]
- (e) is formed in photochemical smog  
.....[1]
- (f) completely combusts to form only a gas that turns limewater milky?  
.....[1]


[Total: 6]

2 The elements in Group VII of the Periodic Table are called the halogens.

(a) Explain why the elements in Group VII have similar chemical properties.

.....  
 ..... [1]

(b) A redox reaction happens when chlorine gas is bubbled through aqueous potassium iodide.



(i) Describe what is observed during this reaction.

..... [1]

(ii) Use the equation to explain that oxidation takes place in this reaction.

.....  
 ..... [1]

(iii) Use the equation to explain that reduction takes place in this reaction.

.....  
 ..... [1]

(c) Describe the chemical test for chlorine gas.

test .....

observation .....

..... [2]

(d) Chlorine reacts with iron to form iron(III) chloride in a closed container.

(i) The pressure of chlorine is increased.

Describe and explain what happens to the rate of this reaction.

.....  
 .....  
 .....  
 ..... [2]

(ii) Iron(III) chloride can act as a catalyst for some reactions.

Explain how a catalyst increases the rate of a reaction.

.....

.....

.....

..... [2]

[Total: 10]

3 Barium chloride is a soluble salt and barium sulfate is an insoluble salt.

(a) Barium sulfate can be prepared by the reaction between aqueous barium chloride and dilute sulfuric acid.

(i) Describe the preparation of a pure, dry sample of barium sulfate from aqueous barium chloride and dilute sulfuric acid.

.....  
 .....  
 .....  
 .....  
 ..... [3]

(ii) Write the ionic equation, including state symbols, for this reaction.

..... [2]

(b) Barium chloride can be prepared by reacting barium carbonate with dilute hydrochloric acid.



Excess barium carbonate is reacted with 40.0 cm<sup>3</sup> of 1.50 mol/dm<sup>3</sup> hydrochloric acid.

After purification the percentage yield of barium chloride was 75.0%.

Calculate the mass of barium chloride prepared.

Give your answer to **three** significant figures.

[*M<sub>r</sub>*: BaCl<sub>2</sub>, 208]

mass of barium chloride ..... g [3]

(c) A barium ion has the formula  ${}^{138}_{56}\text{Ba}^{2+}$ .


Complete the table about this ion.

subatomic particles	number of subatomic particles
electrons	.....
neutrons	.....
protons	.....


[3]

[Total: 11]

4 This question is about ethanol and ethanoic acid.



ethanol



ethanoic acid

(a) Describe the manufacture of ethanol by fermentation.

.....

.....

.....

.....

.....

.....

.....

.....

.....[3]

(b) Ethanol can be oxidised by oxygen to form ethanoic acid.

Construct the equation for this reaction.

.....[1]

(c) A sample of ethanol is heated with ethanoic acid in the presence of concentrated sulfuric acid catalyst.


Draw the structure of the product of this reaction.

Show all of the atoms and all of the bonds.

[1]



(d) Ethanol is used as a fuel.



The complete combustion of one mole of ethanol releases 1350 kJ of energy.

A sample of ethanol reacts with excess oxygen to make 0.240 dm<sup>3</sup> of carbon dioxide, measured at room temperature and pressure.


Calculate the energy released, in kJ, in this reaction.

energy released ..... kJ [2]

[Total: 7]

5 Silicon is a non-metal found in Group IV of the Periodic Table.

(a) Silicon is manufactured by the reduction of silicon dioxide with carbon.



What is the maximum mass of silicon that can be made from 300g of silicon dioxide?

mass of silicon ..... g [2]

(b) Silicon has a giant molecular structure.

Suggest **two** physical properties of silicon.

1 .....

2 .....

[2]

(c) Silane has the molecular formula  $\text{SiH}_4$ .

(i) Draw the 'dot-and-cross' diagram for silane.

You only need to show the outer shell electrons of silicon.


[1]

- (ii) Using ideas about structure and bonding, suggest why silane has a low boiling point.

.....  
 .....  
 .....  
 ..... [2]

- (d) Siloxanes are condensation polymers.

The partial structure of a siloxane can be represented as shown.




- (i) What is meant by the term *condensation polymerisation*?

.....  
 .....  
 .....  
 ..... [2]

- (ii) Draw the repeat unit of this siloxane.

[1]

(e) Butene forms an addition polymer called poly(butene).



Draw a partial structure for poly(butene). You must show at least two repeat units.

[1]

[Total: 11]

## Section B

Answer **three** questions from this section in the spaces provided.

The total mark for this section is 30.

- 6 Copper pyrites is an ore containing compounds of copper. One of the compounds in the ore is  $\text{CuFeS}_2$ .

- (a) Calculate the mass of copper in 20.0 tonnes of  $\text{CuFeS}_2$ .

mass of copper ..... tonnes [2]

- (b)  $\text{CuFeS}_2$  is heated in air. Copper(II) oxide, iron(III) oxide and sulfur dioxide are formed.

Construct the equation for this reaction.

.....[2]

- (c) Copper(II) oxide can be reduced by heating with carbon to form copper.

Construct the equation for this reaction.

.....[1]

- (d) Copper made by this reduction is impure.

Impure copper can be purified using electrolysis.

- (i) Name the electrolyte used.

.....[1]

- (ii) What are the electrodes made from?

positive electrode (anode) .....

negative electrode (cathode) .....

[2]

- (iii) Write the equations for the reactions at both electrodes.

positive electrode (anode) .....

negative electrode (cathode) .....


[2]

[Total: 10]

**[Turn over**

7 A scientist investigates the reaction of iron with steam in a closed system.

A dynamic equilibrium mixture is established.



(a) Explain why the concentrations of steam and of hydrogen do not change once the dynamic equilibrium mixture has been established.

.....  
 .....  
 ..... [1]

(b) The pressure of the equilibrium mixture is increased.

The temperature of the closed system is kept constant.

Predict and explain what will happen, if anything, to the composition of the equilibrium mixture.

.....  
 .....  
 .....  
 ..... [2]

(c) The temperature of the equilibrium mixture is increased.

The pressure within the closed system is kept constant.

(i) The position of equilibrium shifts to the left hand side.

What conclusion can be made about the enthalpy change of the reaction?

..... [1]

(ii) Describe and explain what happens to the rate of reaction.

.....  
 .....  
 ..... [1]

(d) Dilute sulfuric acid reacts with  $\text{Fe}_3\text{O}_4$  to form three compounds, **A**, **B** and **C**.

- **A** is iron(II) sulfate.
- **B** is iron(III) sulfate.
- **C** is a colourless liquid.

(i) Name compound **C**.

.....[1]

(ii) Construct the equation for this reaction.

.....[2]

(iii) Describe a chemical test for iron(III) ions.


test .....

observation .....

.....[2]

[Total: 10]

8 Methanesulfonic acid has the structure shown.



(a) Write the molecular formula for methanesulfonic acid.

.....[1]

(b) Methanesulfonic acid is a stronger acid than ethanoic acid.

Explain the meaning of this statement.

.....  
 .....  
 .....[1]

(c) What is the difference between an aqueous acid and an aqueous alkali, in terms of the ions present?

.....  
 .....  
 .....[2]

(d) What is the mass of methanesulfonic acid needed to make  $150\text{cm}^3$  of a  $0.150\text{mol/dm}^3$  solution?

mass ..... g [3]



- (e) In a titration, 0.00150 moles of NaOH is exactly neutralised by 0.175 mol/dm<sup>3</sup> methanesulfonic acid.

One mole of sodium hydroxide reacts with one mole of methanesulfonic acid.

Calculate the volume, in cm<sup>3</sup>, of aqueous methanesulfonic acid needed in this titration.

volume ..... cm<sup>3</sup> [1]

- (f) Methanesulfonic acid reacts with magnesium to make a salt and a gas.

(i) Name the gas made in this reaction.

.....[1]


(ii) The formula of the anion in the salt is CH<sub>3</sub>SO<sub>3</sub><sup>-</sup>.

Write the formula of the salt formed.

.....[1]

[Total: 10]

9 Methylpropane is a saturated hydrocarbon.



methylpropane

(a) Methylpropane reacts with chlorine in the presence of ultraviolet light to give several compounds.

(i) One of these compounds has a relative molecular mass of 127.

What is the molecular formula of this compound?

molecular formula .....

Suggest a structure for this compound. Show all of the atoms and all of the bonds.

[2]

(ii) Another compound formed has the percentage composition by mass:

29.7% carbon; 4.3% hydrogen; 65.9% chlorine.

Calculate the molecular formula for this compound.


molecular formula ..... [2]

- (iii) A compound is formed when one molecule of methylpropane reacts with five molecules of chlorine.


Write the molecular formula for this compound.

.....[1]

- (b) The structures of propene and cyclopropane are shown.



propene



cyclopropane

These two compounds are isomers of each other.

- (i) What is meant by the term *isomerism*?

.....  
 .....[1]

- (ii) Both compounds are hydrocarbons.

What is meant by the term *hydrocarbons*?

.....  
 .....[1]

- (iii) Propene is an unsaturated hydrocarbon and cyclopropane is a saturated hydrocarbon.

What is the difference between an unsaturated and a saturated hydrocarbon?

.....  
 .....[1]

- (iv) Describe a chemical test to distinguish between unsaturated and saturated hydrocarbons.

.....  
 .....  
 .....[2]

[Total: 10]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cie.org.uk](http://www.cie.org.uk) after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

## The Periodic Table of Elements

		Group							
I	II	III	IV	V	VI	VII	VIII		
3 <b>Li</b> lithium 7	4 <b>Be</b> beryllium 9	1 <b>H</b> hydrogen 1	5 <b>B</b> boron 11	6 <b>C</b> carbon 12	7 <b>N</b> nitrogen 14	8 <b>O</b> oxygen 16	9 <b>F</b> fluorine 19	10 <b>Ne</b> neon 20	
11 <b>Na</b> sodium 23	12 <b>Mg</b> magnesium 24	<b>Key</b> atomic number atomic symbol name relative atomic mass		13 <b>Al</b> aluminium 27	14 <b>Si</b> silicon 28	15 <b>P</b> phosphorus 31	16 <b>S</b> sulfur 32	17 <b>Cl</b> chlorine 35.5	18 <b>Ar</b> argon 40
19 <b>K</b> potassium 39	20 <b>Ca</b> calcium 40	21 <b>Sc</b> scandium 45	22 <b>Ti</b> titanium 48	23 <b>V</b> vanadium 51	24 <b>Cr</b> chromium 52	25 <b>Mn</b> manganese 55	26 <b>Fe</b> iron 56	27 <b>Co</b> cobalt 59	28 <b>Ni</b> nickel 59
37 <b>Rb</b> rubidium 85	38 <b>Sr</b> strontium 88	39 <b>Y</b> yttrium 89	40 <b>Zr</b> zirconium 91	41 <b>Nb</b> niobium 93	42 <b>Mo</b> molybdenum 96	43 <b>Tc</b> technetium —	44 <b>Ru</b> ruthenium 101	45 <b>Rh</b> rhodium 103	46 <b>Pd</b> palladium 106
55 <b>Cs</b> caesium 133	56 <b>Ba</b> barium 137	57–71 lanthanoids	72 <b>Hf</b> hafnium 178	73 <b>Ta</b> tantalum 181	74 <b>W</b> tungsten 184	75 <b>Re</b> rhenium 186	76 <b>Os</b> osmium 190	77 <b>Ir</b> iridium 192	78 <b>Pt</b> platinum 195
87 <b>Fr</b> francium —	88 <b>Ra</b> radium —	89–103 actinoids	104 <b>Rf</b> rutherfordium —	105 <b>Db</b> dubnium —	106 <b>Sg</b> seaborgium —	107 <b>Bh</b> bohrium —	108 <b>Hs</b> hassium —	109 <b>Mt</b> meitnerium —	110 <b>Ds</b> darmstadtium —
							112 <b>Cn</b> copernicium —	111 <b>Rg</b> roentgenium —	110 <b>Ds</b> darmstadtium —
							80 <b>Hg</b> mercury 201	79 <b>Au</b> gold 197	78 <b>Pt</b> platinum 195
							48 <b>Zn</b> zinc 65	47 <b>Cu</b> copper 64	46 <b>Pd</b> palladium 106
							30 <b>Zn</b> zinc 65	29 <b>Cu</b> copper 64	28 <b>Ni</b> nickel 59
							31 <b>Ga</b> gallium 70	30 <b>Zn</b> zinc 65	29 <b>Cu</b> copper 64
							32 <b>Ge</b> germanium 73	31 <b>Ga</b> gallium 70	30 <b>Zn</b> zinc 65
							33 <b>As</b> arsenic 75	32 <b>Ge</b> germanium 73	31 <b>Ga</b> gallium 70
							34 <b>Se</b> selenium 79	33 <b>As</b> arsenic 75	32 <b>Ge</b> germanium 73
							51 <b>Sb</b> antimony 122	50 <b>Sn</b> tin 119	49 <b>In</b> indium 115
							52 <b>Te</b> tellurium 128	51 <b>Sb</b> antimony 122	50 <b>Sn</b> tin 119
							83 <b>Bi</b> bismuth 209	82 <b>Pb</b> lead 207	81 <b>Tl</b> thallium 204
							84 <b>Po</b> polonium —	83 <b>Bi</b> bismuth 209	82 <b>Pb</b> lead 207
							116 <b>Lv</b> livermorium —	115 <b>Po</b> polonium —	114 <b>Fl</b> flerovium —

57 <b>La</b> lanthanum 139	58 <b>Ce</b> cerium 140	59 <b>Pr</b> praseodymium 141	60 <b>Nd</b> neodymium 144	61 <b>Pm</b> promethium —	62 <b>Sm</b> samarium 150	63 <b>Eu</b> europium 152	64 <b>Gd</b> gadolinium 157	65 <b>Tb</b> terbium 159	66 <b>Dy</b> dysprosium 163	67 <b>Ho</b> holmium 165	68 <b>Er</b> erbium 167	69 <b>Tm</b> thulium 169	70 <b>Yb</b> ytterbium 173	71 <b>Lu</b> lutetium 175
89 <b>Ac</b> actinium —	90 <b>Th</b> thorium 232	91 <b>Pa</b> protactinium 231	92 <b>U</b> uranium 238	93 <b>Np</b> neptunium —	94 <b>Pu</b> plutonium —	95 <b>Am</b> americium —	96 <b>Cm</b> curium —	97 <b>Bk</b> berkelium —	98 <b>Cf</b> californium —	99 <b>Es</b> einsteinium —	100 <b>Fm</b> fermium —	101 <b>Md</b> mendelevium —	102 <b>No</b> nobelium —	103 <b>Lr</b> lawrencium —

lanthanoids

actinoids

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).