



# Cambridge O Level

CANDIDATE  
NAME
CENTRE  
NUMBER

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|

CANDIDATE  
NUMBER

|  |  |  |  |
|--|--|--|--|
|  |  |  |  |
|--|--|--|--|

**ADDITIONAL MATHEMATICS**
**4037/12**

Paper 1

**October/November 2024**
**2 hours**

You must answer on the question paper.

No additional materials are needed.

## INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

## INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].

 This document has **16** pages. Any blank pages are indicated.



## Mathematical Formulae

### 1. ALGEBRA

#### Quadratic Equation

For the equation  $ax^2 + bx + c = 0$ ,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

#### Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n$$

where  $n$  is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!}$

*Arithmetic series*  $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

*Geometric series*  $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

### 2. TRIGONOMETRY

#### Identities

$$\begin{aligned}\sin^2 A + \cos^2 A &= 1 \\ \sec^2 A &= 1 + \tan^2 A \\ \operatorname{cosec}^2 A &= 1 + \cot^2 A\end{aligned}$$

#### Formulae for $\triangle ABC$

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

$$\Delta = \frac{1}{2}bc \sin A$$





- 1 The curve  $y = a \cos bx + c$ , where  $a$ ,  $b$  and  $c$  are integers, passes through the points  $\left(-\frac{\pi}{6}, -2\right)$  and  $\left(\frac{\pi}{9}, \frac{1}{2}\right)$ . The curve has a period of  $\frac{2\pi}{3}$ .
- (a) Find the values of  $a$ ,  $b$  and  $c$ . [4]

- (b) Find the least value of  $y$  on the curve for  $0 \leq x \leq \frac{\pi}{2}$ , and state the value of  $x$  at which this occurs. [3]






2 It is given that  $y = f(x)$ , where  $f(x) = (2x - 5)(x - 1)^2$ .

(a) Find the coordinates of the stationary points on the curve  $y = f(x)$ .

[4]

(b) On the axes, sketch the graph of  $y = f(x)$ , stating the intercepts with the axes.

[3]




(c) Hence find the values of  $k$  for which  $f(x) = k$  has exactly one solution.

[2]



- 3 In this question, all lengths are in centimetres and all angles are in radians.



The diagram shows a circle with centre  $O$  and radius 12, and a circle with centre  $C$  and radius 5. The circles intersect at the points  $A$  and  $B$ , such that  $OA$  and  $OB$  are tangents to the circle with centre  $C$ .

- (a) Show that the obtuse angle  $ACB$  is 2.35 radians, correct to 2 decimal places. [2]

- (b) Find the perimeter of the shaded region. [2]

- (c) Find the area of the shaded region. [3]






4 The function  $f$  is such that  $f(x) = 4 \ln(3x - 2)$ , for  $x > a$ , where  $a$  is as small as possible.

(a) (i) Write down the value of  $a$ . [1]

(ii) Write down the range of  $f$ . [1]

(iii) Find  $f^{-1}(x)$ , stating its domain and range. [4]

(iv) On the axes sketch the graphs of  $y = f(x)$  and  $y = f^{-1}(x)$ , stating the intercepts with the axes. [4]





(b) Given that  $g(x) = (2x + 1)^{\frac{1}{2}} + 4$ , for  $x > 0$ , solve the equation  $gg(x) = 9$ .

[3]





5 (a) Show that  $\frac{1 + \cot^2 \theta}{\cot^2 \theta} = \sec^2 \theta$ .

[1]

(b) Write down the derivative of  $\tan \theta$  with respect to  $\theta$ .

[1]

(c) Using **part (a)** and **part (b)**, find the exact value of  $\int_0^{\frac{\pi}{3}} \left( \frac{1 + \cot^2 \theta}{\cot^2 \theta} - \sin \theta \right) d\theta$ .

[4]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





- 6 (a) Find, in descending powers of  $x$ , the first 3 terms in the expansion of  $\left(x + \frac{2}{x^2}\right)^{10}$ . Simplify each term as far as possible. [3]

- (b) Find the term independent of  $x$  in the expansion of  $\left(4x^2 + \frac{1}{2x^2}\right)^8$ . [2]





- 7 It is given that  $y = \frac{\ln(3x^2 - 1)}{x + 2}$ , for  $x > \frac{1}{\sqrt{3}}$ . When  $x = 1$ ,  $y$  is increasing at the rate of  $h$  units per second. Find, in terms of  $h$ , the corresponding rate of change in  $x$ , giving your answer in exact form.

[6]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN






- 8 The tangent to the curve  $y = e^x(2x+5)^{\frac{1}{2}}$  at the point where  $x = 2$  meets the  $x$ -axis at the point  $X$  and the  $y$ -axis at the point  $Y$ . Find the coordinates of the mid-point of  $XY$ , giving your answer in exact form. [8]





9



The diagram shows part of the curve  $y = \frac{4}{2x+1}$  and the straight line  $2y = 6x + 1$ . Find the area of the shaded region, giving your answer in the form  $\ln a + b$ , where  $a$  is an integer and  $b$  is a rational number. [8]





Continuation of working space for question 9.





- 10 (a) The first 3 terms of an arithmetic progression are  $2 \tan 2x$ ,  $5 \tan 2x$ ,  $8 \tan 2x$ . Find the values of  $x$ , where  $-180^\circ \leq x \leq 180^\circ$ , for which the sum to 30 terms is  $455\sqrt{3}$ . [5]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





(b) The first 3 terms of a geometric progression are

$$5 \cos^2\left(\theta - \frac{\pi}{2}\right), \quad 20 \cos^4\left(\theta - \frac{\pi}{2}\right), \quad 80 \cos^6\left(\theta - \frac{\pi}{2}\right), \quad \text{where } -\frac{\pi}{6} \leq \theta \leq \frac{7\pi}{6}.$$

Find the values of  $\theta$  for which this geometric progression has a sum to infinity.

[6]





Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

