

GCE AS

Mathematics

Summer 2009

Mark Schemes

Issued: October 2009

**NORTHERN IRELAND GENERAL CERTIFICATE OF SECONDARY EDUCATION
(GCSE) AND NORTHERN IRELAND GENERAL CERTIFICATE OF EDUCATION (GCE)**

MARK SCHEMES (2009)

Foreword

Introduction

Mark Schemes are published to assist teachers and students in their preparation for examinations. Through the mark schemes teachers and students will be able to see what examiners are looking for in response to questions and exactly where the marks have been awarded. The publishing of the mark schemes may help to show that examiners are not concerned about finding out what a student does not know but rather with rewarding students for what they do know.

The Purpose of Mark Schemes

Examination papers are set and revised by teams of examiners and revisers appointed by the Council. The teams of examiners and revisers include experienced teachers who are familiar with the level and standards expected of 16 and 18-year-old students in schools and colleges. The job of the examiners is to set the questions and the mark schemes; and the job of the revisers is to review the questions and mark schemes commenting on a large range of issues about which they must be satisfied before the question papers and mark schemes are finalised.

The questions and the mark schemes are developed in association with each other so that the issues of differentiation and positive achievement can be addressed right from the start. Mark schemes therefore are regarded as a part of an integral process which begins with the setting of questions and ends with the marking of the examination.

The main purpose of the mark scheme is to provide a uniform basis for the marking process so that all the markers are following exactly the same instructions and making the same judgements in so far as this is possible. Before marking begins a standardising meeting is held where all the markers are briefed using the mark scheme and samples of the students' work in the form of scripts. Consideration is also given at this stage to any comments on the operational papers received from teachers and their organisations. During this meeting, and up to and including the end of the marking, there is provision for amendments to be made to the mark scheme. What is published represents this final form of the mark scheme.

It is important to recognise that in some cases there may well be other correct responses which are equally acceptable to those published: the mark scheme can only cover those responses which emerged in the examination. There may also be instances where certain judgements may have to be left to the experience of the examiner, for example, where there is no absolute correct response – all teachers will be familiar with making such judgements.

The Council hopes that the mark schemes will be viewed and used in a constructive way as a further support to the teaching and learning processes.

CONTENTS

	Page
Unit C1	1
Unit C2	5
Unit F1	11
Unit M1	19
Unit S1	27



Rewarding Learning

ADVANCED SUBSIDIARY (AS)
General Certificate of Education
Summer 2009

Mathematics

Assessment Unit C1

assessing

Module C1: AS Core Mathematics 1

[AMC11]

FRIDAY 5 JUNE, AFTERNOON

MARK SCHEME

		AVAILABLE MARKS
1 (i)	$m_{AB} = \frac{3 - (-3)}{-1 - 2} = \frac{6}{-3} = -2$	M1W1
	$y - 3 = -2(x + 1)$ $y + 2x = 1$	M1W1
(ii)	$4y - 2x = 6$	
	$y + 2x = 1$	M1
	$5y = 7$	W1
	$y = 1.4$ $x = -0.2$	W1
2	$4x - y + 2z = 13$ $2x + y + 2z = 6$ $\frac{2x - 2y = 7}{2x - 2y = 7}$	M1W1
	$4x - y + 2z = 13$ $4x - 4y - 2z = 6$ $\frac{8x - 5y = 19}{8x - 5y = 19}$	MW1
3 (a)	$\frac{dy}{dx} = 6x^2 - 4$	M1W1
	$x = 1 \rightarrow m_t = 2$	MW1
	$x = 1 \rightarrow y = 3$	MW1
	$y - 3 = 2(x - 1)$ $y = 2x + 1$	M1W1
	(b) (i) $\frac{dy}{dx} = 8x - x^{-2}$	MW2
(ii)	$8x = \frac{1}{x^2}$	M1
	$x^3 = \frac{1}{8}$	
	$x = \frac{1}{2}$	W1
	$y = 4(\frac{1}{4}) + 2 = 3$	M1W1
	$\frac{d^2y}{dx^2} = 8 + 2x^{-3}$	M1
	$x = \frac{1}{2} \Rightarrow \frac{d^2y}{dx^2}$ is positive	
	$(\frac{1}{2}, 3)$ is a min tp	W1

7

7

14

			AVAILABLE MARKS
4	(i) $f(l) = 9 - 9 - 1 + 1 = 0 \quad \therefore (x - 1)$ is a factor	M1W1	
	(ii) $\begin{array}{r} 9x^2 - 1 \\ x - 1 \overline{) 9x^3 - 9x^2 - x + 1} \\ \underline{9x^3 - 9x^2} \\ -x + 1 \\ \underline{-x + 1} \\ 0 \end{array}$	M1W1	
	$(x - 1)(9x^2 - 1) = (x - 1)(3x - 1)(3x + 1)$	MW1	
	(iii) $\frac{(x - 1)(3x - 1)(3x + 1)}{(3x - 1)} \div \frac{(x - 1)}{4}$	M1	
	$\frac{(x - 1)(3x - 1)(3x + 1)}{(3x - 1)} \times \frac{4}{(x - 1)}$	M1	
	$4(3x + 1)$	W2	9
5	(i) $C = 80$ initial temperature 80°C	MW1	
	(ii) $80 - 10t + \frac{1}{2}t^2 = 50$ $t^2 - 20t + 60 = 0$	M1	
	$t = \frac{20 \pm \sqrt{400 - 240}}{2}$	M1	
	$t = \frac{20 \pm \sqrt{160}}{2} = 10 \pm 2\sqrt{10}$	W1	
	$t = 10 - 2\sqrt{10}$ only	W1	
	(iii) $\frac{dC}{dt} = -10 + t$	M1W1	
	$t = 3 \Rightarrow \frac{dC}{dt} = -7$	M1W1	9
6	(i) $3xy = 66$ $y = \frac{22}{x}$	M1 W1	
	(ii) $SA = 2xy + 6x + 6y = 101$	M1MW1	
	(iii) $44 + 6x + 6\left(\frac{22}{x}\right) = 101$ $6x + \frac{132}{x} - 57 = 0$	MW1	
	$6x^2 - 57x + 132 = 0$ $2x^2 - 19x + 44 = 0$ $(2x - 11)(x - 4) = 0$	MW1	
	$x = 5.5$ or $x = 4$ $y = 4$ or $y = 5.5$	M1 W1	
	Dimensions $4 \text{ cm} \times 5.5 \text{ cm} \times 3 \text{ cm}$	W1	9

7 (a) $\frac{5(1 - \sqrt{3}) - (\sqrt{3} + 1)}{(\sqrt{3} + 1)(1 - \sqrt{3})}$

$$\frac{4 - 6\sqrt{3}}{-2} = 3\sqrt{3} - 2$$

M1 MW1

(b) $\frac{(5)^x}{(5^2)^{x-1}} = 5^{\frac{1}{2}}$

$$5^{x-2(x-1)} = 5^{\frac{1}{2}}$$

$$5^{2-x} = 5^{\frac{1}{2}}$$

$$2 - x = \frac{1}{2}$$

$$x = 1\frac{1}{2}$$

MW2


M1W1

M1W1

M1

W1

(c)



M1W1

12

8 $b^2 - 4ac < 0$

$$(3 - k)^2 - 28 < 0$$

$$k^2 - 6k - 19 < 0$$

If $k^2 - 6k - 19 = 0$

$$\text{Then } k = \frac{6 \pm \sqrt{36 + 76}}{2}$$


$$k = \frac{6 \pm \sqrt{112}}{2} = 3 \pm 2\sqrt{7}$$

M2W1

W1

M1

W1



M1

$$3 - 2\sqrt{7} < k < 3 + 2\sqrt{7}$$

W1

8

Total

75



Rewarding Learning

**ADVANCED SUBSIDIARY (AS)
General Certificate of Education
Summer 2009**

Mathematics

Assessment Unit C2

assessing

Module C2: Core Mathematics 2

[AMC21]

FRIDAY 22 MAY, MORNING

MARK SCHEME

GCE Advanced/Advanced Subsidiary (AS) Mathematics

Mark Schemes

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

1 (a) (i) $3x^3 + 2x + 4x^{-2}$

MW1

(ii) $\int 3x^3 + 2x + 4x^{-2} dx$

$$\frac{3x^4}{4} + x^2 - 4x^{-1} + c$$

MW4

(b)

x	y
0	4.0000
0.2	3.8462
0.4	3.4483
0.6	2.9412
0.8	2.4390
1	2.0000

$$h = \frac{1}{5}$$

MW1

x values, y values

M1W2

$$\int \approx \frac{1}{2}h(y_0 + 2(y_1 + y_2 + y_3 + y_4) + y_5)$$

$$= \frac{1}{2} \times 0.2(4.0000 + 2(3.8462 + 3.4483 + 2.9412 + 2.4390) + 2.0000)$$

M1

$$= 3.13(3.13494)$$

MW1

11

2 (a) (i) $u_2 = \frac{2}{3}u_1 = \frac{2}{3}(1) = \frac{2}{3}$

MW1

$$u_3 = \frac{2}{3}\left(\frac{2}{3}\right) = \frac{4}{9}$$

MW1

$$u_4 = \frac{2}{3}\left(\frac{4}{9}\right) = \frac{8}{27}$$

MW1

(ii) convergent

MW1

(b) (i) $\frac{2}{3}$

MW1

(ii) $S_\infty = \frac{a}{1-r} = \frac{1}{1-\frac{2}{3}}$

M1

$$S_\infty = 3$$

W1

7

AVAILABLE
MARKS

3 $(1 + nx)^{10}$
 $= 1 + 10nx + \frac{10 \cdot 9n^2x^2}{2 \cdot 1} \dots$
 $= 1 + 10nx + 45n^2x^2 \dots$
Hence, $45n^2 = 3(10n)$
 $45n^2 - 30n = 0$
 $15n(3n - 2) = 0$
 $n = \frac{2}{3} \quad n \neq 0$

M1W1

W1


M1W1

W1

AVAILABLE MARKS

6

4 (i)



exponential shape, (0, 1)

M1W1

shift up parallel to x -axis, (0, 2)

M1W1

(ii) $1 + 2^x = 6$

$2^x = 5$

MW1

$\log_{10} 2^x = \log_{10} 5$

M1

$x \log_{10} 2 = \log_{10} 5$

M1

$x = \frac{\log_{10} 5}{\log_{10} 2}$

W1

$x = 2.32$

8

			AVAILABLE MARKS
5	(a)	$\tan \theta + \frac{1}{\tan \theta} \equiv \frac{1}{\sin \theta \cos \theta}$	
		L.H.S. $\frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta}$	M1W1
		$= \frac{\sin^2 \theta + \cos^2 \theta}{\cos \theta \sin \theta}$	MW1M1
	$= \frac{1}{\sin \theta \cos \theta}$	W1	
	(b)	$\sin^2 x = \frac{1}{4}$	
		$\sin x = \frac{1}{2}$ or $\sin x = -\frac{1}{2}$	MW2
		$x = 30^\circ$ or $x = -30^\circ$	MW2
	(c)	$\cos 2x = 0.4$	
		$2x = 1.159, 5.124$	M1W2
		$x = 0.580, 2.56$	MW1
6	(i)	Area = $\int_0^1 (4 + x^2) dx$	M2W1
		$= \left[4x + \frac{x^3}{3} \right]_0^1$	MW1
		$= [4 + \frac{1}{3}] - [0 - 0]$	
	$= 4\frac{1}{3}$	W1	
	(ii)	y coord = $5 = 4 + x^2 \dots x = 1$ or -1	MW1
		Area of rectangle = $5 \times 1 = 5$	MW1
		Area required = $2(5 - 4\frac{1}{3}) = \frac{4}{3}$	M1MW1
			13
			9

		AVAILABLE MARKS	
7	<p>(i) Gradient AC = $\frac{1-(-1)}{2-(-1)} = \frac{2}{3}$</p> <p>(ii) Gradient BC = $-\frac{3}{2}$</p> <p>Gradient BC = $\frac{(k+5)-(-1)}{k-(-1)} = \frac{k+6}{k+1} = \frac{-3}{2}$</p> $2(k+6) = -3(k+1)$ $2k+12 = -3k-3$ $5k = -15$ $k = -3$ <p>(iii) Coordinates of B = (-3, 2)</p> <p>Centre $\frac{(2+(-3))}{2}, \frac{(1+2)}{2} = (-0.5, 1.5)$</p> <p>Radius = $\sqrt{(2-(-0.5))^2 + (1-1.5)^2} = \sqrt{(2.5)^2 + (-0.5)^2} = \sqrt{6.5}$</p> <p>Equation $(x+0.5)^2 + (y-1.5)^2 = 6.5$</p>	<p>M1W1</p> <p>MW1</p> <p>M1W1</p> <p>MW1</p> <p>MW2</p> <p>M1W1</p> <p>MW1</p>	<p>11</p>
8	<p>(i) Area = $\frac{1}{2}r^2x$</p> <p>(ii) Area = $\frac{1}{2}r^2 \sin x$</p> <p>(iii) Area of major segment = $\frac{1}{2}r^2(2\pi - x) + \frac{1}{2}r^2 \sin x$</p> <p>Area of minor segment = $\frac{1}{2}r^2x - \frac{1}{2}r^2 \sin x$</p> <p>Hence, $\frac{1}{2}r^2(2\pi - x) + \frac{1}{2}r^2 \sin x = 5(\frac{1}{2}r^2x - \frac{1}{2}r^2 \sin x)$</p> $(2\pi - x) + \sin x = 5(x - \sin x)$ $2\pi - x + \sin x = 5x - 5 \sin x$ $6 \sin x = 6x - 2\pi$ $\sin x = x - \frac{\pi}{3}$	<p>MW1</p> <p>MW1</p> <p>M1W1</p> <p>M1W1</p> <p>M1W1</p> <p>MW2</p> <p>Total</p>	<p>10</p> <p>75</p>



Rewarding Learning

**ADVANCED SUBSIDIARY (AS)
General Certificate of Education
Summer 2009**

Mathematics

Assessment Unit F1

assessing

Module FP1: Further Pure Mathematics 1

[AMF11]

TUESDAY 23 JUNE, MORNING

**MARK
SCHEME**

GCE Advanced/Advanced Subsidiary (AS) Mathematics

Mark Schemes

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

		AVAILABLE MARKS
1	(a) Rotation of 45° anticlockwise about the origin	MW2
	(b) $\begin{pmatrix} -1 & 1 \\ 6 & -2 \end{pmatrix} \begin{pmatrix} t \\ mt \end{pmatrix} = \begin{pmatrix} x \\ mx \end{pmatrix}$	M1
	Hence $\begin{matrix} -t + mt = x \\ 6t - 2mt = mx \end{matrix}$	MW1
	Dividing $\Rightarrow \frac{6 - 2m}{-1 + m} = \frac{m}{1}$	M1
	$\Rightarrow 6 - 2m = -m + m^2$	W1
	$\Rightarrow m^2 + m - 6 = 0$	
	$\Rightarrow (m + 3)(m - 2) = 0$	
	$\Rightarrow m = -3, 2$	W1
	Hence the equations of the lines are $y = -3x, y = 2x$	W1
		8
2	(i) $\mathbf{M}^2 = \begin{pmatrix} -1 & 1 \\ -2 & 4 \end{pmatrix} \begin{pmatrix} -1 & 1 \\ -2 & 4 \end{pmatrix}$	M1
	$= \begin{pmatrix} -1 & 3 \\ -6 & 14 \end{pmatrix}$	W1
	$3\mathbf{M} + 2\mathbf{I} = \begin{pmatrix} -3 & 3 \\ -6 & 12 \end{pmatrix} + \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$	M1
	$= \begin{pmatrix} -1 & 3 \\ -6 & 14 \end{pmatrix}$	W1
	Hence $\mathbf{M}^2 = 3\mathbf{M} + 2\mathbf{I}$	
	(ii) $\mathbf{M}^4 = (\mathbf{M}^2)^2 = (3\mathbf{M} + 2\mathbf{I})^2$	M1
	$= 9\mathbf{M}^2 + 12\mathbf{M} + 4\mathbf{I}$	W1
	$= 9(3\mathbf{M} + 2\mathbf{I}) + 12\mathbf{M} + 4\mathbf{I}$	M1
	$= 39\mathbf{M} + 22\mathbf{I}$	W1
		8

3 (i) $[(a, b)*(c, d)]*(e, f) = (ad + bc, bd)*(e, f)$ M1
 $= (adf + bfc + bde, bdf)$ W1

$(a, b)*[(c, d)*(e, f)] = (a, b)*(cf + de, df)$ M1
 $= (adf + bcf + bde, bdf)$ W1

Hence the associative law holds.

(ii) $(a, b)*(c, d) = (a, b)$ if (c, d) is the identity M1M1
Hence $ad + bc = a$ and $bd = b$

$\Rightarrow b(d - 1) = 0$
 $\Rightarrow d = 1$ since $b \neq 0$ W1

$\Rightarrow a + bc = a$ since $d = 1$
 $\Rightarrow c = 0$ since $b \neq 0$ W1

Also $(0, 1)*(a, b) = (a, b)$
Hence the identity element is $(0, 1)$

(iii) $(a, b)*(c, d) = (0, 1)$ if (c, d) is the inverse of (a, b) M1
Hence $ad + bc = 0$ and $bd = 1$ W1

$\Rightarrow d = \frac{1}{b}$

$\Rightarrow \frac{a}{b} + bc = 0$
 $\Rightarrow c = -\frac{a}{b^2}$ W1

Hence the inverse is $\left(-\frac{a}{b^2}, \frac{1}{b}\right)$

AVAILABLE
MARKS

11

4 (i) $\det(\mathbf{N} - \lambda\mathbf{I}) = 0$

$$\Rightarrow \begin{vmatrix} 2-\lambda & 0 & -6 \\ 3 & 1-\lambda & 4 \\ -1 & 0 & 1-\lambda \end{vmatrix} = 0$$

$$\Rightarrow (2-\lambda)\{(1-\lambda)^2 - 0\} - 6\{0 + (1-\lambda)\} = 0$$

$$\Rightarrow (1-\lambda)\{(2-\lambda)(1-\lambda) - 6\} = 0$$

$$\Rightarrow (1-\lambda)(2 - 3\lambda + \lambda^2 - 6) = 0$$

$$\Rightarrow (1-\lambda)(\lambda^2 - 3\lambda - 4) = 0$$

$$\Rightarrow (1-\lambda)(\lambda - 4)(\lambda + 1) = 0$$

$$\Rightarrow \lambda = 1, 4, -1$$

(ii) $\begin{pmatrix} 2 & 0 & -6 \\ 3 & 1 & 4 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = 4 \begin{pmatrix} x \\ y \\ z \end{pmatrix}$

$$2x - 6z = 4x$$

$$\Rightarrow 3x + y + 4z = 4y$$

$$-x + z = 4z$$

① and ③ $\Rightarrow x = -3z$

② $\Rightarrow -9z - 3y + 4z = 0$

$$\Rightarrow y = -\frac{5}{3}z$$

Hence an eigenvector is $\begin{pmatrix} -9 \\ -5 \\ 3 \end{pmatrix}$

A unit eigenvector is therefore $\frac{1}{\sqrt{115}} \begin{pmatrix} -9 \\ -5 \\ 3 \end{pmatrix}$

M1

AVAILABLE
MARKS

M1

M1

W1

W3

M1

M1

W1

W1

11

5 (a) $(a + bi)^2 = 21 - 20i$
 $\Rightarrow a^2 + 2abi - b^2 = 21 - 20i$
Hence $a^2 - b^2 = 21$

and $2ab = -20$
 $\Rightarrow b = -\frac{10}{a}$

M1

M1W1

MW1

Substituting into equation ① gives

$a^2 - \frac{100}{a^2} = 21$

M1

$\Rightarrow a^4 - 21a^2 - 100 = 0$

W1

$\Rightarrow (a^2 - 25)(a^2 + 4) = 0$

$\Rightarrow a = \pm 5$

W1

$\Rightarrow b = \frac{-10}{\pm 5} = \mp 2$

W1


(b) (i) 

Circle centre $(\sqrt{3}, 1)$ and radius $\sqrt{2}$

MW2



MW1

AVAILABLE
MARKS



$$\arg z \leq \alpha + \theta$$

M1



$$\tan \theta = \frac{1}{\sqrt{3}}$$

$$x^2 = 3 + 1 = 4$$

$$\Rightarrow \theta = \frac{\pi}{6}$$

MW1

$$\Rightarrow x = 2$$

MW1

$$\sin \alpha = \frac{\sqrt{2}}{2}$$

$$\Rightarrow \alpha = \frac{\pi}{4}$$

MW1

$$\text{Hence } \arg z \leq \frac{\pi}{4} + \frac{\pi}{6}$$

$$\Rightarrow \arg z \leq \frac{5\pi}{12}$$

W1

16

6 (i)	C_1 has centre $(-1, 7)$	MW1	AVAILABLE MARKS
	Gradient of radius $= \frac{7-6}{-1-2} = \frac{1}{-3}$	M1W1	
	Gradient of tangent $= 3$	MW1	
	Equation of tangent is $y - 6 = 3(x - 2)$	M1	
	which gives $y = 3x$	W1	
(ii)	Let equation of tangent be $y = mx$		
	Then $x^2 + m^2x^2 + 2x - 14mx + 40 = 0$	M1	
	$\Rightarrow x^2(1 + m^2) + x(2 - 14m) + 40 = 0$	W1	
	If line is a tangent then $b^2 - 4ac = 0$		
	$\Rightarrow (2 - 14m)^2 = 4(40)(1 + m^2)$	M1	
	$\Rightarrow 4 - 56m + 196m^2 = 160 + 160m^2$	W1	
	$\Rightarrow 36m^2 - 56m - 156 = 0$		
	$\Rightarrow 9m^2 - 14m - 39 = 0$	W1	
	$\Rightarrow (m - 3)(9m + 13) = 0$		
	$m = 3, -\frac{13}{9}$	W1	
	Therefore the other tangent has equation $y = -\frac{13}{9}x$	W1	
(iii)	$x^2 + y^2 + 2x - 14y + 40 = 0$		
	$x^2 + y^2 - 10x - 8y + 16 = 0$		
	Subtract to give $12x - 6y + 24 = 0$	M1	
	$\Rightarrow y = 2x + 4$	W1	
	Substitute into equation ① to give		
	$x^2 + (2x + 4)^2 + 2x - 14(2x + 4) + 40 = 0$	M1	
	$\Rightarrow x^2 + 4x^2 + 16x + 16 + 2x - 28x - 56 + 40 = 0$		
	$\Rightarrow 5x^2 - 10x = 0$	W1	
	$\Rightarrow 5x(x - 2) = 0$		
	$\Rightarrow x = 0, 2$	W2	
	$\Rightarrow y = 4, 8$	W2	
	Therefore the points of intersection are $(0, 4)$ and $(2, 8)$		

Total

21

75



Rewarding Learning

**ADVANCED SUBSIDIARY (AS)
General Certificate of Education
Summer 2009**

Mathematics

Assessment Unit M1

assessing

Module M1: Mechanics 1

[AMM11]

FRIDAY 15 MAY, MORNING

MARK SCHEME

GCE Advanced/Advanced Subsidiary (AS) Mathematics

Mark Schemes

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:


It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).


1 $\rightarrow 8 - 15 \cos 50^\circ = -1.642 \text{ N}$
 $\uparrow 15 \sin 50^\circ - 5 = 6.491 \text{ N}$



$$R = \sqrt{6.491^2 + 1.642^2} = 6.70 \text{ N}$$

$$\theta = \tan^{-1} \frac{6.491}{1.642} = 75.8^\circ$$

2 (i)



(ii) $M(P) \ 80g \times 3 = Y \times 5$

$$48gN = Y$$

$$X + Y = 80g$$

$$\uparrow \quad X = 32gN$$

(iii)



Reaction at P = 0 when about to tilt

$$M(Q) \ 3Mg = 80g \times 2$$

$$M = \frac{160}{3} \text{ kg}$$

M1
M1W1
MW1

M1W1
M1W1

MW2

M2W1
W1
M1
W1

M1
M1W1
W1


AVAILABLE
MARKS

8

12

		AVAILABLE MARKS		
3	(i)	$u = 14 \text{ m s}^{-1}$		
		$a = -9.8 \text{ m s}^{-2}$		
		$v = 0$	MW1	
		$s = h$		
		By $v^2 = u^2 + 2as$	M1W1	
		$0 = 14^2 + 2(-9.8)h$		
		$h = 10 \text{ m}$	W1	
		(ii)	$u = 14$	
		$a = 9.8$		
		$s = 8.4$		
	$t = ?$			
	By $s = ut + \frac{1}{2}at^2$			
	$8.4 = 14t + \frac{1}{2}(-9.8)t^2$	M1W1		
	$8.4 = 14t - 4.9t^2$			
	$7t^2 - 20t + 12 = 0$			
	$(7t - 6)(t - 2) = 0$			
	$t = \frac{6}{7}$ or $t = 2$	W1		
	So above 8.4 m for $2 - \frac{6}{7} = \frac{8}{7} \text{ s}$	M1W1		
4	(i)		MW2	
	(ii)	Using $F = ma$		
		car $2400 - T - 300 = 800 \times 2$	M1W1	
		trailer $T - R = 200 \times 2$	M1W1	
		system $2400 - 300 - R = 2000$	M1	
		$R = 100 \text{ N}$	W1	
		$T = 400 + R$	W1	
		$T = 500 \text{ N}$		
			9	

5



(i) By conservation of momentum

$$0.4u + 0.6 \times 0 = 0.4 \times 0 + 0.6 \times v_2$$

M2W1

$$\frac{0.4u}{0.6} = v_2$$

$$v_2 = \frac{2u}{3} \text{ m s}^{-1}$$

W1

(ii) Using $I =$ increase in momentum of A

$$I = 0.4 \times v_1 - 0.4u$$

M1

$$I = -0.4u \text{ N s}$$

W1W1

$$\text{or } I = 0.6v_2 - 0.6 \times 0$$

M1

$$= 0.6 \times \frac{2u}{3}$$

W1


$$I = 0.4u \text{ N s}$$

so impulse on A is $-I = -0.4u \text{ N s}$

W1

7

6 (i)



MW2

(ii) Using $F = ma$

$$\uparrow R - 10g \cos 30^\circ = 0$$

$$R = 10g \cos 30^\circ$$

MW1

$$F = \mu R$$

$$= 0.3 \times 10g \cos 30^\circ$$

M1

$$= 25.46$$

$$\uparrow 100 - \mu R - 10g \sin 30^\circ = 10a$$

M1

$$100 - 0.3 \times 10g \cos 30^\circ - 10g \sin 30^\circ = 10a$$

$$100 - 25.46 - 49 = 10a$$

W2

$$25.54 = 10a$$

$$2.55 \text{ m s}^{-2} = a$$

W1

8

			AVAILABLE MARKS
7 (i)	$v = t^2 - 5t + 6$ $v = 0 \Rightarrow t^2 - 5t + 6 = 0$ $(t - 2)(t - 3) = 0$ $t = 2s \text{ or } 3s$	M1 W2	
(ii)	$a = \frac{dv}{dt} = (2t - 5) \text{ m s}^{-2}$	M1W1	
(iii)	$s = \int v \, dt$ $= \int t^2 - 5t + 6 \, dt$ $s = \frac{t^3}{3} - \frac{5t^2}{2} + 6t + c$ $t = 6, s = 0 \text{ so}$ $0 = \frac{6^3}{3} - 5 \times \frac{6^2}{2} = 6 \times 6 + c$ $0 = 72 - 90 + 36 + c$ $c = -18$ $s = \frac{t^3}{3} - \frac{5t^2}{2} + 6t - 18$	M1 W1 M1W1	
(iv)	$t = 0 \quad s = -18$ $t = 2 \quad s = \frac{2^3}{3} - \frac{5(2)^2}{2} + 6(2) - 18 = -13\frac{1}{3}$ $t = 3 \quad s = \frac{(3)^3}{3} - \frac{5(3)^2}{2} + 6(3) - 18 = -13\frac{1}{2}$ Distance travelled $4\frac{2}{3} + \frac{1}{6} = 4\frac{5}{6} \text{ m}$	MW2 M1W1	
			13

8 Assume police car catches motorcyclist when $t = T$.

Motorcyclist $u = 30 \text{ m s}^{-1}$

$$a = 0.2 \text{ m s}^{-2}$$

$$s = d$$

$$t = T$$

Using $s = ut + \frac{1}{2}at^2$

$$d = 30T + \frac{1}{2}(0.2)T^2$$

$$d = 30T + 0.1T^2$$

Police car $u = 0$

$$a = 1 \text{ m s}^{-2}$$

$$t = T - 5$$

$$s = d$$

Using $s = ut + \frac{1}{2}at^2$

$$d = 0(T - 5) + \frac{1}{2}(1)(T - 5)^2$$

Hence $30T + 0.1T^2 = \frac{1}{2}(T^2 - 10T + 25)$

$$30T + 0.1T^2 = 0.5T^2 - 5T + 12.5$$

$$0.4T^2 - 35T + 12.5 = 0$$

$$4T^2 - 350T + 125 = 0$$

$$T = 87.1 \text{ or } 0.359 \text{ (not valid)}$$

Hence $T = 87.1 \text{ s}$

MW1

M1

W1

MW1

MW1

W1


M1

W1

W1

AVAILABLE
MARKS

Alternative solution



Draw level when $t = T =$

Area under graphs equal

$$\frac{1}{2}(T - 5)^2 = \frac{1}{2}(30 + 0.2T + 30) T$$

$$T^2 - 10T + 25 = 60T + 0.2T^2$$

$$0.8T^2 - 70T + 25 = 0$$

$$T = 87.1 \text{ or } 2.9 \text{ (not valid)}$$

Hence $T = 87.1\text{s}$

MW1M3W2

W1

W1

W1

9

Total

75

AVAILABLE
MARKS



Rewarding Learning

**ADVANCED SUBSIDIARY (AS)
General Certificate of Education
Summer 2009**

Mathematics

Assessment Unit S1

assessing

Module S1: Statistics 1

[AMS11]

MONDAY 1 JUNE, MORNING

MARK SCHEME

GCE Advanced/Advanced Subsidiary (AS) Mathematics

Mark Schemes

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:

It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

		AVAILABLE MARKS	
3	(i) $8k = 1$ $k = 0.125$	M1 W1	
	(ii) Symmetrically mid-way between 5 and 12	M1	
	(iii) $E(X^2) = 0.125(5^2 + 6^2 + 7^2 + 8^2 + 9^2 + 10^2 + 11^2 + 12^2)$ $= 77.5$ $\text{Var}(X) = E(X^2) - [E(X)]^2$ $= 77.5 - 8.5^2$ $= 5.25$	M1 W1 M1 W1	
	(iv) $E(Y) = 2 \times 8.5 - 5 = 12$ $\text{Var}(Y) = 2^2 \times 5.25 = 21$	MW1 M1W1	
4	X is r.v. "No. of goals scored during a match"		
(i)	$\lambda = \frac{21}{35} = 0.6 \quad X \sim \text{Po}(0.6)$ $P(\text{scores}) = P(X \geq 1) = 1 - P(X = 0)$ $P(X = 0) = \frac{e^{-0.6} 0.6^0}{0!}$ $= 0.5488 \dots$ $P(\text{scores}) = 0.451188 \dots = 0.451 \text{ (3 s.f.)}$	MW1 M1 M1 W1	
	(ii) $P(X = 1) = \frac{e^{-0.6} \times 0.6^1}{1!} = 0.6e^{-0.6} = 0.32928$ $P(X = 2) = \frac{e^{-0.6} \times 0.6^2}{2!} = 0.18e^{-0.6} = 0.09878$ $P(X = 1 \text{ or } 2) = 0.428 \text{ (3 s.f.)}$	MW1 MW1 W1	
	(iii) $P(X \geq 3) = 1 - [P(X = 0) + P(X = 1) + P(X = 2)]$ $= 1 - (0.5488 + 0.32928 + 0.09878)$ $= 0.02311 \dots$ $E(\text{Bonus}) = 1000 \times P(X = 1 \text{ or } 2) + 5000 \times P(X \geq 3)$ $= 1000 \times 0.42807 + 5000 \times 0.02311 \dots$ $= \text{£}543.62$	M1 W1 W1 M1 W1 W1	
			10
			12

5 (i) Let X be r.v. mass of student, $X \sim N(\mu, 12^2)$

$$P(X < 111.74) = 0.95 \Rightarrow P\left(Z < \frac{111.74 - \mu}{12}\right) = 0.95$$

M1

$$\Rightarrow \frac{111.74 - \mu}{12} = \Phi^{-1}(0.95)$$

M1

$$\frac{111.74 - \mu}{12} = 1.645$$

W1

$$\Rightarrow \mu = 111.74 - 12 \times 1.645 = 92$$

W1

(ii) $P(X < 89) = P\left(Z < \frac{89 - 92}{12}\right) = P(Z < -0.25)$

W1

$$= 1 - \Phi(0.25)$$

M1

$$= 1 - 0.5987$$

W1

$$= 0.4013 \quad 0.401 \text{ (3 s.f.)}$$

W1

(iii) $P(89 < X < 98) = P\left(\frac{89 - 92}{12} < Z < \frac{98 - 92}{12}\right)$

$$= P(-0.25 < Z < 0.5)$$

W1

$$= \Phi(0.5) - \Phi(0.25)$$

M1

$$= 0.6915 - 0.4013 = 0.2902 = 0.290 \text{ (3 s.f.)}$$

W1, W1

(iv) $P(X < W) = 0.8$

$$\Rightarrow P\left(Z < \frac{W - 92}{12}\right) = 0.8$$

M1

$$\Rightarrow \Phi\left(\frac{W - 92}{12}\right) = 0.8$$

$$\frac{W - 92}{12} = \Phi^{-1}(0.8)$$

MW1

$$= 0.842$$

W1

$$W = 12 \times 0.842 + 92 = 102.104$$

W1

$$102 \text{ (3 s.f.)}$$

16

		AVAILABLE MARKS	
6	(i) $f(2) = 2k$	MW1	12
	(ii) Area = 1	M1	
	$\frac{1}{2} \times 3 \times 2k = 1$	MW1	
	$\Rightarrow k = \frac{1}{3}$	W1	
	(iii) $P(1 \leq X \leq 3) = 1 - P(X \leq 1)$	M1	
	$= 1 - \frac{1}{2} \times 1 \times \frac{1}{3}$	MW1	
	$= 1 - \frac{1}{6}$		
	$= \frac{5}{6}$	W1	
	(iv) Let $m =$ median of $X [m < 2]$	M1	
	$f(m) = \frac{1}{3}m$	W1	
$P(X < m) = \frac{1}{2}$	M1	12	
$\therefore \frac{1}{2} \times m \times \frac{1}{3}m = \frac{1}{2}$	W1		
$m^2 = 3$			
$m = \sqrt{3}$	W1		
7	(i) $P(F \cap C) = P(F) \times P(C F)$		M1
	$0.082 = P(F) \times 0.2$		
	$P(F) = \frac{0.082}{0.2} = 0.41$		W1
	(ii) $P(F \cap C) = P(C) \times P(F C)$		M1
	$0.082 = P(C) \times 0.25$		
	$P(C) = \frac{0.082}{0.25} = 0.328$		W1
	(iii) $P(F \cup C) = P(F) + P(C) - P(F \cap C)$	M1	
	$= 0.41 + 0.328 - 0.082 = 0.656$	W1	
	$P(\bar{F} \cap \bar{C}) = 1 - 0.656 = 0.344$	MW1	
	(iv) $P(F \bar{C}) = \frac{P(F \cap \bar{C})}{P(\bar{C})}$	M1	
$P(\bar{C}) = 1 - P(C) = 1 - 0.328 = 0.672$	MW1	12	
$P(F \cap \bar{C}) = P(F) - P(F \cap C) = 0.41 - 0.082 = 0.328$	MW1		
$P(F \bar{C}) = \frac{0.328}{0.672} = 0.488$ (3 s.f.)	W2		
Total			75

