



Rewarding Learning

ADVANCED
General Certificate of Education
2010

Mathematics

Assessment Unit F3

assessing

Module FP3: Further Pure Mathematics 3

[AMF31]



THURSDAY 27 MAY, MORNING

TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number on the Answer Booklet provided.

Answer **all seven** questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

You are permitted to use a graphic or a scientific calculator in this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 75

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

A copy of the **Mathematical Formulae and Tables booklet** is provided.

Throughout the paper the logarithmic notation used is $\ln z$ where it is noted that

$\ln z \equiv \log_e z$

Answer all seven questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

1 Using integration by parts, find

$$\int e^x \cos x \, dx \quad [6]$$

2 The points A (2, 0, -1), B (4, 3, 1) and the origin determine a plane Π_1

(i) Verify that an equation of the plane is

$$x - 2y + 2z = 0 \quad [2]$$

The plane Π_2 has an equation $\mathbf{r} \cdot \mathbf{n} = d$ where

$$\mathbf{n} = 3\mathbf{i} - 2\mathbf{j} - \mathbf{k}$$

and d is a constant.

(ii) If the point B lies on the plane Π_2 , find the value of d . [2]

The planes Π_1 and Π_2 intersect in the line L.

(iii) Find, in Cartesian form, an equation for the line L. [6]

3 For each non-negative integer n , let

$$I_n = \int_0^1 x^n e^{-x} dx$$

(i) Show that for $n \geq 1$

$$I_n = nI_{n-1} - e^{-1} \quad [5]$$

(ii) Hence find the exact value of

$$\int_0^1 x^4 e^{-x} dx \quad [5]$$

4 By using the substitution $u = e^x$ find

$$\int \frac{dx}{5 \cosh x + 4 \sinh x} \quad [8]$$

5 (i) Show that

$$\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}} \quad [4]$$

The tangent at the point where $x = \frac{1}{8}$ on the curve $y = \cos^{-1} 4x$, cuts the y -axis at the point P.

(ii) Show that

$$OP = \frac{\pi + \sqrt{3}}{3}$$

where O is the origin.

[6]

- 6 (i) Using the exponential definitions of $\sinh x$ and $\cosh x$ show that for $|x| < 1$

$$\tanh^{-1} x = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right) \quad [5]$$

- (ii) Sketch the graph of $y = \tanh^{-1} x$ clearly labelling any asymptotes. [2]

- (iii) Find $\frac{d}{dx} (\tanh^{-1} x)$ [4]

- (iv) Solve the equation

$$x = \tanh \left(\ln \sqrt{6x} \right)$$

where $0 < x < 1$ [5]

- 7 Relative to an origin O the points A and B have position vectors \mathbf{a} and \mathbf{b} respectively where

$$\mathbf{a} = -\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

and $\mathbf{b} = \mathbf{i} + 2\mathbf{j} - \mathbf{k}$

- (i) Find the area of the triangle OAB. [5]

The point C is the intersection of the two lines

$$(\mathbf{r} - \mathbf{a}) \times \mathbf{m} = \mathbf{0}$$

and $(\mathbf{r} - \mathbf{b}) \times \mathbf{n} = \mathbf{0}$

where $\mathbf{m} = \mathbf{i} + \mathbf{k}$ and $\mathbf{n} = -\mathbf{i} - \mathbf{j} + 4\mathbf{k}$

- (ii) Find \mathbf{c} , the position vector of the point C. [7]

- (iii) Show that the volume of the tetrahedron OABC is $\frac{4}{3}$ [3]

$$\left[\text{The volume of a tetrahedron is } \frac{1}{6} |\mathbf{c} \cdot (\mathbf{a} \times \mathbf{b})| \right]$$