



ADVANCED
General Certificate of Education
2011

Mathematics
Assessment Unit M2
assessing
Module M2: Mechanics 2
[AMM21]



THURSDAY 16 JUNE, AFTERNOON

* Please note amendment to Question 4(i) on page 3.

TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number on the Answer Booklet provided.
Answer **all seven** questions.
Show clearly the full development of your answers.
Answers should be given to three significant figures unless otherwise stated.
You are permitted to use a graphic or a scientific calculator in this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 75
Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.
Answers should include diagrams where appropriate and marks may be awarded for them.
Take $g = 9.8 \text{ m s}^{-2}$, unless specified otherwise.

A copy of the **Mathematical Formulae and Tables booklet** is provided.

Throughout the paper the logarithmic notation used is $\ln z$ where it is noted that $\ln z \equiv \log_e z$

Answer all seven questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

1 Two forces $(2\mathbf{i} - 11\mathbf{j})$ N and $(6\mathbf{i} + 7\mathbf{j})$ N act on a body of mass 4 kg.

(i) Find the acceleration of the body. [3]

The body starts from rest.

(ii) Find the **speed** of the body after 4 s. [5]

2 A smooth bend in a track is banked at 20° to the horizontal. A car of mass 1000 kg travels round the bend at a speed of 15 m s^{-1} on an arc of a horizontal circle, of radius r metres, as shown in **Fig. 1** below.




Fig. 1

Model the car as a particle.

(i) Draw a diagram showing all the external forces acting on the car. [2]

(ii) Find the normal reaction between the car and the road. [3]

(iii) Find r . [5]

- 3 A pair of scale pans, A and B, each of mass m kg, are attached to the ends of a light inextensible string which passes over a smooth fixed pulley. They are held at the same level as shown in **Fig. 2** below.




Fig. 2

A mass of $2m$ kg is now placed on A.
The system is released from rest.

- (i) Calculate, in terms of m , the kinetic energy of B when it is moving at 4 m s^{-1} [2]
- (ii) Using the principle of conservation of energy, find how far B has risen when it has a speed of 4 m s^{-1} [7]
- (iii) State one modelling assumption you have made when answering this question. [1]

* In the light of further quality assurance, you should attempt this question with part (i) revised as shown below.

- 4 A fire engine's pump raises 100 litres of water through a vertical distance of 10 m.
The water issues from the end of a hose at 25 m s^{-1}
1 litre of water has a mass of 1 kg.
- (i) If the cross-sectional area of the end of the hose is 0.01 m^2 , find the time taken for the 100 litres to issue from the end of the hose. [2]
- (ii) Find the power developed by the fire engine's pump. [5]

- 5 A lorry of mass 2000 kg is travelling along a straight horizontal test track. When its velocity is 30 m s^{-1} its engine is switched off. During its subsequent motion a horizontal resistance of magnitude $100v^2 \text{ N}$ opposes its motion, where $v \text{ m s}^{-1}$ is the lorry's speed at any time t seconds. Model the lorry as a particle.

- (i) Show that the lorry's equation of motion can be modelled by

$$\frac{dv}{dt} = -\frac{v^2}{20} \quad [3]$$

- (ii) Find the value of t when $v = 0.5$ [8]

6 A ball is projected at an angle θ above the horizontal with an initial velocity of $u \text{ m s}^{-1}$

(i) Prove that the greatest height, h metres, of the ball above the horizontal is given by

$$h = \frac{u^2 \sin^2 \theta}{2g}$$

[4]

A ball bounces off a roof with an initial velocity of 3 m s^{-1} at an angle of 30° above the horizontal.

The roof is 4 m vertically above horizontal ground as illustrated in **Fig. 3** below.




Fig. 3

(ii) Find the greatest height above the ground reached by the ball.

[4]

(iii) Show that the ball will reach the ground after 1.07 s

[4]

(iv) Find the horizontal distance travelled by the ball before it reaches the ground.

[2]

7 The velocity $v \text{ m s}^{-1}$ of a particle R at any time t seconds is given by

$$v = (t^2 - 4t)\mathbf{i} + \left(\frac{t^3}{3} - t^2\right)\mathbf{j} + \mathbf{k}$$

(i) Find an expression for the acceleration of R at any time t . [3]

(ii) Find t when the acceleration is zero. [4]

R passes through a fixed point O when $t = 0$

(iii) Find an expression for the displacement of R from O at any time t . [4]

(iv) Find the **distance** of R from O when $t = 3$ [4]

THIS IS THE END OF THE QUESTION PAPER

Permission to reproduce all copyright material has been applied for.
In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA
will be happy to rectify any omissions of acknowledgement in future if notified.