



Rewarding Learning

ADVANCED
General Certificate of Education
2014

Mathematics

Assessment Unit M4

assessing

Module M4: Mechanics 4

[AMM41]

MONDAY 23 JUNE, MORNING

MARK
SCHEME

GCE ADVANCED/ADVANCED SUBSIDIARY (AS) MATHEMATICS

Introduction

The mark scheme normally provides the most popular solution to each question. Other solutions given by candidates are evaluated and credit given as appropriate; these alternative methods are not usually illustrated in the published mark scheme.

The marks awarded for each question are shown in the right-hand column and they are prefixed by the letters **M**, **W** and **MW** as appropriate. The key to the mark scheme is given below:

M indicates marks for correct method.

W indicates marks for working.

MW indicates marks for combined method and working.

The solution to a question gains marks for correct method and marks for an accurate working based on this method. Where the method is not correct no marks can be given.

A later part of a question may require a candidate to use an answer obtained from an earlier part of the same question. A candidate who gets the wrong answer to the earlier part and goes on to the later part is naturally unaware that the wrong data is being used and is actually undertaking the solution of a parallel problem from the point at which the error occurred. If such a candidate continues to apply correct method, then the candidate's individual working must be followed through from the error. If no further errors are made, then the candidate is penalised only for the initial error. Solutions containing two or more working or transcription errors are treated in the same way. This process is usually referred to as "follow-through marking" and allows a candidate to gain credit for that part of a solution which follows a working or transcription error.

Positive marking:


It is our intention to reward candidates for any demonstration of relevant knowledge, skills or understanding. For this reason we adopt a policy of **following through** their answers, that is, having penalised a candidate for an error, we mark the succeeding parts of the question using the candidate's value or answers and award marks accordingly.

Some common examples of this occur in the following cases:

- (a) a numerical error in one entry in a table of values might lead to several answers being incorrect, but these might not be essentially separate errors;
- (b) readings taken from candidates' inaccurate graphs may not agree with the answers expected but might be consistent with the graphs drawn.

When the candidate misreads a question in such a way as to make the question easier only a proportion of the marks will be available (based on the professional judgement of the examining team).

1 Separating the Pythagorean Triples and so combining the forces in pairs. M1



MW1
MW1
MW1

M1
W1
W1
MW1

Alternative solution

Trig. values

MW1

$$\text{Res } \leftrightarrow 29 \times \frac{20}{29} + 24 \times \frac{3}{5} + 32 \times \frac{4}{5} = 60$$

M1 W1

$$\text{Res } \uparrow 29 \times \frac{21}{29} - 21 - 24 \times \frac{4}{5} + 32 \times \frac{3}{5} = 0$$

MW1

$$\begin{aligned} \text{Moments } \curvearrowright \text{ B } \quad 60d &= 21 \times 2.0 - 32 \times 1.2 \\ &= 3.6 \\ d &= 0.06 \end{aligned}$$

M1 MW1

W1

Resultant 60 N, // BD $d = 0.06$

MW1

8

2 (i) Perp from D to P is $\sqrt{3} l$ MW1

Moments \curvearrowright D

$$\begin{aligned} P \times \sqrt{3} l &= 1000 \times 2\sqrt{3} l \\ P &= 2000 \text{ N} \end{aligned}$$

M1

W1

(ii) BA must have a \uparrow component so tension M1
BC must have a \leftarrow component as BA now has a \rightarrow component so a thrust M1

(iii) at B resolving

$$\uparrow T_1 \sin 30^\circ = 1000 \text{ N}$$

M1

$$T_1 = 2000 \text{ N}$$

W1

$$\leftrightarrow T_1 \cos 30^\circ = T_2$$

$$T_2 = 2000 \times \frac{\sqrt{3}}{2}$$

$$= 1000\sqrt{3} \text{ N}$$

MW1

(iv) at C Res \downarrow $T_3 = 0$
 at A Res $\perp_{R} BA$ $T_4 \cos 30^\circ = T_3 \cos 30^\circ = 0$
 or $T_4 \cos 30^\circ + T_3 \cos 30^\circ = 0 \quad \therefore T_4 = 0$


MW1

MW1

AVAILABLE
MARKS

10

3



(i) at P M1
 $\frac{1}{2} mv^2 + mgr(1 + \cos \theta) = 0 + 2 mgr$ MW2
 $KE = \frac{1}{2} mv^2 = mgr(1 - \cos \theta)$ W1

(ii) at P M1
 Res \swarrow $\frac{mv^2}{r} = mg \cos \theta - R$ M1 W1
 $R = mg \cos \theta - 2mg(1 - \cos \theta)$ MW1
 $= mg(3 \cos \theta - 2)$ W1

(iii) $mg(3 \cos \theta_c - 2) = 0$ M1
 $\cos \theta_c = \frac{2}{3}$
 $\theta_c = \cos^{-1} \frac{2}{3}$
 $= 48.2^\circ$ W1

11

4 (i) $\frac{GMm}{r^2} = mg \quad \therefore G = \frac{gr^2}{M}$ MW1

$[G] = [LT^{-2}][L]^2[M]^{-1}$ M1

$= [M]^{-1}[L]^3[T]^{-2}$ W1

(ii) $[T] = [M]^x[L]^y[M^{-1}L^3T^{-2}]^z$ M1


$= [M]^{x-z}[L]^{y+3z}[T]^{-2z}$ W1

$\therefore [M] \quad x - z = 0$ M1

$[L] \quad y + 3z = 0$

$[T] \quad -2z = 1$

$\therefore z = -\frac{1}{2} = x \quad y = -3z = \frac{3}{2}$ W2

(iii)  $\frac{GMm_1}{d^2} = m_1d\omega^2$ M1 W1

$\omega^2 = \frac{GM}{d^3}$ W1

$T = \frac{2\pi}{\omega}$


$= 2\pi\sqrt{\frac{d^3}{GM}}$ M1

$= 2\pi M^{-\frac{1}{2}}d^{\frac{3}{2}}G^{-\frac{1}{2}}$

$\therefore k = 2\pi$ W1

AVAILABLE MARKS

13



Cons mom. $\cancel{mv_2} + k\cancel{mv_1} = k\cancel{mv_1} + \cancel{mv_2}$
 Newton's law $v_2 - v_1 = (-e)(-u) = eu$
 $(1+k)v_1 = (k-e)u$

M1 W1
 M1 W1

$v_1 = \frac{(k-e)u}{1+k}$
 $v_2 = v_1 + eu$

W1

$= \frac{(k-e)u}{1+k} + \frac{(1+k)eu}{1+k}$
 $= \frac{(1+e)ku}{1+k}$

W1

(ii) $v_3 \leftarrow = ev_2 = \frac{e(1+e)ku}{1+k}$

MW1

If $k > e$ $v_1 \rightarrow v_3 \leftarrow$
 \therefore a collision

M1

(iii) If $k < e$,
 then for another collision we need

M1

$\frac{e(1+e)ku}{1+k} > \frac{(e-k)u}{1+k}$

W1

i.e. $e^2k + ek - e + k > 0$
 $ke^2 - (1-k)e + k > 0$

W1

11

- 6 (i) $y_1 - y_2$ is the strip length and is needed to find the mass. M1
 $\frac{1}{2}(y_1 + y_2)$ is the distance the strip's c.o.m. is from OX M1
 and needed to calculate its moment about OX


(ii) $I_2 = \int_0^2 px(2x - x^2) dx = \int_0^2 p(2x^2 - x^3) dx$ MW1

$= p \left[\frac{2x^3}{3} - \frac{x^4}{4} \right]_0^2$ MW1


$= p \left(\frac{16}{3} - \frac{16}{4} \right) = \frac{16p}{12} = \frac{4p}{3}$ MW1

$I_1 = \frac{4p}{3} \quad \therefore \quad \bar{x} = \frac{I_2}{I_1} = 1$ MW1

(iii) $\bar{y} = \frac{I_3}{I_1} = \frac{28p}{15} \times \frac{3}{4p}$
 $= 1.4$ MW1



$d = (1.4 - 1) \cos 45^\circ = 0.4 \times \frac{1}{\sqrt{2}} = 0.2\sqrt{2}$ M1W1



Moments B \curvearrowright M1

$2\sqrt{2}P = 0.2\sqrt{2} \times 250$ MW1

$P = 25 \text{ N}$ W1

7 (i) Resolving vertically

$$mg = (E + F) \sin \alpha + (N + P) \cos \alpha$$

M1 W1

Resolving horizontally

$$\frac{mv^2}{r} = (E + F) \cos \alpha - (N + P) \sin \alpha$$

M2 W1

$$E + F = \mu(N + P)$$

MW1

$$\frac{mv^2}{r} = \frac{\mu \cos \alpha - \sin \alpha}{\mu \sin \alpha + \cos \alpha}$$

MW1

$$= \frac{\mu - \tan \alpha}{\mu \tan \alpha + 1}$$

$$\therefore v^2 = rg \left(\frac{\mu - \tan \alpha}{\mu \tan \alpha + 1} \right)$$

W1

(ii)
$$v^2 = 98 \left(\frac{\frac{6}{7} - \frac{1}{4}}{\frac{6}{7} \times \frac{1}{4} + 1} \right)$$

MW1

$$v^2 = 49$$

$$v = 7$$

W1

Total

AVAILABLE
MARKS

10

75