

Mark Scheme (Results)

January 2014

IAL Chemistry (WCH04/01)

Unit 4: General Principles of Chemistry I

ALWAYS LEARNING PEARSON

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2014
Publications Code IA037633
All the material in this publication is copyright
© Pearson Education Ltd 2014

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:
 - i) ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear
 - ii) select and use a form and style of writing appropriate to purpose and to complex subject matter
 - iii) organise information clearly and coherently, using specialist vocabulary when appropriate

Using the Mark Scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit.

() means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in bold indicate that the <u>meaning</u> of the phrase or the actual word is essential to the answer.

ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

Quality of Written Communication

Ouestions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities. Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

Section A (multiple choice)

Question	Correct Answer	Reject	Mark
Number 1	C		1
<u> </u>	C		l
Question	Correct Answer	Reject	Mark
Number	_		
2 (a)	В		1
Question	Correct Answer	Reject	Mark
Number		reject	, idirk
2(b)	D		1
Ougation	Courset Anouse	Deiest	Moule
Question Number	Correct Answer	Reject	Mark
2(c)	В		1
Question	Correct Answer	Reject	Mark
Number 3	D		1
J		<u>l</u>	
Question	Correct Answer	Reject	Mark
Number	6		4
4	С		1
Question	Correct Answer	Reject	Mark
Number		_	
5(a)	D		1
Question	Correct Answer	Reject	Mark
Number	Correct / tilswei	reject	riark
5(b)	Α		1
Ougation	Courset Anouse	Deiest	Moule
Question Number	Correct Answer	Reject	Mark
6	С		1
Ougation	Courset Anouse	Deiest	Moule
Question Number	Correct Answer	Reject	Mark
7	D		1
Question	Correct Answer	Reject	Mark
Number 8	В		1
U	1 5	1	_ '
Question	Correct Answer	Reject	Mark
Number	B		
9(a)	В		1

Question Number	Correct Answer	Reject	Mark
9(b)	Α		1
	•	·	•
Question Number	Correct Answer	Reject	Mark
9(c)	С		1
	•		·
Question Number	Correct Answer	Reject	Mark
10	D		1
Question Number	Correct Answer	Reject	Mark
11	D		1
Question Number	Correct Answer	Reject	Mark
12	В		1
		•	·
Question Number	Correct Answer	Reject	Mark
13	D		1
Question Number	Correct Answer	Reject	Mark
14	С		1
			·
Question Number	Correct Answer	Reject	Mark
15	Α		1

Total for Section A = 20 marks

Section B

Question Number	Acceptable Answers	Reject	Mark
	CH CH CH C-N	CH CH CH CN	
16(a)	$CH_3CH_2CH_2C\equiv N$	CH ₃ CH ₂ CH ₂ CN	
	ALLOW		
	displayed formula	molecular formula	1

Question Number	Acceptable Answers	Reject	Mark
16(b)	I GNORE conditions and solvents, even if incorrect	incorrect formulae, including subscripts written as large numbers or superscripts eg LiAlH4/LiAlH ⁴ any charges	
	Step 1 LiAlH $_4$ I GNORE dry ether/ followed by H $_2$ O ALLOW lithium tetrahydridoaluminate((III)) lithium aluminium hydride (1) Step 2 PCI $_5$	NaBH₄ H₂/ hydrogen	
	ALLOW phosphorus(V) chloride/ phosphorus pentachloride HCl /(concentrated) hydrochloric acid PCl ₃ / phosphorus(III) chloride/ phosphorus trichloride SOCl ₂ / thionyl chloride (1)	dilute hydrochloric acid	
	Step 4 HCI/ HCI(aq)/ HCI in water or H ₂ O ALLOW any strong acid/ H ⁺ / NaOH/ sodium hydroxide followed by HCI / hydrochloric acid (1)	just 'dilute acid' just 'concentrated acid' just 'H ₂ O/ water'	
	Step 5 CH_3CH_2OH/C_2H_5OH (and any strong acid) ALLOW ethanol (1)	OHCH₂CH₃	4

Question Number	Acceptable Answers	Reject	Mark
16(c)	2CH ₃ CH ₂ COOH + Na ₂ CO ₃ →		
	$2CH_3CH_2CH_2COO^{(-)}Na^{(+)} + CO_2 + H_2O$		
	ALLOW		
	butanoic acid as CH ₃ CH ₂ CH ₂ CO ₂ H/ CH ₃ (CH ₂) ₂ COOH/CH ₃ (CH ₂) ₂ CO ₂ H/ C ₃ H ₇ COOH/C ₃ H ₇ CO ₂ H		
	and the salt as $CH_3CH_2CH_2CO_2H$ / $CH_3(CH_2)_2$ $COO^{(-)}Na^{(+)}$ / $CH_3(CH_2)_2CO_2^{(-)}Na^{(+)}$ / $C_3H_7COO^{(-)}Na^{(+)}$ / $C_3H_7CO_2^{(-)}Na^{(+)}$		
	all product formulae correct (1) correct balanced equation (1)		
	ALLOW correct ionic equation for (1) $2CH_3CH_2COOH + CO_3^{2-} \rightarrow$		
	$2CH_3CH_2CH_2COO^- + CO_2 + H_2O$		
	I GNORE state symbols even if incorrect		2

Question Number	Acceptable Answers	Reject	Mark
16(d)	Any two correct points from: First point butanoic acid has 4 peaks, butan-1-ol has 5 peaks OR butanoic acid has one peak fewer OR butan-1-ol has one peak more ALLOW butanoic acid has fewer peaks/ butan-1-ol has more peaks (1) I GNORE butanoic acid has 4 proton environments and butan-1-ol has 5	incorrect numbers of peaks quoted different number of peaks	
	Second point ratio of peak heights/ area under each peak is 3:2:2:1 for butanoic acid and 3:2:2:2:1 for butan-1-ol (1) Third point the OH (hydrogens) have different chemical shifts	area under peaks in the ratio 8:10	
	OR butanoic acid has a (COOH) peak at 10-12 (ppm) (and butan-1-ol does not) OR butan-1-ol has (an OH) peak at 2-4 (ppm) (and butanoic acid does not) (1)	incorrect data quoted	
	Fourth point peak at 3.0-1.8 (ppm) for H-C-C=O in acid and not in the alcohol OR peak at 3.0-4.4 (ppm) for H-C-O- in alcohol and not in acid OR the hydrogens on the alpha carbon have different chemical shifts (1)	incorrect data quoted	
	I GNORE reference to splitting patterns		2

Question Number	Acceptable Answers	Reject	Mark
16(e)	First mark – bond and range C=O(stretching) in butanoic acid (has an absorption at) 1725 – 1700 (cm ⁻¹) (1)	COOH/ incorrect name of bond/ 1740 – 1720 (cm ⁻¹)/ other incorrect range	
	Second mark – bond and both ranges O-H/ OH (stretching) in butan-1-ol 3750 – 3200 (cm ⁻¹) and O-H/ OH (stretching) in butanoic acid 3300 – 2500 (cm ⁻¹) ALLOW COOH in butanoic acid (1)	incorrect name of bonds	
	ALLOW any wavenumber or range of wavenumbers within the ranges above and ranges written in reverse order		
	If no other marks are awarded, then ALLOW 1 mark if all 3 ranges are identified but bonds are missing/incorrect		
	I GNORE reference to fingerprint region		2

Question Number	Acceptable Answers	Reject	Mark
16(f)	مأم		
	I GNORE bond lengths and bond angles ALLOW any orientation		1

Question Number	Acceptable Answers	Reject	Mark
16(g)	First step – PCI ₅ / phosphorus(V) chloride/ phosphorus pentachloride	HCI	
	ALLOW PCl ₃ / phosphorus(III) chloride/ phosphorus trichloride SOCl ₂ / thionyl chloride (1)		
	Second step - conditional on first mark CH ₃ CH ₂ OH/ C ₂ H ₅ OH/ ethanol (1)		
	Advantage - stand alone mark higher yield (of ester) OR		
	reaction goes to completion/ not an equilibrium reaction/ not reversible OR		
	no heat energy needed/ reacts at room temperature/ no (concentrated acid) catalyst needed (1)		
	I GNORE atom economy/ faster/		
	requires less energy		3

Total for Question 16 = 15 marks

Question Number	Acceptable Answers		Reject	Mark
17(a)				
	Method 1 – gas collection Diagram 2 marks stoppered/ sealed side arm test tube/ stoppered/ sealed test tube with delivery tube/ stoppered/ sealed side arm flask/ stoppered/ sealed flask with delivery tube	(1)		
	gas syringe OR collection of gas over water in a measuring cyling upturned burette/ graduated gas tube I GNORE heat	der/ (1)		
	Measurements volume of gas and time	(1)		
	Method 2 - mass loss Diagram 2 marks digital balance flask with cotton wool/ mineral wool in neck	(1)	amount of gas	
	OR open flask/ beaker	(1)		
	Measurements mass (loss) and time	(1)		
	Method 3 – colour change Diagram 2 marks colorimeter light and filter shown	(1) (1)		
	Measurements transmission/ absorbance and time	(1)		3

Question Number	Acceptable Answers	Reject	Mark
17(b)(i)	s ⁻¹ ALLOW 1/s sec ⁻¹ any actual unit of time to power -1	time ⁻¹ t ⁻¹	1

Question Number	Acceptable Answers	Reject	Mark
17(b)*(ii)	First mark 1 st Step – slow 2 nd Step – fast 3 rd Step – fast		
	Second mark – stand alone the slow(est)/ first step is the rate determining step (1)		
	Third mark – consequential on correct first mark (1 mol) N_2O_5 is in the rate equation so the reaction with N_2O_5 is the slow/ rate determining step OR only the species in the rate equation is in the first/ slow/ rate determining step ALLOW (there is only 1 mol of) one species/ N_2O_5 in the first/ slow/ rate determining step (1)		
	ALLOW 1 st Step – fast 2 nd Step – slow 3 rd Step – fast (1)		
	the slow(est) step/second step is the rate determining step (1)		
	there is only (1 mol of) one species in the steps up to and including the rate determining step (1)		3

Question Number	Acceptable Answers	Reject	Mark
17(c)(i)	(thermostatically controlled) water bath/ ice bath	direct heating with flame	1
	ALLOW oil bath	electrical heater	

Question Number	Acceptable Answers		Reject	Mark
17(c)(ii)	$(1/T) 3.13 \times 10^{-3} / 3.125 \times 10^{-3}$	(1)	3.12 x 10 ⁻³	
	(lnk) -7.1/ -7.05/ -7.0528	(1)	-7.0	2

Number 17(c) * (iii) In K -1 3.1 3.2 3.3 3.4 3.5 -1 1/0 5 1 -5 -1 -1 -1 -1 -1 -1 -6 -1 -1 -1 -1 -1 -1 -7 -1 -1 -1 -1 -1 -8 -1 -1 -1 -1 -8 -1 -1 -1 -1 -9 -1 -1 -1 -1 -9 -1 -1 -1 -9 -1 -1 -1 -9 -1 -1 -1 -9 -1 -1 -1 -9 -1 -1 -1 -9 -1 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1 -1 -9 -1	
Graph – 3 marks First mark axes correct with sensible scales i.e. points/line covering at least 3 large squares on the x axis and 5 on the y axis, with lnk values becoming more negative down the axis and the negative signs shown (1) Second mark both axes labelled, with units on x axis and no units on y axis x axis: 3.1-3.5 1/T /10 ⁻³ K ⁻¹ OR 3.1-3.5 1/Tx10 ³ /K ⁻¹ OR 0.0031-0.0035 OR 3.1 x 10 ⁻³ -3.5 x 10 ⁻³ 1/T /K ⁻¹ ALLOW x axis labels at bottom of page (1) Third mark points correctly plotted and straight line drawn (1)	

```
Gradient - 2 marks - this may be shown on the
gradient = -11550 to -12760 (K)
negative sign
                                                        (1)
value (ignore sf)
                                                        (1)
maximum 1 mark if an incorrect unit is given
ALLOW these marks if the correct sign and value for the
gradient are shown in the calculation for E_a
Calculation - 2 marks
If E_a = (+)96.0 \text{ to } (+)106 \text{ kJ mol}^{-1} \text{ or } (+)96000 \text{ to}
(+)106000 J mol<sup>-1</sup>, award 2 marks
If E_a is in this range but is not given to 3 sf, or the
units are incorrect or missing, award 1 mark
If not.
E_a = -8.31 \text{ x their gradient}
OR
                                                       (1)
gradient = -Ea/R
value to 3 sf and units and consequential sign if negative
                                                        (1)
ALLOW
correct answer to 3 sf, in range, with sign and units,
but no working for gradient or E_a
                                                                  7
```

Total for Question 17 = 17 marks

Acceptable Answers	Reject	Mark
I GNORE sf except 1		
If answer is 8.485 x 10 ⁻³ (moldm ⁻³), award 2 marks		
If not, $[OH^{-}(aq)] = \sqrt{(K_b [NH_3])}$		
$= \sqrt{(1.8 \times 10^{-5} \times 4.0)} \tag{1}$		
= $8.485 \times 10^{-3} \text{ (mol dm}^{-3}\text{)}$ (1)		2
	IGNORE sf except 1 If answer is 8.485×10^{-3} (moldm ⁻³), award 2 marks If not, $[OH^{-}(aq)] = \sqrt{(K_b [NH_3])}$ $= \sqrt{(1.8 \times 10^{-5} \times 4.0)}$ (1)	I GNORE sf except 1 If answer is 8.485×10^{-3} (moldm ⁻³), award 2 marks If not, $[OH^{-}(aq)] = \sqrt{(K_b [NH_3])}$ $= \sqrt{(1.8 \times 10^{-5} \times 4.0)}$ (1)

Question Number	Acceptable Answers		Reject	Mark
18(a)(ii)	I GNORE sf except 1			
	If answer is 11.9(3)/12, award 2 marks			
	If not,			
	EITHER – Method 1 $[H^{+}] = \frac{1 \times 10^{-14}}{[OH^{-}]}$			
	$= \underbrace{1 \times 10^{-14}}_{8.485 \times 10^{-3}}$	(1)		
	= 1.179×10^{-12} ALLOW ecf from their answer to (i)			
	pH = $-\log 1.179 \times 10^{-12} = 11.9(3)$			
	ALLOW ecf from their answer for [H ⁺]	(1)		
	OR — Method 2			
	pOH = $-\log 8.485 \times 10^{-3} = 2.07$ ALLOW ecf from their answer to (i)	(1)		
	pH = $(14 - 2.07 =) 11.9(3)$ ALLOW ecf from their answer to pOH	(1)		
				2

Question Number	Acceptable Answers	Mark
18(8)(1)	$(pH_{12} - log 4.0)$ = 0.6(021)	1
	11 - 10 - 9 - 8 - 7 - PH 6 -	
	5 - 4 - 3 - 2 - 1 - 0 1 - 0 - 5 10 15 20 25 30 Volume of HCI(aq)/cm ³	
	First mark graph starting at 11.9/ answer to a(ii), ± 1 small square, provided above 7 (1) Second mark buffering region to 25 cm ³ ALLOW any line showing a decrease in pH from 0 to 25 cm ³ of HCl	
	added (1) Third mark straight vertical portion between 8 and 1, midpoint below 7 and between 2 and 7 pH units long (1)	
	Fourth mark finishing at $+0.5$ to -0.8 , with at least 27.5 cm ³ of HCl added ALLOW final pH as answer to (b)(i), within 1 pH unit, if pH is less than answer to (b)(i) or within 1 small square if pH is more than answer to (b)(i) (1)	
	ALLOW If graph is drawn with aqueous ammonia added to hydrochloric acid, only the second and third marks are available for the correct vertical portion at 25cm ³	4

Question Number	Acceptable Answers	Reject	Mark
18(b)*(iii)		universal indicator loses all 3 marks	
	First mark any indicator from 4 to 10 or 12, 13 in the Data booklet – see end ALLOW ecf from the vertical portion on their graph (1)	litmus loses first mark only	
	Second mark alkaline to acidic colour change for their stated indicator ALLOW acidic to alkaline colour change if their curve shows alkali added to acid (1)		
	Third mark pH range (of indicator) is within the vertical section of the graph OR pKin (± 1) is in the vertical section of the graph OR pKin is nearest to the pH at the end/equivalence point ALLOW indicator will change colour in the vertical		
	section of the graph ALLOW Indicator will change colour at the end/ equivalence point ALLOW (because it is a) titration of a strong acid with a weak base (1)		3

Question Number	Acceptable Answers	Mark
18(c)(i)	I GNORE sf except 1	
	If answer is 3.84 (mol dm ⁻³), award 3 marks If not,	
	number of moles of acid = $\frac{24.0 \times 4}{1000} = 0.096 \tag{1}$	
	EI THER number of moles ammonia = $0.096 \text{ in } 25 \text{ cm}^3$ (1)	
	concentration of ammonia $= 0.096 \times 1000$	
	$ = 3.84 \text{ (mol dm}^{-3}) $ (1)	
	OR number of moles ammonia = $0.288 \text{ in } 75 \text{ cm}^3$ (1)	
	concentration of ammonia $= 0.288 \times 1000$	
	$75 = 3.84 \text{ (mol dm}^{-3}\text{)} $ (1)	
	I GNORE unit unless incorrect	
	ALLOW ecf in both methods on their number of moles of ammonia	3

Question Number	Acceptable Answers	Mark
18(c)(ii)	I GNORE sf except 1 (concentration of ammonia in trichloromethane =) 0.16 (mol dm ⁻³)	
	ALLOW ecf from their answer to (c)(i), provided it is less than 4.0 and given to 2 or more sf	1

Question Number	Acceptable Answers	Reject	Mark
18(c)(iii)	Expression for Kc and answer needed for the mark $K_c = \frac{[NH_3(aq)]}{[NH_3(CHCl_3)]}$ ALLOW one state symbol missing $= \frac{3.84}{0.16}$ = 24(.0)	K _c expressions without both state symbols	
	I GNORE sf, including 1 sf, and units ALLOW ecf from answers to (c)(i) and (c)(ii)		1

Question Number	Acceptable Answers	Reject	Mark
18(c)(iv)	(ammonia/ it is much more soluble in water) as can form hydrogen bonds with water		
	ALLOW more/ stronger hydrogen bonds with water (than with trichloromethane) I GNORE answers based on polarity/ hydrophilic		
	randing answers susce on polarity, mydrophine		1

Total for Question 18 = 18 marks

Question Number	Acceptable Answers	Mark
19(a)(i)	Penalise lack of + sign once only in (a)(i) or (ii) in each final answer	
	IGNORE sf in (a)(i), (ii), and (iii) in each final answer, except 1 sf	
	FIRST, CHECK THE FINAL ANSWER +479.7 J mol ⁻¹ K ⁻¹ scores 3 marks	
	479.7 J mol ⁻¹ K ⁻¹ scores 2 marks (+ sign missing)	
	+479.7/ 479.7 scores 2 marks (units and/or + missing)	
	+1709.7 J mol ⁻¹ K ⁻¹ scores 2 marks – multiple of 12 used for oxygen	
	$1709.7 \ \mathrm{J} \ \mathrm{mol}^{-1} \ \mathrm{K}^{-1}/\ +1709.7/\ 1709.7 \ \mathrm{score} \ 1 \ \mathrm{mark} - \mathrm{multiple} \ \mathrm{of} \ 12$ used for oxygen and positive sign and/or units	
	If these answers are not given, award marks as follows:	
	First mark correct data for CO_2 (213.6) and H_2O (69.9) (1)	
	Second mark correct multiples (12, 11, 1 and 24) and Hess's Law applied $\Delta S^{\circ}_{\text{system}} = 12 \times 213.6 + 11 \times 69.9$ $-(392.4 + 24 \times 102.5)$	
	ALLOW ecf from incorrect data for CO_2 and/or H_2O (1)	
	Third mark correct answer with sign and units = +479.7 J mol ⁻¹ K ⁻¹	
	ALLOW ecf from incorrect data for CO_2 and/or H_2O and incorrect multiples (1)	3

Question Number	Acceptable Answers		Reject	Mark
19(a)(ii)	If answer is + 18925.2 J mol ⁻¹ K ⁻¹ / + 18.9252 kJ mol ⁻¹ K ⁻¹ , then award 2 marks If not, $\Delta S^{e}_{surroundings} = \frac{-\Delta H^{e}}{T}$ = - $\frac{(-5639.7) \times 1000}{298}$ = + 18925.2 J mol ⁻¹ K ⁻¹ / + 18.9252 kJ mol ⁻¹ K ⁻¹	(1)	+18925.1 J mol ⁻¹ K ⁻¹ / +18.9251 kJ mol ⁻¹ K ⁻¹	2
				2

Question Number	Acceptable Answers	Mark
19(a)(iii)	First mark $(\Delta S^{e}_{total} = \Delta S^{e}_{surroundings} + \Delta S^{e}_{system} = 18925.2 + 479.7)$	
	= $(+)19404.9$ (J mol ⁻¹ K ⁻¹)/ $(+)19.4049$ (kJ mol ⁻¹ K ⁻¹)	
	if units given they must be correct	
	ALLOW (+)19500 (J mol ⁻¹ K ⁻¹)/ (+)19.5 (kJ mol ⁻¹ K ⁻¹) (from 19.0 + 0.480)	
	ALLOW ecf on adding answers to (a)(i) and (a)(ii) in the same units (1)	
	Note If answer to (a)(i) was +1709.7, $\Delta S^{\circ}_{\text{total}} = +20634.9 \text{ (J mol}^{-1} \text{ K}^{-1}\text{) } / +20.6349 \text{ (kJ mol}^{-1} \text{ K}^{-1}\text{)}$	
	Second mark ($\Delta S^{\rm e}_{\rm total}$ is positive so) reaction is (thermodynamically) spontaneous/ feasible/ goes to completion	
	ALLOW thermodynamically unstable	
	If their sign for ΔS^{e}_{total} is negative, then ALLOW reaction is not spontaneous/ not feasible/ does not go to completion (1)	2

Question Number	Acceptable Answers	Reject	Mark
19(a)(iv)	I GNORE comments on $\Delta S^{ m e}_{ m system}$		
	First mark $(\Delta S^{e}_{surroundings} = -\Delta H^{e}/T \text{ so increase in T makes})$		
	$\Delta S^{\rm e}_{ m surroundings}$ less positive/ decreases	more exothermic	
	ALLOW more negative (1)		
	Second mark $(\Delta S^{\circ}_{total} = \Delta S^{\circ}_{surroundings} + \Delta S^{\circ}_{system}$ so increase in T makes)		
	$\Delta S^{ m e}_{ m total}$ less positive/ decreases		
	ALLOW more negative NOTE no ecf on $\Delta S^{\circ}_{\text{surroundings}}$ increases (1)		
	Third mark (because ΔS°_{total} is so large and positive to start with) there is an insignificant effect on the extent of the reaction ALLOW ΔS°_{total} is still positive so reaction still goes to completion/is spontaneous ALLOW ecf on ΔS°_{total} increases (1)		
			3

Question Number	Acceptable Answers	Reject	Mark
19(a)(v)	First mark (stable because) high activation energy/ E_a (for combustion of sucrose) ALLOW sucrose is kinetically stable/ inert (1) Second mark (hazardous because small particles/ powder have/ has) larger surface area and react faster (1) I GNORE any reference to temperature If answers are not linked to stability and hazardous, still		
	award both marks even if the points are written in the wrong order		2

Question Number	Acceptable Answers	Reject	Mark
19(a)(vi)	Any two of:		
	obesity/ weight gain/ stored as fat/ get fat (1)		
	tooth decay/ cavities/ toothache (1)		
	diabetes/ glycosuria (1)		
	heart/ cardiovascular condition/ disease/ attack (1)		
	strokes (1)		
	damage to the immune system (1)		
	high insulin levels (1)		
	high blood pressure (1)		
	kidney damage (1)		
	liver disease (1)		
	headaches/ migraines (1)		
	arthritis (1)		
	high cholesterol (1)	
	I GNORE risk of cancer/ high blood sugar/ stomach ulcers		2

Question Number	Acceptable Answers		Reject	Mark
19(b)(i)	circles or asterisks on carbons 2-5		all 6 carbons	
	all four correct	(2)	circled (0)	
	3 or 2 correct	(1)		
	1 or 0 correct	(0)		
	ALLOW 5 carbons circled	(1)		2

Question Number	Acceptable Answers	Reject	Mark
19(b)(ii)	rotate the plane of (plane-) polarized light	just 'rotate light'	
	ALLOW rotate plane-polarized light		
	I GNORE optically active/ optical activity/ non- superimposable		1

Question Number	Acceptable Answers	Reject	Mark
19(b)(iii)	First mark – colour change from a blue (solution) to a red/ orange/ brown/ yellow precipitate ALLOW solid or (s) for precipitate which could be shown in formula or equation (1)	incorrect observation for one of the reagents for first mark only, eg. silver mirror formed	
	Second mark – functional group (glucose/it is) an aldehyde / (has) a CHO group (1)		
	Third mark - oxidation/reduction		
	copper(II)/Cu ²⁺ is reduced (to copper(I)/Cu ⁺ oxide by the aldehyde group) /Cu ²⁺ + $e^{(-)} \rightarrow Cu^+$		
	OR the aldehyde/ glucose is oxidized (to the carboxylate/carboxylic acid)/ RCHO + [O] → RCOOH		
	OR Benedict's and Fehling's (solutions) are oxidizing agents		
	ALLOW equation showing oxidation of aldehyde and reduction of Cu ²⁺ even if not balanced (1)		3

Total for Question 19 = 20 marks

Pearson Education Limited. Registered company number 872828 with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE