

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

Thursday 7 November 2019

Morning (Time: 1 hour 15 minutes)

Paper Reference **WCH06/01**

Chemistry

Advanced

Unit 6: Chemistry Laboratory Skills II

Candidates must have: Scientific calculator

Total Marks

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P58301A

©2019 Pearson Education Ltd.

1/1/1/



Pearson

Answer ALL the questions. Write your answers in the spaces provided.

1 A pale green crystalline solid **A** contains two cations and one anion.

(a) When **A** is warmed with aqueous sodium hydroxide, a gas is evolved that turns damp red litmus paper blue.

(i) Identify, by name or formula, the gas evolved.

(1)

(ii) Give the name or formula of the cation in **A** that is identified by this test.

(1)

(b) **A** dissolves in distilled water to form a very pale green solution **B**.

B reacts with aqueous sodium hydroxide to form a green precipitate, which turns into a brown solid **C**, on standing in air.

(i) Give the name or formula of the cation in **B** that is identified by this test.

(1)

(ii) Identify, by name or formula, the brown solid **C**.

(1)

(c) **B** gives a white precipitate when aqueous barium chloride acidified with dilute hydrochloric acid is added.

Give the name or formula of the anion in **B** that is identified by this test.

(1)

(d) Suggest the **formula** of solid **A**. Do not include any water of crystallisation.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (e) A sample of 0.025 mol of solid **A** with a mass of 9.80 g is heated gently to remove the water of crystallisation and leave 0.025 mol of the anhydrous solid.

The mass of anhydrous solid is 7.10 g.

Calculate the number of moles of water of crystallisation combined with 1 mol of the anhydrous solid.

(2)

(Total for Question 1 = 8 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



2 **W** is a white solid with the molecular formula $C_9H_8O_2$.

(a) A series of tests is carried out on **W**.
Complete the table.

Test	Observation	Inference
(i) Ignite a sample of W	Very smoky flame	W could be an alkene or compound (1)
(ii) Add a little W to bromine water and shake the mixture	Yellow solution turns into a colourless solution	W contains the group (1)
(iii) Heat W until it melts then add phosphorus(V) chloride	Steamy fumes form	W contains the group (1)
(iv) Heat W until it melts then add solid	Bubbles of carbon dioxide form	W contains the group (2)

(b) Complete the table, which contains information about the mass spectrum of **W**.

Peak	Inference
(i) A peak occurs at $m/e =$	The peak is due to $C_6H_5^+$ (1)
(ii) A peak occurs at $m/e = 103$	The peak is due to an ion with the formula (1)



(c) The **low** resolution proton nmr spectrum of **W** has four peaks each with relative area 1 and two peaks each with relative area 2.

(i) State the number of proton environments in **W**.

(1)

(ii) State what can be deduced from the relative peak areas.

(1)

(d) **W** exists as two geometric isomers.

Use all the information in this question to deduce the structure of **one** of these isomers.

(2)

(Total for Question 2 = 11 marks)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



3 A student used two methods to determine the concentration of vanadium(III) ions in an aqueous solution **X**.

(a) **Method 1** used a titration procedure.

10.0 cm³ of **X** was titrated with 0.0400 mol dm⁻³ acidified potassium manganate(VII).

The equation for the reaction is



The results of four titrations are shown.

Titration	Rough	1	2	3
Final burette reading / cm ³	21.10	41.30	19.85	20.10
Initial burette reading / cm ³	0.50	21.10	0.25	0.00
Titre / cm ³				20.10
Titres used to calculate mean				

- (i) Complete the table and calculate the mean titre.
Show which titres you have used in your calculation by putting a tick (✓) in the appropriate boxes in the table.

(2)

Mean titre = cm³



- (ii) Calculate the concentration, in mol dm^{-3} , of $\text{V}^{3+}(\text{aq})$ ions in solution **X**.
Give your answer to **three** significant figures.

(3)

- (iii) Each burette reading was accurate to $\pm 0.05 \text{ cm}^3$.

Calculate the percentage uncertainty in the titre value for Titration **3**.

(1)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA




(b) **Method 2** used an electrochemical cell.

An electrochemical cell was made from the electrode systems represented by these half-equations:



The E_{cell} value was measured using the apparatus shown.

Solution **Y** was made by mixing 50 cm^3 of an aqueous solution of V^{2+} ions with 50 cm^3 of the same solution **X** as used in **Method 1**.



- (i) Complete the diagram by adding labels on the dotted lines provided. Conditions are not required.

(4)

- (ii) The salt bridge consisted of a strip of filter paper soaked in a saturated solution of potassium nitrate.

Give a reason why potassium hydroxide solution should **not** be used for the salt bridge.

(1)

.....

.....

.....



- (iii) In this cell, the zinc half-cell was at standard temperature and concentration. When the cell reaction occurred, the zinc was oxidised and $E_{\text{cell}} = +0.44\text{V}$.

Write the overall equation for the cell reaction.
State symbols are not required.

(1)

- (iv) The standard electrode potential, E^\ominus , for the $\text{Zn}^{2+}(\text{aq})|\text{Zn}(\text{s})$ half-cell = -0.76V .

The $\text{V}^{3+}(\text{aq})|\text{V}^{2+}(\text{aq})$ half-cell was **not** at standard concentration in this experiment.

Calculate the electrode potential, E , for the $\text{V}^{3+}(\text{aq})|\text{V}^{2+}(\text{aq})$ half-cell in this experiment.

(1)

- (v) The **standard** electrode potential, E^\ominus , for the $\text{V}^{3+}(\text{aq})|\text{V}^{2+}(\text{aq})$ half-cell = -0.26V .

Solution **Y** was 1 mol dm^{-3} with respect to $\text{V}^{2+}(\text{aq})$.

For the half-cell in this experiment, the electrode potential is given by

$$E = E^\ominus + 0.059 \log [\text{V}^{3+}(\text{aq})]$$

Use this, and your answer to (b)(iv), to calculate the concentration of $\text{V}^{3+}(\text{aq})$ in solution **Y**. You **must** show your working.

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(c) The concentration of $V^{3+}(aq)$ obtained in (a)(ii) was approximately double that obtained in (b)(v).

Explain why these two values were different.

(1)

.....

.....


.....

.....

(Total for Question 3 = 16 marks)



- 4 This question is about the preparation of iodobenzene from phenylamine, and its purification. The preparation occurs in two steps.



Some data about phenylamine and iodobenzene are given in the table.

Compound	Molar mass / g mol^{-1}	Density / g cm^{-3}	Boiling temperature / $^{\circ}\text{C}$
Phenylamine	93.0	1.02	184
Iodobenzene	203.9	1.83	188

- (a) In Step 1 of the preparation, phenylamine is converted into benzenediazonium chloride. Give the reagents and condition for Step 1.

(2)

.....

.....

.....

.....


(b) In Step 2 of the preparation, aqueous potassium iodide is added slowly to the reaction mixture from Step 1.

The mixture is left to stand for 10 minutes and then it is heated for 20 minutes. The iodobenzene formed is steam distilled from the mixture.

(i) Suggest a reason why the aqueous potassium iodide is added slowly.

(1)

(ii) The apparatus used for steam distillation is shown.



Complete the labelling of the diagram, **A**, **B**, **C** and **D**.

(3)

- A
- B
- C
- D



(iii) State the purpose of the part of the apparatus labelled **E**.

(1)

(iv) The distillate collected contains iodobenzene and water.

Describe how iodobenzene is obtained from the distillate.

[Refer to the data given at the start of Question 4]

(2)

(v) The iodobenzene obtained from the distillate is a cloudy liquid.

Name a substance that should be added to make the liquid clear.

(1)

(vi) The clear liquid is distilled to obtain pure iodobenzene.

Give a suitable temperature **range** for collecting the pure iodobenzene.

[Refer to the data given at the start of Question 4]

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(c) This preparation and purification process has an expected yield of 70%.

Calculate the **volume** of phenylamine needed to produce 25.0 cm^3 of iodobenzene.

[Refer to the data given at the start of Question 4]

(4)

(Total for Question 4 = 15 marks)

TOTAL FOR PAPER = 50 MARKS



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



The Periodic Table of Elements

	1	2	Key										0 (8)					
			(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)
			relative atomic mass															
			atomic symbol															
			name															
			atomic (proton) number															
6.9	Li	9.0	45.0	47.9	50.9	52.0	54.9	55.8	58.9	58.7	63.5	65.4	10.8	12.0	14.0	16.0	19.0	4.0
	lithium	beryllium	Scandium	titanium	vanadium	chromium	manganese	iron	cobalt	nickel	copper	zinc	boron	carbon	nitrogen	oxygen	fluorine	helium
23.0	Na	24.3	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	27.0	28.1	31.0	32.1	35.5	39.9
	sodium	magnesium	Yttrium	zirconium	niobium	molybdenum	technetium	ruthenium	rhodium	palladium	silver	cadmium	aluminium	silicon	phosphorus	sulfur	chlorine	argon
39.1	K	40.1	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	69.7	72.6	74.9	79.0	79.9	83.8
	potassium	calcium	Scandium	titanium	vanadium	chromium	manganese	iron	cobalt	nickel	copper	zinc	gallium	germanium	arsenic	selenium	bromine	krypton
85.5	Rb	87.6	88.9	91.2	92.9	95.9	[98]	101.1	102.9	106.4	107.9	112.4	114.8	118.7	121.8	127.6	126.9	131.3
	rubidium	strontium	Yttrium	zirconium	niobium	molybdenum	technetium	ruthenium	rhodium	palladium	silver	cadmium	indium	tin	antimony	tellurium	iodine	xenon
132.9	Cs	137.3	138.9	178.5	180.9	183.8	186.2	190.2	192.2	195.1	197.0	200.6	204.4	207.2	209.0	209.0	[210]	[222]
	caesium	barium	lanthanum	hafnium	tantalum	tungsten	rhenium	osmium	iridium	platinum	gold	mercury	thallium	lead	bismuth	polonium	astatine	radon
[223]	Fr	[226]	[227]	[261]	[262]	[266]	[264]	[277]	[268]	[271]	[272]	Elements with atomic numbers 112-116 have been reported but not fully authenticated						
	francium	radium	actinium	rutherfordium	roentgenium	seaborgium	bohrium	hassium	meitnerium	darmstadtium	roentgenium							
140	Ce	141	144	[147]	150	152	157	159	163	165	167	169	173	175				
	cerium	praseodymium	neodymium	promethium	samarium	europium	gadolinium	terbium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium				
232	Th	[231]	238	[237]	[242]	[243]	[247]	[245]	[251]	[254]	[253]	[256]	[254]	[257]				
	thorium	protactinium	uranium	neptunium	plutonium	americium	curium	berkelium	californium	einsteinium	fermium	mendelevium	nobelium	lawrencium				

* Lanthanide series

* Actinide series

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

