

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--	--

## Pearson Edexcel International Advanced Level

Time 1 hour 30 minutes

Paper  
reference

**WCH12/01**



### Chemistry

#### International Advanced Subsidiary/Advanced Level UNIT 2: Energetics, Group Chemistry, Halogenoalkanes and Alcohols

#### You must have:

Scientific calculator, Data Booklet, ruler

Total Marks

#### Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
  - *there may be more space than you need.*

#### Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets
  - *use this as a guide as to how much time to spend on each question.*
- In the question marked with an **asterisk (\*)** marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- A Periodic Table is printed on the back cover of this paper.

#### Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P69501A

©2022 Pearson Education Ltd.

L:1/1/1/




P 6 9 5 0 1 A 0 1 2 8



Pearson

**SECTION A****Answer ALL the questions in this section.****You should aim to spend no more than 20 minutes on this section.****For each question, select one answer from A to D and put a cross in the box  . If you change your mind, put a line through the box  and then mark your new answer with a cross  .**

- 1 Which are correct for the reaction shown?



	Type of reaction	Type of enthalpy change
<input type="checkbox"/> A	endothermic	formation
<input checked="" type="checkbox"/> B	endothermic	neutralisation
<input type="checkbox"/> C	exothermic	formation
<input type="checkbox"/> D	exothermic	neutralisation

**(Total for Question 1 = 1 mark)**

- 2 Which equation does **not** represent a standard enthalpy change of atomisation?

- A  $\text{Mg(s)} \rightarrow \text{Mg(g)}$
- B  $\text{Cl}_2(\text{g}) \rightarrow 2\text{Cl(g)}$
- C  $\frac{1}{2}\text{O}_2(\text{g}) \rightarrow \text{O(g)}$
- D  $\text{Hg(l)} \rightarrow \text{Hg(g)}$

**(Total for Question 2 = 1 mark)****Use this space for any rough working. Anything you write in this space will gain no credit.**

- 3** 5.20 g of sodium hydrogencarbonate is added to an excess of acid.

The temperature increases and the energy change is calculated to be 1030 J.


What is the enthalpy change per mole of sodium hydrogencarbonate?

$$[M_r \text{ NaHCO}_3 = 84.0]$$

- A**  $-12.3 \text{ kJ mol}^{-1}$
- B**  $-16.6 \text{ kJ mol}^{-1}$
- C**  $-63.8 \text{ kJ mol}^{-1}$
- D**  $-16\,600 \text{ kJ mol}^{-1}$

(Total for Question 3 = 1 mark)

- 4** The equation for the complete combustion of methanal is shown.



Some bond enthalpy data are shown.

Bond	Bond enthalpy / $\text{kJ mol}^{-1}$
C—H	413
O=O	498
C=O in $\text{CO}_2$	805
O—H	464

What is the C=O bond enthalpy in methanal?

- A**  $623 \text{ kJ mol}^{-1}$
- B**  $678 \text{ kJ mol}^{-1}$
- C**  $805 \text{ kJ mol}^{-1}$
- D**  $1036 \text{ kJ mol}^{-1}$

(Total for Question 4 = 1 mark)



P 6 9 5 0 1 A 0 3 2 8

5 Which sequence shows the molecules in order of **increasing** boiling temperature?

- A  $\text{H}_2\text{O} < \text{Br}_2 < \text{Cl}_2 < \text{CH}_4$
- B  $\text{Br}_2 < \text{CH}_4 < \text{Cl}_2 < \text{H}_2\text{O}$
- C  $\text{Cl}_2 < \text{CH}_4 < \text{H}_2\text{O} < \text{Br}_2$
- D  $\text{CH}_4 < \text{Cl}_2 < \text{Br}_2 < \text{H}_2\text{O}$


(Total for Question 5 = 1 mark)

6 Which is **not** correct about ice?

- A ice has a lower density than water
- B  $\text{H}_2\text{O}$  molecules are further apart in ice than in water
- C the H–O–H bond angle is the same in ice and in water
- D  $\text{H}_2\text{O}$  molecules in ice are held together by hydrogen bonds

(Total for Question 6 = 1 mark)

7 Which intermolecular forces exist **between** the molecules of the compound shown?



- A hydrogen bonding and London forces only
- B hydrogen bonding and permanent dipole-permanent dipole forces only
- C London forces and permanent dipole-permanent dipole forces only
- D hydrogen bonding, permanent dipole-permanent dipole forces and London forces

(Total for Question 7 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



8 This question is about alkanes.





(a) Which of these alkanes has the **highest** boiling temperature?

(1)

- A butane
- B hexane
- C pentane
- D propane

(b) Which of these alkanes has the **lowest** boiling temperature?

(1)

- A 
- B 
- C 
- D 

(Total for Question 8 = 2 marks)

9 Which solvent dissolves the greatest amount of hydrocarbon  $C_{35}H_{72}$ ?

- A butan-1-ol
- B ethanoic acid
- C hexane
- D water

(Total for Question 9 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



**10** Which reagent would convert an alcohol into an alkene?

- A** acidified potassium dichromate(VI)
- B** anhydrous calcium sulfate
- C** concentrated phosphoric acid
- D** ethanolic potassium hydroxide

(Total for Question 10 = 1 mark)

**11** Which name is correct for the ion  $\text{SO}_4^{2-}$ ?

- A** sulfate(II)
- B** sulfate(IV)
- C** sulfate(VI)
- D** sulfate(VIII)

(Total for Question 11 = 1 mark)

**12** In which compound is the oxidation number of nitrogen +5?

- A**  $\text{Ca}(\text{NO}_3)_2$
- B**  $\text{Mg}_3\text{N}_2$
- C**  $\text{N}_2\text{O}_3$
- D**  $\text{NaNO}_2$

(Total for Question 12 = 1 mark)

**13** In which reaction is the copper species acting as an oxidising agent?


- A**  $\text{Cu}^{2+} + 2\text{Ag} \rightarrow 2\text{Ag}^+ + \text{Cu}$
- B**  $2\text{Cu}^+ + \text{O}^{2-} \rightarrow \text{Cu}_2\text{O}$
- C**  $3\text{Cu} + \text{O}_2 \rightarrow \text{Cu}_2\text{O} + \text{CuO}$
- D**  $\text{Cu} + \text{Hg}^{2+} \rightarrow \text{Hg} + \text{Cu}^{2+}$

(Total for Question 13 = 1 mark)

**Use this space for any rough working. Anything you write in this space will gain no credit.**



14 Two half-equations for a reaction are shown.



What is the overall ionic equation for this reaction?

- A  $\text{Cu} + \text{NO}_3^- + 4\text{H}^+ \rightarrow \text{Cu}^{2+} + \text{NO} + 2\text{H}_2\text{O}$
- B  $2\text{Cu} + \text{NO}_3^- + 4\text{H}^+ \rightarrow 2\text{Cu}^{2+} + \text{NO} + 2\text{H}_2\text{O}$
- C  $3\text{Cu} + 2\text{NO}_3^- + 8\text{H}^+ \rightarrow 3\text{Cu}^{2+} + 2\text{NO} + 4\text{H}_2\text{O}$
- D  $6\text{Cu} + 2\text{NO}_3^- + 8\text{H}^+ \rightarrow 6\text{Cu}^{2+} + 2\text{NO} + 4\text{H}_2\text{O}$

(Total for Question 14 = 1 mark)

15 A titre of  $13.25 \text{ cm}^3$  was obtained using a  $50 \text{ cm}^3$  burette.


What is the percentage uncertainty in the titre?

[Each reading of the burette has an uncertainty of  $\pm 0.05 \text{ cm}^3$ ]

- A  $\pm 0.38\%$
- B  $\pm 0.75\%$
- C  $\pm 1.5\%$
- D  $\pm 7.5\%$

(Total for Question 15 = 1 mark)

16 Barium hydroxide reacts with sulfuric acid as shown.



Which is the ionic equation for this reaction?

- A  $\text{Ba}^{2+}(\text{aq}) + 2\text{OH}^-(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{BaSO}_4(\text{s}) + 2\text{H}_2\text{O}(\text{l})$
- B  $\text{Ba}^{2+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{BaSO}_4(\text{s})$
- C  $\text{OH}^-(\text{aq}) + \text{H}^+(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l})$
- D  $\text{Ba}^{2+}(\text{aq}) + 2\text{OH}^-(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{SO}_4^{2-}(\text{aq}) \rightarrow \text{Ba}^{2+}(\text{s}) + \text{SO}_4^{2-}(\text{s}) + 2\text{H}_2\text{O}(\text{l})$

(Total for Question 16 = 1 mark)



P 6 9 5 0 1 A 0 7 2 8

**17** Four tests used to identify ions are shown:


- 1 flame test
- 2 addition of acidified barium nitrate solution
- 3 addition of acidified silver nitrate solution
- 4 addition of sodium hydroxide solution, then testing any gas with indicator paper

Which tests could be used to positively identify the ions in ammonium chloride?

- A** 1 and 2
- B** 1 and 3
- C** 2 and 4
- D** 3 and 4

(Total for Question 17 = 1 mark)

**18** Which conditions give the highest yield for the forward reaction?




- A** high temperature, high pressure
- B** high temperature, low pressure
- C** low temperature, high pressure
- D** low temperature, low pressure

(Total for Question 18 = 1 mark)

**Use this space for any rough working. Anything you write in this space will gain no credit.**



**19** Nitrogen dioxide and dinitrogen tetroxide exist in equilibrium.



When an equilibrium is set up in a gas syringe, the mixture is pale brown.

When the mixture is compressed the colour becomes

- A** darker
- B** lighter
- C** darker and then lighter
- D** lighter and then darker

(Total for Question 19 = 1 mark)

---

**TOTAL FOR SECTION A = 20 MARKS**



P 6 9 5 0 1 A 0 9 2 8

**SECTION B**


**Answer ALL the questions. Write your answers in the spaces provided.**

**20** Ammonia reacts with 1-chlorobutane.

(a) State the type and mechanism of this reaction.

(2)

(b) A student drew the first step of the mechanism for the reaction.




(i) Identify **two** omissions in the student's mechanism.

(2)

(ii) To obtain butylamine, sodium hydroxide solution is added.

Complete the next step of the mechanism to form butylamine,  
showing curly arrows, relevant lone pairs and the reaction products.

(3)



(c) The reactions of ammonia and of hydroxide ions with halogenoalkanes are similar.

Compare the rate of reaction of ammonia with 1-chlorobutane and with 2-bromo-2-methylpropane.


Justify your answer.

(3)

**(Total for Question 20 = 10 marks)**



- 21 Enthalpy changes of combustion can be determined using calorimetry or calculated using Hess cycles. Apparatus for a calorimetry experiment is shown.



A sample of 2-methylpropan-2-ol was burned in a spirit burner and used to heat 75 g of water. The results are shown.

	At the start	At the end	Change
Mass of spirit burner / g	267.35	266.78	
Temperature of water / °C	19.5	65.3	

- (a) (i) Complete the table.

(1)

- (ii) Calculate the enthalpy change of combustion,  $\Delta_c H$ , of 2-methylpropan-2-ol.

Give a sign and units in your answer.

[Specific heat capacity of water =  $4.18 \text{ J g}^{-1} \text{ }^{\circ}\text{C}^{-1}$ ]

(4)




- (b) The standard enthalpy change of combustion,  $\Delta_c H^\ominus$ , can be calculated using standard enthalpy changes of formation.

Compound	$\Delta_f H^\ominus / \text{kJ mol}^{-1}$
2-methylpropan-2-ol	-359
carbon dioxide	-394
water	-286

- (i) State why no  $\Delta_f H^\ominus$  value has been given for oxygen.

(1)

- (ii) Complete the Hess cycle.



(2)

- (iii) Calculate the standard enthalpy change of combustion of 2-methylpropan-2-ol using the data in the table and the completed Hess cycle.

(2)



- (c) The value for  $\Delta_c H$  obtained in part (a)(ii) is much less exothermic than  $\Delta_c H^\ominus$  calculated in (b)(iii).

Suggest **two** reasons for this other than non-standard conditions.

(2)

---

---

---

---

---

**(Total for Question 21 = 12 marks)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



**22** This question is about the elements in Group 7.

- (a) Use your knowledge of the trends in the properties of Group 7 elements to predict the colour and physical state of astatine at room temperature.

(1)

.....  
.....

- (b) (i) State the meaning of the term electronegativity.

(1)

.....  
.....

- (ii) Explain the trend in electronegativity down Group 7.

(2)

.....  
.....  
.....  
.....  
.....



**\*c) Compare and contrast the reactions of chlorine with**

- water
  - cold, dilute aqueous alkali
  - hot, concentrated aqueous alkali

Include an equation for each reaction, stating the type of reaction and the oxidation numbers of the chlorine involved. State symbols are not required.

(6)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 22 = 10 marks)



P 6 9 5 0 1 A 0 1 7 2 8

- 23** Magnesium ethanedioate ( $\text{MgC}_2\text{O}_4$ ) decomposes on gentle heating to form magnesium carbonate and carbon monoxide.



- (a) (i) State why the thermal decomposition of magnesium ethanedioate should be carried out in a fume cupboard.

(1)

---

---

- (ii) After heating a 6.0 g sample of magnesium ethanedioate for three minutes, the decomposition was 70% complete.

Calculate the total mass of the solid mixture that remains.

(4)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



- (b) Magnesium carbonate undergoes thermal decomposition at a higher temperature than magnesium ethanedioate.



Explain the trend in the thermal decomposition of Group 2 carbonates going down the group.

(3)

**(Total for Question 23 = 8 marks)**

**TOTAL FOR SECTION B = 40 MARKS**




P 6 9 5 0 1 A 0 1 9 2 8

**SECTION C**

**Answer all the questions. Write your answers in the spaces provided.**

- 24** Some diesel cars contain an extra catalytic converter for the reduction of nitrogen oxides ( $\text{NO}_x$ ) in exhaust gases.  
A solution of urea is used for this process.



- (a) Urea has a melting temperature of  $133^\circ\text{C}$ .

Explain why this value is higher than expected for a relatively small molecule.

(3)

---

---

---

---

---

- (b) A saturated solution of urea has a concentration of  $9.07 \text{ mol dm}^{-3}$  at  $25^\circ\text{C}$ .

Calculate the mass of urea in  $150 \text{ cm}^3$  of a saturated solution.

(2)




- (c) State why  $\text{NO}_x$  emissions are harmful to the environment.

(1)

- (d) An infrared spectrum of urea is shown.

Refer to your Data Booklet.



- (i) Draw a circle around an absorption in the spectrum that could be due to the stretching of the N—H bond.

(1)

- (ii) Identify the bond responsible for the absorption at  $1683\text{ cm}^{-1}$ .

(1)

- (e) In a diesel car exhaust system, the urea reacts with water to form ammonia and carbon dioxide. The enthalpy change for this reaction is  $+133\text{ kJ mol}^{-1}$ .

- (i) Complete the equation for this **reversible** reaction.

State symbols are not required.

(1)



P 6 9 5 0 1 A 0 2 1 2 8

- (ii) Sketch the reaction profile for the forward reaction on the axes provided.

Include labels for  $\Delta H$  and the activation energy ( $E_a$ ).



(3)

- (f) The catalytic converter contains metal oxides. When the exhaust gases pass through the catalytic converter, ammonia reacts with  $\text{NO}_x$  gases to form nitrogen and water.
- (i) Explain why it is **not** correct to state that urea is acting as a catalyst in the reaction.


(1)



- (ii) Explain how a catalyst increases the rate of a chemical reaction.

Use the Maxwell-Boltzmann distribution shown and refer to the collision theory.

(3)



- (g) The catalytic converter works best at a temperature of around 350 °C.

- (i) Suggest how the catalytic converter reaches this temperature.

(1)



- (ii) The chemical reactions in the exhaust system of a diesel car, using a catalytic converter, form  $89.3 \text{ m}^3$  of nitrogen per hour.

Calculate the number of molecules of nitrogen formed per hour.

[Molar volume at  $350^\circ\text{C}$  =  $51.1 \text{ dm}^3 \text{ mol}^{-1}$

Avogadro constant,  $L = 6.02 \times 10^{23} \text{ mol}^{-1}$ ]

(3)

**(Total for Question 24 = 20 marks)**

**TOTAL FOR SECTION C = 20 MARKS**

**TOTAL FOR PAPER = 80 MARKS**




**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**BLANK PAGE**



P 6 9 5 0 1 A 0 2 5 2 8

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**



**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**DO NOT WRITE IN THIS AREA**

**BLANK PAGE**



P 6 9 5 0 1 A 0 2 7 2 8

# The Periodic Table of Elements

1 2

(1)	(2)
6.9 Li lithium 3	9.0 Be beryllium 4
23.0 Na sodium 11	24.3 Mg magnesium 12

## Key

relative atomic mass
atomic symbol
name
atomic (proton) number

1.0 H hydrogen 1
------------------

3 4 5 6 7 0 (8)

10.8 B boron 5	12.0 C carbon 6	14.0 N nitrogen 7	16.0 O oxygen 8	19.0 F fluorine 9	20.2 Ne neon 10
27.0 Al aluminium 13	28.1 Si silicon 14	31.0 P phosphorus 15	32.1 Cl chlorine 17	35.5 Ar argon 18	
39.9 Ge germanium 32	40.8 As arsenic 33	41.9 Se selenium 34	42.9 Br bromine 35	43.8 Kr krypton 36	
50.7 Ga gallium 31	52.6 Ge germanium 32	53.5 Cu copper 29	54.4 Zn zinc 30	55.4 Ga gallium 31	
54.9 Mn manganese 25	55.8 Fe iron 26	56.7 Ni nickel 28	57.5 Cu copper 29	58.7 Ni nickel 28	
56.9 Cr chromium 24	57.0 Ti titanium 23	58.9 Co cobalt 27	59.7 Fe iron 26	60.6 Co cobalt 27	
58.9 Sc scandium 21	59.9 V vanadium 22	60.8 Mn manganese 25	61.7 Fe iron 26	62.6 Sc scandium 21	
60.0 Ca calcium 20	61.0 Ti titanium 22	61.9 Cr chromium 23	62.8 V vanadium 23	63.0 Ca calcium 20	
61.9 K potassium 19	62.9 Y yttrium 39	63.9 Nb niobium 41	64.8 Tc technetium 43	65.9 K potassium 19	
62.9 Rb rubidium 37	63.9 Sr strontium 38	64.9 Zr zirconium 40	65.9 Ru ruthenium 44	66.9 Rb rubidium 37	
63.9 Cs caesium 55	63.9 Ba barium 56	64.9 La lanthanum 57	65.9 Hf hafnium 72	66.9 Cs caesium 55	
64.9 Fr francium 87	65.9 Ra radium 88	65.9 Ac actinium 89	66.9 Rf rutherfordium 104	67.9 Db dubnium 105	
65.9 Ce cerium 58	66.9 Pr praseodymium 59	67.9 Nd neodymium 60	68.9 Pm promethium 61	69.9 Sm samarium 62	
66.9 Tb terbium 65	67.9 Dy dysprosium 66	68.9 Gd gadolinium 64	69.9 Eu europium 63	70.9 Ho holmium 67	
67.9 Ds darmstadtium 109	68.9 Mt meitnerium 108	68.9 Hs hassium 107	69.9 Ts tsenium 108	70.9 Rg roentgenium 110	
68.9 Rg roentgenium 111					

Elements with atomic numbers 112-116 have been reported  
but not fully authenticated

71.5 Lu lutetium 71	72.5 Hg mercury 80	73.5 Tl thallium 81	74.5 Pb lead 82	75.5 Bi bismuth 83	76.5 Po polonium 84	77.5 At astatine 85	78.5 Rn radon 86
79.5 Te tellurium 52	80.5 Sn tin 50	81.5 In indium 49	82.5 Cd cadmium 48	83.5 Sb antimony 51	84.5 Te tellurium 52	85.5 I iodine 53	86.5 Xe xenon 54
85.5 Kr krypton 36	86.5 Ge germanium 34	87.5 Ar argon 18	88.5 Zn zinc 30	89.5 Ga gallium 31	90.5 Ge germanium 32	91.5 F fluorine 9	92.5 He helium 2
93.5 Rb radon 86	94.5 Po polonium 84	95.5 At astatine 85	96.5 Br bromine 35	97.5 Cl chlorine 17	98.5 S sulfur 16	99.5 F fluorine 9	100.5 Ne neon 10

\* Lanthanide series

\* Actinide series

