

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Centre Number

Candidate Number

--	--	--	--	--

--	--	--	--

Pearson Edexcel International Advanced Level

Time 1 hour 20 minutes

Paper
reference

WCH16/01

Chemistry

International Advanced Level

UNIT 6: Practical Skills in Chemistry II

You must have:

Scientific calculator, ruler

Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P70957A

©2022 Pearson Education Ltd.


Q:1/1/1/




Pearson

Answer ALL the questions. Write your answers in the spaces provided.

- 1 A flowchart of a series of reactions of chromium and some of its compounds is shown.



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(a) Give the formula of a complex ion of chromium present in solution **A**. (1)

(b) Give the formula of precipitate **B**. (1)

(c) Identify reagent **C**, by name or formula. (1)

(d) Suggest the identity, by name or formula, of precipitate **D**. (1)

(e) State the colour of solution **E**. (1)

(f) State the colour of solution **F**. (1)

(Total for Question 1 = 6 marks)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



2 This question is about the identification of three organic compounds, **X**, **Y** and **Z**.

A molecule of each compound has only one **type** of functional group.
Each compound contains six carbon atoms and two oxygen atoms.

In the mass spectrum of **X**, the molecular ion peak is at $m/z = 114$.

(a) State the molecular formula of **X**.

(1)

(b) Three chemical tests are carried out on **X**.

Test 1 When Brady's reagent (2,4-dinitrophenylhydrazine solution) is added to **X**,
an orange precipitate is observed.

Test 2 When **X** is heated with an acidified solution of potassium dichromate(VI),
no change is observed.

Test 3 When **X** is added to an alkaline solution of iodine, the formation of a
pale yellow precipitate is observed.

Explain what can be deduced about the functional group present in **X**, by
considering the results of each of these tests.

(3)



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

- (c) The **high** resolution proton NMR spectrum of **X** contains only two singlets with relative peak areas of 3:2.

Draw the structure of **X**, identifying the two proton environments.

(2)



(d) The molecular formula of **Y** is $C_6H_{12}O_2$.

(i) When aqueous sodium carbonate is added to **Y**, effervescence is observed.

Identify, by name or formula, the functional group present in **Y**.

(1)

(ii) The ^{13}C NMR spectrum of **Y** contains **four** peaks.

Give **two** possible structures for **Y**.

(2)

Structure 1	Structure 2

(iii) Explain how the **low** resolution **proton** NMR spectrum of **Y** would confirm which of your structures in (d)(ii) is correct.

Chemical shifts are not required.

(2)

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(e) The molecular formula of **Z** is $C_6H_{12}O_2$.

The infrared spectrum of **Z** is shown.



Group	Wavenumber range / cm^{-1}
C—H stretching vibrations	
Alkane	2962–2853
Alkene	3095–3010
Aldehyde	2900–2820 and 2775–2700
O—H stretching vibrations	
Alcohols	3750–3200
Carboxylic acids	3300–2500
C=O stretching vibrations	
Aldehydes	1740–1720
Ketones	1720–1700
Carboxylic acids	1725–1700
Esters	1750–1735



- (i) **Z** has a fruity smell.

Deduce the identity of the functional group present in **Z**, using all the information given.

Include any relevant wavenumber ranges in your answer.

(2)

- (ii) The high resolution proton NMR spectrum of **Z** contains four peaks (J, K, L and M). Peak J has the highest chemical shift, showing that this proton environment is close to an electronegative atom.

The splitting pattern of the peaks is shown.

Peak	J	K	L	M
Splitting pattern	septet	quartet	doublet	triplet

Draw the displayed structure of **Z**, labelling the proton environment responsible for each of the peaks J, K, L and M.

(3)

(Total for Question 2 = 16 marks)

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 7 0 9 5 7 A 0 9 2 0

- 3 A student investigated the kinetics of the reaction between bromide ions, Br^- , and bromate(V) ions, BrO_3^- , in aqueous acid.



To determine the rate equation for the reaction, the student varied the concentration of bromide ions, bromate(V) ions and acid in turn.

The effect of the concentration of bromide ions was investigated first.

Procedure

- Step 1** Add 10.0 cm^3 of $0.0050\text{ mol dm}^{-3}$ potassium bromate(V), 15.0 cm^3 of acidified methyl orange indicator solution and 5.0 cm^3 of $0.00010\text{ mol dm}^{-3}$ aqueous phenol to a beaker labelled **P**.
- Step 2** Prepare the contents of beaker **Q** for Run 1 as specified in the table.
- Step 3** Pour the contents of beaker **Q** into beaker **P** and start a timer. Pour the contents of beaker **P** back into beaker **Q** and place beaker **Q** on a white tile.
- Step 4** Stop the timer as soon as the mixture turns colourless. Record the time, along with the temperature of the solution.
- Step 5** Repeat Steps **1** to **4** for the remaining runs.

Contents of beaker Q		
Run	Volume of 0.01 mol dm^{-3} KBr / cm^3	Volume of $\text{H}_2\text{O} / \text{cm}^3$
1	10.0	0
2	8.0	2.0
3	6.0	4.0
4	5.0	5.0
5	4.0	6.0
6	3.0	7.0



(a) Give **one** reason why the contents of beaker **P** are poured back into beaker **Q** in Step **3**.

(1)

(b) Methyl orange indicator is bleached colourless by bromine.
Under the experimental conditions, the reciprocal of the time taken for the methyl orange to be bleached ($1/t$) is proportional to the initial rate.

(i) Explain why phenol is added to the reaction mixture.

(2)

(ii) Give the colour of the reaction mixture before it turns colourless.

(1)

(iii) Give the reason why the temperature is measured in Step **4**.

(1)




P 7 0 9 5 7 A 0 1 1 2 0

(c) The student's results are shown.

Volume of KBr / cm ³	10.0	8.0	6.0	5.0	4.0	3.0
Time, t / s	23	24	32	39	48	64
$(1/t)$ / s ⁻¹	0.043	0.042	0.031	0.026		
Temperature / °C	18	22	22	22	22	22

(i) Complete the table by calculating the remaining values of $1/t$. (1)

(ii) Plot a graph of $1/t$ against the volume of KBr. Include a line of best fit. (3)





(iii) State why the volume of KBr is proportional to the concentration of bromide ions in the reaction mixture.

(1)

(iv) State the order of reaction with respect to bromide ions, using your graph.

(1)

(d) After investigating the effect on the rate of the concentration of bromate(V) ions and the concentration of hydrogen ions, the student obtained the graphs shown.



Deduce the rate equation for the reaction, using these data and your answer to (c)(iv).


(1)

(Total for Question 3 = 12 marks)



P 7 0 9 5 7 A 0 1 3 2 0

- 4 This question is about the preparation of phenylamine by the reduction of nitrobenzene.



Outline procedure

- Step 1** Add 2.1 cm³ of nitrobenzene, 5 g of granulated tin and 10 cm³ of concentrated hydrochloric acid to a round-bottomed flask.
- Step 2** Add a few anti-bumping granules to the flask and heat the contents under reflux for 15 minutes. Leave to cool.
- Step 3** Dissolve 7.5 g of sodium hydroxide in 10 cm³ of distilled water and add to the flask. An initial precipitate forms before redissolving.
- Step 4** Add 15 cm³ of distilled water to the flask and steam distil the contents, collecting the cloudy distillate in a conical flask.
- Step 5** Add 3 g of powdered sodium chloride to the distillate and swirl to dissolve, before transferring the contents to a separating funnel. Add 8 cm³ of ether to the funnel and shake, occasionally relieving the pressure. Allow the layers to separate.
- Step 6** Discard the aqueous layer before transferring the ether layer to a clean flask containing a few pellets of potassium hydroxide.
- Step 7** Decant the contents of the flask from Step 6 into a pear-shaped flask and distil off all the ether.
- Step 8** Distil the remaining contents of the pear-shaped flask, collecting the fraction boiling between 180 °C and 185 °C.

Some data relating to the organic chemicals involved in the preparation are shown.

Chemical	Hazard	M_r	Density / g cm ⁻³	Boiling temperature / °C
Nitrobenzene	Toxic by inhalation and skin absorption	123.0	1.20	211
Phenylamine	Toxic by inhalation and skin absorption	93.0	1.03	184
Ether	Highly flammable	74.0	0.71	35



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(a) Give **two** reasons why gloves should be worn in Step 1.

(2)

.....

.....

.....

(b) (i) State the role of tin in the preparation.

(1)

.....

.....

(ii) Suggest, by name or formula, the identity of the initial precipitate formed in Step 3.

(1)

.....

.....

(c) State why the distillate in Step 4 is cloudy.

(1)

.....

.....

(d) Suggest why sodium chloride is added to the distillate in Step 5.

(1)

.....

.....

.....



(e) (i) Draw a **labelled** diagram of the separating funnel **at the end** of Step 5.

(2)

(ii) State how you would relieve the pressure in the separating funnel in Step 5.

(1)

.....

.....

.....

(f) Suggest **one** reason for adding pellets of potassium hydroxide in Step 6.

(1)

.....

.....



(g) State how the mixture would be heated to distil off the ether in Step 7.
Justify your answer.

(2)

.....

.....

.....

(h) Calculate the mass of phenylamine formed in this preparation.

The limiting reagent is nitrobenzene and the overall yield by mass is 43 %.

Refer to all the information at the start of the question.

(4)

(Total for Question 4 = 16 marks)

TOTAL FOR PAPER = 50 MARKS

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

BLANK PAGE



The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0
H
hydrogen
1

Key

relative atomic mass
atomic symbol
name
atomic (proton) number

(1)	6.9 Li lithium 3	(2)	9.0 Be beryllium 4	(3)	45.0 Sc scandium 21	(4)	47.9 Ti titanium 22	(5)	50.9 V vanadium 23	(6)	52.0 Cr chromium 24	(7)	54.9 Mn manganese 25	(8)	55.8 Fe iron 26	(9)	58.9 Co cobalt 27	(10)	58.7 Ni nickel 28	(11)	63.5 Cu copper 29	(12)	65.4 Zn zinc 30	(13)	10.8 B boron 5	(14)	12.0 C carbon 6	(15)	14.0 N nitrogen 7	(16)	16.0 O oxygen 8	(17)	19.0 F fluorine 9	(18)	4.0 He helium 2	
	23.0 Na sodium 11		24.3 Mg magnesium 12		40.1 Ca calcium 20		47.9 Ti titanium 22		50.9 V vanadium 23		52.0 Cr chromium 24		54.9 Mn manganese 25		55.8 Fe iron 26		58.9 Co cobalt 27		58.7 Ni nickel 28		63.5 Cu copper 29		65.4 Zn zinc 30		27.0 Al aluminium 13		28.1 Si silicon 14		31.0 P phosphorus 15		32.1 S sulfur 16		35.5 Cl chlorine 17		39.9 Ar argon 18	
	39.1 K potassium 19		39.1 K potassium 19		39.1 K potassium 19		47.9 Ti titanium 22		50.9 V vanadium 23		52.0 Cr chromium 24		54.9 Mn manganese 25		55.8 Fe iron 26		58.9 Co cobalt 27		58.7 Ni nickel 28		63.5 Cu copper 29		65.4 Zn zinc 30		69.7 Ga gallium 31		72.6 Ge germanium 32		74.9 As arsenic 33		79.0 Se selenium 34		79.9 Br bromine 35		83.8 Kr krypton 36	
	85.5 Rb rubidium 37		87.6 Sr strontium 38		88.9 Y yttrium 39		91.2 Zr zirconium 40		92.9 Nb niobium 41		95.9 Mo molybdenum 42		[98] Tc technetium 43		101.1 Ru ruthenium 44		102.9 Rh rhodium 45		106.4 Pd palladium 46		107.9 Ag silver 47		112.4 Cd cadmium 48		114.8 In indium 49		118.7 Sn tin 50		121.8 Sb antimony 51		127.6 Te tellurium 52		126.9 I iodine 53		131.3 Xe xenon 54	
	132.9 Cs caesium 55		137.3 Ba barium 56		138.9 La* lanthanum 57		178.5 Hf hafnium 72		180.9 Ta tantalum 73		183.8 W tungsten 74		186.2 Re rhenium 75		190.2 Os osmium 76		192.2 Ir iridium 77		195.1 Pt platinum 78		197.0 Au gold 79		200.6 Hg mercury 80		204.4 Tl thallium 81		207.2 Pb lead 82		209.0 Bi bismuth 83		[209] Po polonium 84		[210] At astatine 85		[222] Rn radon 86	
	[223] Fr francium 87		[226] Ra radium 88		[227] Ac* actinium 89		[261] Rf rutherfordium 104		[262] Db dubnium 105		[266] Sg seaborgium 106		[264] Bh bohrium 107		[277] Hs hassium 108		[268] Mt meitnerium 109		[271] Ds darmstadtium 110		[272] Rg roentgenium 111															

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140	Ce cerium 58	141	Pr praseodymium 59	144	Nd neodymium 60	150	Sm samarium 62	152	Eu europium 63	157	Gd gadolinium 64	163	Dy dysprosium 66	165	Ho holmium 67	167	Er erbium 68	169	Tm thulium 69	173	Yb ytterbium 70	175	Lu lutetium 71
232	Th thorium 90	[231]	Pa protactinium 91	238	U uranium 92	[242]	Pu plutonium 94	[243]	Am americium 95	[247]	Cm curium 96	[251]	Cf californium 98	[254]	Es einsteinium 99	[253]	Fm fermium 100	[256]	Md mendelevium 101	[254]	No nobelium 102	[257]	Lr lawrencium 103

* Lanthanide series
* Actinide series



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA