


Question number	Scheme	Marks
1.	$x^2 - 9 = (x - 3)(x + 3)$ seen Attempt at forming single fraction $\frac{x(x - 3) + (x + 12)(x + 1)}{(x + 1)(x + 3)(x - 3)} ; = \frac{2x^2 + 10x + 12}{(x + 1)(x + 3)(x - 3)}$ Factorising numerator = $\frac{2(x + 2)(x + 3)}{(x + 1)(x + 3)(x - 3)}$ or equivalent = $\frac{2(x + 2)}{(x + 1)(x - 3)}$	B1 M1; A1 M1 M1 A1 (6) (6 marks)
2.	$(1 + px)^n \equiv 1 + npx, + \frac{n(n - 1)p^2 x^2}{2} + \dots$ Comparing coefficients: $np = -18$, $\frac{n(n - 1)}{2} = 36$ Solving $n(n - 1) = 72$ to give $n = 9$; $p = -2$	B1, B1 M1, A1 M1 A1; A1 ft (7) (7 marks)
3. (a)	<p>V graph with 'vertex' on x-axis $\{-\frac{1}{2}a, (0)\}$ and $\{(0), a\}$ seen</p>	M1 A1 (2)
	<p>Correct graph (could be separate)</p>	B1 (1)
(c)	Meet where $\frac{1}{x} = 2x + a \Rightarrow x 2x + a - 1 = 0$; only one meet	B1 (1)
(d)	$2x^2 + x - 1$ Attempt to solve; $x = \frac{1}{2}$ (no other value)	B1 M1; A1 (3) (7 marks)

Question number	Scheme	Marks														
4.	$\text{Volume} = \pi \int_1^4 \left(1 + \frac{1}{2\sqrt{x}}\right)^2 dx$ $\int \left(1 + \frac{1}{2\sqrt{x}}\right)^2 dx = \int \left(1 + \frac{1}{\sqrt{x}} + \frac{1}{4x}\right) dx$ $= \left[x + 2\sqrt{x} + \frac{1}{4} \ln x \right]$ <p>Using limits correctly</p> $\text{Volume} = \pi \left[\left(8 + \frac{1}{4} \ln 4\right) + 3 \right]$ $= \pi \left[5 + \frac{1}{2} \ln 2 \right]$	M1 B1 M1 A1 A1ft M1 A1 A1 (8) (8 marks)														
5. (a)	<table border="1"> <tr> <td>Distance from one side (m)</td> <td>0</td> <td>2</td> <td>4</td> <td>6</td> <td>8</td> <td>10</td> </tr> <tr> <td>Height (m)</td> <td>0</td> <td>6.13</td> <td>7.80</td> <td>7.80</td> <td>6.13</td> <td>0</td> </tr> </table> <p>"y" = 7.80 when "x" = 4 or 6</p> <p>Symmetry</p>	Distance from one side (m)	0	2	4	6	8	10	Height (m)	0	6.13	7.80	7.80	6.13	0	B1 B1 ft (2)
Distance from one side (m)	0	2	4	6	8	10										
Height (m)	0	6.13	7.80	7.80	6.13	0										
(b)	Estimate area = $\frac{2}{2} [0 + 2(6.13 + 7.80 + 7.80 + 6.13)]$ $= 55.7 \text{ m}^2$	B1 M1 A1ft A1 (4)														
(c)	$140 - (b) = 84.3 \text{ m}^2$	A1 ft (1)														
(d)	Over-estimate; reason, e.g. area under curve is under-estimate (due to curvature)	B1 B1 (2) (9 marks)														

Question number	Scheme	Marks
6. (a)	 <p>Shape $p = \frac{1}{3}$ or $\{\frac{1}{3}, 0\}$ seen</p>	B1 B1 (2)
(b)	<p>Gradient of tangent at $Q = \frac{1}{q}$</p> <p>Gradient of normal = $-q$</p> <p>Attempt at equation of OQ [$y = -qx$] and substituting $x = q$, $y = \ln 3q$</p> <p>or attempt at equation of tangent [$y - 3 \ln q = -q(x - q)$] with $x = 0$, $y = 0$</p> <p>or equating gradient of normal to $(\ln 3q)/q$</p> <p>$q^2 + \ln 3q = 0$ (*)</p>	B1 M1 M1 M1 (4)
(c)	$\ln 3x = -x^2 \Rightarrow 3x = e^{-x^2} ; \Rightarrow x = \frac{1}{3}e^{-x^2}$	M1; A1 (2)
(d)	$x_1 = 0.298280; x_2 = 0.304957, x_3 = 0.303731, x_4 = 0.303958$ Root = 0.304 (3 decimal places)	M1; A1 A1 (3)
		(11 marks)
7. (a)	$\sin x + \sqrt{3} \cos x = R \sin(x + \alpha)$ $= R(\sin x \cos \alpha + \cos x \sin \alpha)$ $R \cos \alpha = 1, R \sin \alpha = \sqrt{3}$ Method for R or α , e.g. $R = \sqrt{(1+3)}$ or $\tan \alpha = \sqrt{3}$ Both $R = 2$ and $\alpha = 60^\circ$	M1 A1 M1 A1 (4)
(b)	$\sec x + \sqrt{3} \operatorname{cosec} x = 4 \Rightarrow \frac{1}{\cos x} + \frac{\sqrt{3}}{\sin x} = 4$ $\Rightarrow \sin x + \sqrt{3} \cos x = 4 \sin x \cos x$ $= 2 \sin 2x$ (*)	B1 M1 M1 (3)
(c)	Clearly producing $2 \sin 2x = 2 \sin(x + 60^\circ)$	A1 (1)
(d)	$\sin 2x - \sin(x + 60^\circ) = 0 \Rightarrow \cos \frac{3x + 60}{2} \sin \frac{x - 60}{2} = 0$ $\cos \frac{3x + 60}{2} = 0 \Rightarrow x = 40^\circ, 160^\circ$ $\sin \frac{x - 60}{2} = 0 \Rightarrow x = 60^\circ$	M1 M1 A1 A1 ft B1 (5)
		(13 marks)

Question number	Scheme	Marks
8. (a)	<p>shape intersections with axes $(c, 0), (0, d)$</p>	B1 B1 (2)
(b)	<p>shape x intersection $(\frac{1}{2}d, 0)$ y intersection $(0, 3c)$</p>	B1 B1 B1 (3)
(c)(i)	$c = 2$	B1
(ii)	$-1 < f(x) \leq$ (candidate's) c value	B1 B1 ft (3)
(d)	$3(2^{-x}) = 1 \Rightarrow 2^{-x} = \frac{1}{3}$ and take logs; $-x = \frac{\ln \frac{1}{3}}{\ln 2}$ d (or x) = 1.585 (3 decimal places)	M1; A1 A1 (3)
(e)	$fg(x) = f[\log_2 x] = [3(2^{-\log_2 x}) - 1]; = [3(2^{\log_2 \frac{1}{x}}) - 1]$ or $\frac{3}{2^{\log_2 x}} - 1$ $= \frac{3}{x} - 1$	M1; A1 A1 (3)
		(14 marks)