















4.




Figure 1

A set square  $S$  is made by removing a circle of centre  $O$  and radius 3 cm from a triangular piece of wood. The piece of wood is modelled as a uniform triangular lamina  $ABC$ , with  $\angle ABC = 90^\circ$ ,  $AB = 12$  cm and  $BC = 21$  cm. The point  $O$  is 5 cm from  $AB$  and 5 cm from  $BC$ , as shown in Figure 1.

(a) Find the distance of the centre of mass of  $S$  from

- (i)  $AB$ ,
- (ii)  $BC$ .

(9)

The set square is freely suspended from  $C$  and hangs in equilibrium.

(b) Find, to the nearest degree, the angle between  $CB$  and the vertical.

(3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---












5.



**Figure 2**

A ladder  $AB$ , of mass  $m$  and length  $4a$ , has one end  $A$  resting on rough horizontal ground. The other end  $B$  rests against a smooth vertical wall. A load of mass  $3m$  is fixed on the ladder at the point  $C$ , where  $AC = a$ . The ladder is modelled as a uniform rod in a vertical plane perpendicular to the wall and the load is modelled as a particle. The ladder rests in limiting equilibrium making an angle of  $30^\circ$  with the wall, as shown in Figure 2.

Find the coefficient of friction between the ladder and the ground.

**(10)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









6.




Figure 3

[In this question, the unit vectors  $\mathbf{i}$  and  $\mathbf{j}$  are in a vertical plane,  $\mathbf{i}$  being horizontal and  $\mathbf{j}$  being vertical.]

A particle  $P$  is projected from the point  $A$  which has position vector  $47.5\mathbf{j}$  metres with respect to a fixed origin  $O$ . The velocity of projection of  $P$  is  $(2u\mathbf{i} + 5u\mathbf{j}) \text{ m s}^{-1}$ . The particle moves freely under gravity passing through the point  $B$  with position vector  $30\mathbf{i}$  metres, as shown in Figure 3.

- (a) Show that the time taken for  $P$  to move from  $A$  to  $B$  is 5 s. (6)
- (b) Find the value of  $u$ . (2)
- (c) Find the speed of  $P$  at  $B$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---













Leave blank

7. A particle  $P$  of mass  $2m$  is moving with speed  $2u$  in a straight line on a smooth horizontal plane. A particle  $Q$  of mass  $3m$  is moving with speed  $u$  in the same direction as  $P$ . The particles collide directly. The coefficient of restitution between  $P$  and  $Q$  is  $\frac{1}{2}$ .

(a) Show that the speed of  $Q$  immediately after the collision is  $\frac{8}{5}u$ . (5)

(b) Find the total kinetic energy lost in the collision. (5)

After the collision between  $P$  and  $Q$ , the particle  $Q$  collides directly with a particle  $R$  of mass  $m$  which is at rest on the plane. The coefficient of restitution between  $Q$  and  $R$  is  $e$ .

(c) Calculate the range of values of  $e$  for which there will be a second collision between  $P$  and  $Q$ . (7)

Lined area for writing the answer to part (c).









**BLANK PAGE**

