Centre No.						Paper Reference							Surname	Initial(s)
Candidate No.						6	6	7	9	/	0	1	Signature	

Paper Reference(s)

6679/01

Edexcel GCE

Mechanics M3

Advanced/Advanced Subsidiary

Thursday 29 January 2009 – Morning

Time: 1 hour 30 minutes

Examiner's use only

Team Leader's use only

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
Total	

Materials required for examination
Mathematical Formulae (Green)

Items included with question papers
Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature.
Check that you have the correct question paper.
Answer ALL the questions.
You must write your answer for each question in the space following the question.
Whenever a numerical value of g is required, take $g = 9.8$ m s^{-2}.
When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.
Full marks may be obtained for answers to ALL questions.
The marks for individual questions and the parts of questions are shown in round brackets: e.g. **(2)**.
There are 7 questions in this question paper. The total mark for this question paper is 75.
There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled.
You should show sufficient working to make your methods clear to the Examiner.
Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.
©2009 Edexcel Limited.

Printer's Log. No.
N29495A
W850/R6679/57570 3/3/3/3/3

N 2 9 4 9 5 A 0 1 2 4

Turn over

edexcel
advancing learning, changing lives

https://xtremepape.rs/

Leave blank

1. A particle P of mass 3 kg is moving in a straight line. At time t seconds, $0 \leqslant t \leqslant 4$, the only force acting on P is a resistance to motion of magnitude $\left(9+\frac{15}{(t+1)^2}\right)$ N. At time t seconds the velocity of P is v m s^{-1}. When $t = 4$, $v = 0$.

Find the value of v when $t = 0$.

(7)

N 2 9 4 9 5 A 0 2 2 4

https://xtremepape.rs/

Leave blank

Question 1 continued

Q1

(Total 7 marks)

N 2 9 4 9 5 A 0 3 2 4

Turn over

https://xtremepape.rs/

Leave blank

2.



Figure 1

A particle P of mass m is attached to one end of a light elastic string, of natural length a and modulus of elasticity $3mg$. The other end of the string is attached to a fixed point O.

The particle P is held in equilibrium by a horizontal force of magnitude $\frac{4}{3}mg$ applied to P.

This force acts in the vertical plane containing the string, as shown in Figure 1. Find

(a) the tension in the string,

(5)

(b) the elastic energy stored in the string.

(4)

N 2 9 4 9 5 A 0 4 2 4

https://xtremepape.rs/

Leave blank

Question 2 continued

Q2

(Total 9 marks)

N 2 9 4 9 5 A 0 5 2 4

Turn over

https://xtremepape.rs/

Leave blank

3. A rough disc rotates about its centre in a horizontal plane with constant angular speed 80 revolutions per minute. A particle P lies on the disc at a distance 8 cm from the centre of the disc. The coefficient of friction between P and the disc is μ. Given that P remains at rest relative to the disc, find the least possible value of μ.

(7)

N 2 9 4 9 5 A 0 6 2 4

https://xtremepape.rs/

Leave blank

Question 3 continued

(Total 7 marks)

Q3

N 2 9 4 9 5 A 0 7 2 4

Turn over

https://xtremepape.rs/

Leave blank

4. A small shellfish is attached to a wall in a harbour. The rise and fall of the water level is modelled as simple harmonic motion and the shellfish as a particle. On a particular day the minimum depth of water occurs at 10 00 hours and the next time that this minimum depth occurs is at 22 30 hours. The shellfish is fixed in a position 5 m above the level of the minimum depth of the water and 11 m below the level of the maximum depth of the water. Find

(a) the speed, in metres per hour, at which the water level is rising when it reaches the shellfish,

(7)

(b) the earliest time after 10 00 hours on this day at which the water reaches the shellfish.

(4)

N 2 9 4 9 5 A 0 8 2 4

https://xtremepape.rs/

Leave blank

Question 4 continued

https://xtremepape.rs/

Leave blank

Question 4 continued

N 2 9 4 9 5 A 0 1 0 2 4

https://xtremepape.rs/

Leave blank

Question 4 continued

(Total 11 marks)

Q4

N 2 9 4 9 5 A 0 1 1 2 4

Turn over

https://xtremepape.rs/

Leave blank

5.



Figure 2

One end A of a light elastic string, of natural length a and modulus of elasticity $6mg$, is fixed at a point on a smooth plane inclined at 30° to the horizontal. A small ball B of mass m is attached to the other end of the string. Initially B is held at rest with the string lying along a line of greatest slope of the plane, with B below A and $AB = a$. The ball is released and comes to instantaneous rest at a point C on the plane, as shown in Figure 2. Find

(a) the length AC,

(5)

(b) the greatest speed attained by B as it moves from its initial position to C.

(7)

N 2 9 4 9 5 A 0 1 2 2 4

https://xtremepape.rs/

Leave blank

Question 5 continued

https://xtremepape.rs/

Leave blank

Question 5 continued

N 2 9 4 9 5 A 0 1 4 2 4

https://xtremepape.rs/

Leave blank

Question 5 continued

(Total 12 marks)

Q5

N 2 9 4 9 5 A 0 1 5 2 4

Turn over

https://xtremepape.rs/

Leave blank

6.



Figure 3

The region R is bounded by part of the curve with equation $y = 4 - x^2$, the positive x-axis and the positive y-axis, as shown in Figure 3. The unit of length on both axes is one metre. A uniform solid S is formed by rotating R through 360° about the x-axis.

(a) Show that the centre of mass of S is $\frac{5}{8}$ m from O.

(10)



Figure 4

Figure 4 shows a cross section of a uniform solid P consisting of two components, a solid cylinder C and the solid S. The cylinder C has radius 4 m and length l metres. One end of C coincides with the plane circular face of S. The point A is on the circumference of the circular face common to C and S. When the solid P is freely suspended from A, the solid P hangs with its axis of symmetry horizontal.

(b) Find the value of l.

(4)

N 2 9 4 9 5 A 0 1 6 2 4

https://xtremepape.rs/

Leave blank

Question 6 continued

https://xtremepape.rs/

Leave blank

Question 6 continued

N 2 9 4 9 5 A 0 1 8 2 4

https://xtremepape.rs/

Leave blank

Question 6 continued

(Total 14 marks)

Q6

N 2 9 4 9 5 A 0 1 9 2 4

Turn over

https://xtremepape.rs/

Leave blank

7.



Figure 5

A particle is projected from the highest point A on the outer surface of a fixed smooth sphere of radius a and centre O. The lowest point B of the sphere is fixed to a horizontal plane. The particle is projected horizontally from A with speed $\frac{1}{2}\sqrt{(ga)}$. The particle leaves the surface of the sphere at the point C, where $\angle AOC = \theta$, and strikes the plane at the point P, as shown in Figure 5.

(a) Show that $\cos\theta = \frac{3}{4}$.

(7)

(b) Find the angle that the velocity of the particle makes with the horizontal as it reaches P.

(8)

N 2 9 4 9 5 A 0 2 0 2 4

https://xtremepape.rs/

Leave blank

Question 7 continued

https://xtremepape.rs/

Leave blank

Question 7 continued

N 2 9 4 9 5 A 0 2 2 2 4

https://xtremepape.rs/

Leave blank

Question 7 continued

Q7

(Total 15 marks)

TOTAL FOR PAPER: 75 MARKS

END

N 2 9 4 9 5 A 0 2 3 2 4

https://xtremepape.rs/

BLANK PAGE

https://xtremepape.rs/