

1. A particle P of mass 0.5 kg is moving along the positive x -axis. At time t seconds, P is moving under the action of a single force of magnitude $[4 + \cos(\pi t)]$ N, directed away from the origin. When $t=1$, the particle P is moving away from the origin with speed 6 m s^{-1} .

Find the speed of P when $t=1.5$, giving your answer to 3 significant figures.

(7)



Question 1 continued

Handwriting practice area with horizontal lines.

Q1

(Total 7 marks)



2. A particle P moves in a straight line with simple harmonic motion of period 2.4 s about a fixed origin O . At time t seconds the speed of P is $v \text{ m s}^{-1}$. When $t=0$, P is at O . When $t=0.4$, $v=4$. Find

(a) the greatest speed of P ,

(7)

(b) the magnitude of the greatest acceleration of P .

(2)



3.




Figure 1

A bowl B consists of a uniform solid hemisphere, of radius r and centre O , from which is removed a solid hemisphere, of radius $\frac{2}{3}r$ and centre O , as shown in Figure 1.

- (a) Show that the distance of the centre of mass of B from O is $\frac{65}{152}r$. (5)




Figure 2

The bowl B has mass M . A particle of mass kM is attached to a point P on the outer rim of B . The system is placed with a point C on its outer curved surface in contact with a horizontal plane. The system is in equilibrium with P , O and C in the same vertical plane. The line OP makes an angle θ with the horizontal as shown in Figure 2. Given that

$$\tan \theta = \frac{4}{5},$$

- (b) find the exact value of k . (5)



Question 3 continued

Lined writing area for the response to Question 3.



Question 3 continued

Lined area for writing the answer to Question 3.

Q3

(Total 10 marks)

--	--



5.




Figure 4

One end A of a light inextensible string of length $3a$ is attached to a fixed point. A particle of mass m is attached to the other end B of the string. The particle is held in equilibrium at a distance $2a$ below the horizontal through A , with the string taut. The particle is then projected with speed $\sqrt{2ag}$, in the direction perpendicular to AB , in the vertical plane containing A and B , as shown in Figure 4. In the subsequent motion the string remains taut. When AB is at an angle θ below the horizontal, the speed of the particle is v and the tension in the string is T .

(a) Show that $v^2 = 2ag(3 \sin \theta - 1)$. (5)

(b) Find the range of values of T . (6)



Question 5 continued

Blank lined area for writing the answer to Question 5.



6. A bend of a race track is modelled as an arc of a horizontal circle of radius 120 m. The track is not banked at the bend. The maximum speed at which a motorcycle can be ridden round the bend without slipping sideways is 28 m s^{-1} . The motorcycle and its rider are modelled as a particle and air resistance is assumed to be negligible.

(a) Show that the coefficient of friction between the motorcycle and the track is $\frac{2}{3}$. **(6)**

The bend is now reconstructed so that the track is banked at an angle α to the horizontal. The maximum speed at which the motorcycle can now be ridden round the bend without slipping sideways is 35 m s^{-1} . The radius of the bend and the coefficient of friction between the motorcycle and the track are unchanged.

(b) Find the value of $\tan \alpha$. **(8)**



7. A light elastic string has natural length a and modulus of elasticity $\frac{3}{2}mg$. A particle P of mass m is attached to one end of the string. The other end of the string is attached to a fixed point A . The particle is released from rest at A and falls vertically. When P has fallen a distance $a + x$, where $x > 0$, the speed of P is v .

(a) Show that $v^2 = 2g(a + x) - \frac{3gx^2}{2a}$. (4)

(b) Find the greatest speed attained by P as it falls. (4)

After release, P next comes to instantaneous rest at a point D .

(c) Find the magnitude of the acceleration of P at D . (6)



Leave
blank

Question 7 continued

Handwriting lines for Question 7 continued.

(Total 14 marks)

Q7

--	--

TOTAL FOR PAPER: 75 MARKS

END

