

# Mark Scheme (Results) Summer 2010

GCE

## Core Mathematics C4 (6666)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our [Ask The Expert](#) email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

Summer 2010

Publications Code UA023705

All the material in this publication is copyright

© Edexcel Ltd 2010

June 2010  
6666 Core Mathematics C4  
Mark Scheme

Question Number	Scheme	Marks
<b>1.</b>	<p>(a) <math>y\left(\frac{\pi}{6}\right) \approx 1.2247, y\left(\frac{\pi}{4}\right) = 1.1180</math>      accept awrt 4 d.p.</p>	B1 B1 <b>(2)</b>
	<p>(b)(i) <math>I \approx \left(\frac{\pi}{12}\right)(1.3229 + 2 \times 1.2247 + 1)</math>      B1 for <math>\frac{\pi}{12}</math>  <math>\approx 1.249</math>      cao      A1</p>	B1 M1 A1
	<p>(ii) <math>I \approx \left(\frac{\pi}{24}\right)(1.3229 + 2 \times (1.2973 + 1.2247 + 1.1180) + 1)</math>      B1 for <math>\frac{\pi}{24}</math>  <math>\approx 1.257</math>      cao      A1</p>	B1 M1 A1 <b>(6)</b> <b>[8]</b>

Question Number	Scheme	Marks
2.	$\frac{du}{dx} = -\sin x$ $\int \sin x e^{\cos x+1} dx = -\int e^u du$ $= -e^u$ $= -e^{\cos x+1}$ $\left[-e^{\cos x+1}\right]_0^{\frac{\pi}{2}} = -e^1 - (-e^2)$ $= e(e-1) *$	<p>B1</p> <p>M1 A1</p> <p>A1ft</p> <p>ft sign error</p> <p>or equivalent with <math>u</math></p> <p>M1</p> <p>A1</p> <p>cso</p> <p><b>(6)</b></p> <p><b>[6]</b></p>

Question Number	Scheme	Marks
3.	$\frac{d}{dx}(2^x) = \ln 2 \cdot 2^x$ $\ln 2 \cdot 2^x + 2y \frac{dy}{dx} = 2y + 2x \frac{dy}{dx}$ <p>Substituting (3, 2)</p> $8 \ln 2 + 4 \frac{dy}{dx} = 4 + 6 \frac{dy}{dx}$ $\frac{dy}{dx} = 4 \ln 2 - 2$ <p style="text-align: right;">Accept exact equivalents</p>	<p>B1</p> <p>M1 A1= A1</p> <p>M1</p> <p>M1 A1 (7)</p> <p>[7]</p>

Question Number	Scheme	Marks
4.	<p>(a) <math>\frac{dx}{dt} = 2 \sin t \cos t, \frac{dy}{dt} = 2 \sec^2 t</math></p> <p><math>\frac{dy}{dx} = \frac{\sec^2 t}{\sin t \cos t} \left( = \frac{1}{\sin t \cos^3 t} \right)</math>      or equivalent</p> <p>(b) At <math>t = \frac{\pi}{3}, x = \frac{3}{4}, y = 2\sqrt{3}</math></p> <p><math>\frac{dy}{dx} = \frac{\sec^2 \frac{\pi}{3}}{\sin \frac{\pi}{3} \cos \frac{\pi}{3}} = \frac{16}{\sqrt{3}}</math></p> <p><math>y - 2\sqrt{3} = \frac{16}{\sqrt{3}} \left( x - \frac{3}{4} \right)</math></p> <p><math>y = 0 \Rightarrow x = \frac{3}{8}</math></p>	<p>B1 B1</p> <p>M1 A1      <b>(4)</b></p> <p>B1</p> <p>M1 A1</p> <p>M1</p> <p>M1 A1      <b>(6)</b></p> <p><b>[10]</b></p>

Question Number	Scheme	Marks
5.	<p>(a) <math>A = 2</math>  <math>2x^2 + 5x - 10 = A(x-1)(x+2) + B(x+2) + C(x-1)</math>  <math>x \rightarrow 1 \quad -3 = 3B \Rightarrow B = -1</math>  <math>x \rightarrow -2 \quad -12 = -3C \Rightarrow C = 4</math></p> <p>(b) <math>\frac{2x^2 + 5x - 10}{(x-1)(x+2)} = 2 + (1-x)^{-1} + 2\left(1 + \frac{x}{2}\right)^{-1}</math>  <math>(1-x)^{-1} = 1 + x + x^2 + \dots</math>  <math>\left(1 + \frac{x}{2}\right)^{-1} = 1 - \frac{x}{2} + \frac{x^2}{4} + \dots</math>  <math>\frac{2x^2 + 5x - 10}{(x-1)(x+2)} = (2+1+2) + (1-1)x + \left(1 + \frac{1}{2}\right)x^2 + \dots</math>  <math>= 5 + \dots</math> ft their <math>A - B + \frac{1}{2}C</math>  <math>= \dots + \frac{3}{2}x^2 + \dots</math> <math>0x</math> stated or implied</p>	<p>B1</p> <p>M1 A1</p> <p>A1 (4)</p> <p>M1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1 ft</p> <p>A1 A1 (7)</p> <p>[11]</p>

Question Number	Scheme	Marks
6.	<p>(a)</p> $f(\theta) = 4\cos^2\theta - 3\sin^2\theta$ $= 4\left(\frac{1}{2} + \frac{1}{2}\cos 2\theta\right) - 3\left(\frac{1}{2} - \frac{1}{2}\cos 2\theta\right)$ $= \frac{1}{2} + \frac{7}{2}\cos 2\theta \quad *$ <p>(b)</p> $\int \theta \cos 2\theta \, d\theta = \frac{1}{2}\theta \sin 2\theta - \frac{1}{2} \int \sin 2\theta \, d\theta$ $= \frac{1}{2}\theta \sin 2\theta + \frac{1}{4}\cos 2\theta$ $\int \theta f(\theta) \, d\theta = \frac{1}{4}\theta^2 + \frac{7}{4}\theta \sin 2\theta + \frac{7}{8}\cos 2\theta$ $\left[ \dots \right]_0^{\frac{\pi}{2}} = \left[ \frac{\pi^2}{16} + 0 - \frac{7}{8} \right] - \left[ 0 + 0 + \frac{7}{8} \right]$ $= \frac{\pi^2}{16} - \frac{7}{4}$	<p>M1 M1</p> <p>cso A1 (3)</p> <p>M1 A1</p> <p>A1</p> <p>M1 A1</p> <p>M1</p> <p>A1 (7)</p> <p>[10]</p>



Question Number	Scheme	Marks
7.	<p>(a) <b>j</b> components <math>3 + 2\lambda = 9 \Rightarrow \lambda = 3</math> <span style="float: right;"><math>(\mu = 1)</math></span>  Leading to <math>C : (5, 9, -1)</math> <span style="float: right;">accept vector forms</span></p> <p>(b) <span style="float: right;">Choosing correct directions or finding <math>\overline{AC}</math> and <math>\overline{BC}</math></span>  <math display="block">\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} \cdot \begin{pmatrix} 5 \\ 0 \\ 2 \end{pmatrix} = 5 + 2 = \sqrt{6}\sqrt{29} \cos \angle ACB</math> <span style="float: right;">use of scalar product</span>  <math>\angle ACB = 57.95^\circ</math> <span style="float: right;">awrt <math>57.95^\circ</math></span></p> <p>(c) <math>A : (2, 3, -4)</math> <math>B : (-5, 9, -5)</math>  <math display="block">\overline{AC} = \begin{pmatrix} 3 \\ 6 \\ 3 \end{pmatrix}, \quad \overline{BC} = \begin{pmatrix} 10 \\ 0 \\ 4 \end{pmatrix}</math> <math display="block">AC^2 = 3^2 + 6^2 + 3^2 \Rightarrow AC = 3\sqrt{6}</math> <math display="block">BC^2 = 10^2 + 4^2 \Rightarrow BC = 2\sqrt{29}</math> <math display="block">\Delta ABC = \frac{1}{2} AC \times BC \sin \angle ACB</math> <math display="block">= \frac{1}{2} 3\sqrt{6} \times 2\sqrt{29} \sin \angle ACB \approx 33.5</math> <span style="float: right;"><math>15\sqrt{5}</math>, awrt 34</span></p>	<p>M1 A1 A1 <b>(3)</b></p> <p>M1 M1 A1 A1 <b>(4)</b></p> <p>M1 A1 A1 M1 A1 <b>(5)</b> <b>[12]</b></p>
	<p><i>Alternative method for (b) and (c)</i></p> <p>(b) <math>A : (2, 3, -4)</math> <math>B : (-5, 9, -5)</math> <math>C : (5, 9, -1)</math>  <math>AB^2 = 7^2 + 6^2 + 1^2 = 86</math>  <math>AC^2 = 3^2 + 6^2 + 3^2 = 54</math>  <math>BC^2 = 10^2 + 0^2 + 4^2 = 116</math> <span style="float: right;">Finding all three sides</span>  <math>\cos \angle ACB = \frac{116 + 54 - 86}{2\sqrt{116}\sqrt{54}} (= 0.53066 \dots)</math>  <math>\angle ACB = 57.95^\circ</math> <span style="float: right;">awrt <math>57.95^\circ</math></span></p> <p>If this method is used some of the working may gain credit in part (c) and appropriate marks may be awarded if there is an attempt at part (c).</p>	<p>M1 M1 A1 A1 <b>(4)</b></p>

Question Number	Scheme	Marks
8.	<p>(a)</p> $\frac{dV}{dt} = 0.48\pi - 0.6\pi h$ $V = 9\pi h \Rightarrow \frac{dV}{dt} = 9\pi \frac{dh}{dt}$ $9\pi \frac{dh}{dt} = 0.48\pi - 0.6\pi h$ <p>Leading to <math>75 \frac{dh}{dt} = 4 - 5h</math> *</p> <p>(b)</p> $\int \frac{75}{4-5h} dh = \int 1 dt$ <p style="text-align: right;">separating variables</p> $-15 \ln(4-5h) = t (+C)$ $-15 \ln(4-5h) = t + C$ <p>When <math>t = 0, h = 0.2</math></p> $-15 \ln 3 = C$ $t = 15 \ln 3 - 15 \ln(4-5h)$ <p>When <math>h = 0.5</math></p> $t = 15 \ln 3 - 15 \ln 1.5 = 15 \ln \left( \frac{3}{1.5} \right) = 15 \ln 2$ <p style="text-align: right;">awrt 10.4</p> <p><i>Alternative for last 3 marks</i></p> $t = \left[ -15 \ln(4-5h) \right]_{0.2}^{0.5}$ $= -15 \ln 1.5 + 15 \ln 3$ $= 15 \ln \left( \frac{3}{1.5} \right) = 15 \ln 2$ <p style="text-align: right;">awrt 10.4</p>	<p>M1 A1</p> <p>B1</p> <p>M1</p> <p>cs0 A1 (5)</p> <p>M1</p> <p>M1 A1</p> <p>M1</p> <p>M1 A1</p> <p>M1 M1</p> <p>A1 (6)</p>



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@inneydirect.com](mailto:publications@inneydirect.com)

Order Code UA023705 Summer 2010

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no. 4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH