

# Mark Scheme (Results) Summer 2010

GCE

## GCE Mechanics M3 (6679/01)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>


Summer 2010

Publications Code UA024475


All the material in this publication is copyright


© Edexcel Ltd 2010


**Summer 2010  
Mechanics M3 6679  
Mark Scheme**

Question Number	Scheme	Marks
<p>Q1</p> <p>(a)</p>	<div style="text-align: center;">  </div> $\cos \alpha = \frac{12}{13}$ $R(\uparrow) \quad T \cos \alpha = mg$ $T \times \frac{12}{13} = mg$ $T = \frac{13}{12} mg \quad \text{oe}$	<p>B1</p> <p>M1</p> <p>A1 (3)</p>
<p>(b)</p>	<p>Eqn of motion <math>T \sin \alpha = m \frac{v^2}{5l}</math></p> $\frac{13mg}{12} \times \frac{5}{13} = m \frac{v^2}{5l}$ $v^2 = \frac{25gl}{12}$ $v = \frac{5}{2} \sqrt{\frac{gl}{3}} \quad \left( \text{accept } 5\sqrt{\frac{gl}{12}} \text{ or } \sqrt{\frac{25gl}{12}} \text{ or any other equiv} \right)$	<p>M1 A1</p> <p>M1 dep</p> <p>A1 (4)</p> <p>[7]</p>

Question Number	Scheme	Marks
Q2 (a)	$F = (-) \frac{k}{x^2}$ $mg = (-) \frac{k}{R^2}$ $F = \frac{mgR^2}{x^2} *$	M1 M1 A1 (3)
(b)	$m\ddot{x} = -\frac{mgR^2}{x^2}$ $v \frac{dv}{dx} = -\frac{gR^2}{x^2}$ $\frac{1}{2}v^2 = \int \left( -\frac{gR^2}{x^2} \right) dx$ $\frac{1}{2}v^2 = \frac{gR^2}{x} (+c)$ $x = R, v = 3U \quad \frac{9U^2}{2} = gR + c$ $\frac{1}{2}v^2 = \frac{gR^2}{x} + \frac{9U^2}{2} - gR$ $x = 2R, v = U \quad \frac{1}{2}U^2 = \frac{gR^2}{2R} + \frac{9U^2}{2} - gR$ $U^2 = \frac{gR}{8}$ $U = \sqrt{\frac{gR}{8}}$	M1 M1 M1 dep on 1st M mark A1 M1 dep on 3rd M mark M1 dep on 3rd M mark A1 (7) <b>[10]</b>



Question Number	Scheme	Marks
Q3	 <p> <math display="block">\text{EPE lost} = \frac{\lambda \times 0.6^2}{2 \times 0.9} - \frac{\lambda \times 0.1^2}{2 \times 0.9} \left( = \frac{7}{36} \lambda \right)</math> <math display="block">R(\uparrow) \quad R = mg \cos \theta</math> <math display="block">= 0.5g \times \frac{4}{5} = 0.4g</math> <math display="block">F = \mu R = 0.15 \times 0.4g</math> <p>P.E. gained = E.P.E. lost – work done against friction</p> <math display="block">0.5g \times 0.7 \sin \theta = \frac{\lambda \times 0.6^2}{2 \times 0.9} - \frac{\lambda \times 0.1^2}{2 \times 0.9} - 0.15 \times 0.4g \times 0.7</math> <math display="block">0.1944\lambda = 0.5 \times 9.8 \times 0.7 \times \frac{3}{5} + 0.15 \times 0.4 \times 9.8 \times 0.7</math> <math display="block">\lambda = 12.70 \dots</math> <math display="block">\lambda = 13 \text{ N} \quad \text{or } 12.7</math> </p>	<p>M1 A1</p> <p>M1</p> <p>M1 A1</p> <p>M1 A1 A1</p> <p>A1</p> <p style="text-align: right;"><b>[9]</b></p>

Question Number	Scheme			Marks	
Q4 (a)		cone	container	cylinder	M1 A1  B1  M1 A1ft A1 (6)
	mass ratio	$\frac{4\pi l^3}{3}$	$\frac{68\pi l^3}{3}$	$24\pi l^3$	
		4	68	72	
	dist from $O$	$l$	$\bar{x}$	$3l$	
Moments:		$4l + 68\bar{x} = 72 \times 3l$ $\bar{x} = \frac{212l}{68} = \frac{53}{17}l$ accept $3.12l$			
(b)	 <p data-bbox="411 1276 667 1312"><math>GX = 6l - \bar{x}</math> seen</p> $\tan \theta = \frac{2l}{6l - \bar{x}}$ $= \frac{2 \times 17}{49}$ $\theta = 34.75\dots = 34.8 \text{ or } 35$			M1 M1 A1  A1 (4) [10]	

Question Number	Scheme	Marks
Q5		
(a)	Energy: $mga \sin \theta = \frac{1}{2}m \times 5ag - \frac{1}{2}mv^2$ $v^2 = 5ag - 2ag \sin \theta$	M1 A1 A1 (3)
(b)	Eqn of motion along radius: $T + mg \sin \theta = \frac{mv^2}{a}$ $T = \frac{m}{a}(5ag - 2ag \sin \theta) - mg \sin \theta$ $T = mg(5 - 3 \sin \theta)$	M1 A1 M1 A1 (4)
(c)	At C, $\theta = 90^\circ$ $T = mg(5 - 3) = 2mg$ $T > 0 \therefore P$ reaches C	M1 A1 A1 (3)
(d)	Max speed at lowest point ( $\theta = 270^\circ$ ; $v^2 = 5ag - 2ag \sin 270$ $v^2 = 5ag + 2ag$ $v = \sqrt{7ag}$ )	M1 A1 (2) [12]

Question Number	Scheme	Marks
Q6 (a)	$\frac{d^2x}{dt^2} = -\frac{3}{(t+1)^2}$ $\frac{dx}{dt} = \int -3(t+1)^{-2} dt$ $= 3(t+1)^{-1} (+c)$ <p><math>t = 0, v = 2 \quad 2 = 3 + c \quad c = -1</math></p> $\frac{dx}{dt} = \frac{3}{t+1} - 1 \quad *$	<p>M1</p> <p>M1 A1</p> <p>M1</p> <p>A1 (5)</p>
(b)	$x = \int \left( \frac{3}{t+1} - 1 \right) dt$ $= 3 \ln(t+1) - t \quad (+c')$ <p><math>t = 0, x = 0 \Rightarrow c' = 0</math></p> $x = 3 \ln(t+1) - t$ $v = 0 \Rightarrow \frac{3}{t+1} = 1$ $t = 2$ $x = 3 \ln 3 - 2$ $= 1.295\dots$ $= 1.30 \text{ m (Allow 1.3)}$	<p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1 (7)</p> <p>[12]</p>



Question Number	Scheme	Marks
<p>Q7</p> <p>(a)</p>	 <p>R(↑) <math>T = 2mg</math></p> <p>Hooke's law: <math>T = \frac{6mge}{3a}</math></p> <p><math>2mg = \frac{6mge}{3a}</math></p> <p><math>e = a</math></p> <p><math>AO = 4a</math></p>	<p>B1</p> <p>M!</p> <p>A1 (3)</p>
<p>(b)</p>	 <p>H.L.</p> <p>Eqn. of motion</p> <p><math>T = \frac{6mg(a-x)}{3a} = \frac{2mg(a-x)}{a}</math></p> <p><math>-2mg + T = 2m\ddot{x}</math></p> <p><math>-2mg + \frac{2mg(a-x)}{a} = 2m\ddot{x}</math></p> <p><math>-\frac{2mgx}{a} = 2m\ddot{x}</math></p> <p><math>\ddot{x} = -\frac{g}{a}x</math></p> <p>period <math>2\pi\sqrt{\frac{a}{g}}</math> *</p>	<p>B1ft</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>A1 (5)</p>

Question Number	Scheme	Marks
(c)	$v^2 = \omega^2 (a^2 - x^2)$ $v_{\max}^2 = \frac{g}{a} \left( \left( \frac{a}{4} \right)^2 - 0 \right)$ $v_{\max} = \frac{1}{4} \sqrt{(ga)}$	<p>M1 A1</p> <p>A1 (3)</p>
(d)	$x = -\frac{a}{8}$ $v^2 = \frac{g}{a} \left( \frac{a^2}{16} - \frac{a^2}{64} \right)$ $= \frac{3ag}{64}$ $v^2 = u^2 + 2as$ $0 = \frac{3ag}{64} - 2gh$ $h = \frac{3a}{128}$ <p>Total height above <math>O = \frac{a}{8} + \frac{3a}{128} = \frac{19a}{128}</math></p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>A1 (4)</p> <p>[15]</p>



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA024475 Summer 2010

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH