Centre No.					Pape	r Refer	ence			Surname	Initial(s)
Candidate No.			6	6	6	7	/	0	1	Signature	

6667/01

Edexcel GCE

Further Pure Mathematics FP1 Advanced/Advanced Subsidiary

Tuesday 22 June 2010 – Afternoon

Time: 1 hour 30 minutes

Materials required for examination
Mathematical Formulae (Pink)

Items included with question papers

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 9 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy. ©2010 Edexcel Limited.

Printer's Log. No.
N35387A W850/R6667/57570 4/3





1

2

3

4

5

6

7

8

1.	z = 2 - 3i	
	(a) Show that $z^2 = -5 - 12i$.	
		(2)
	Find, showing your working,	
	(b) the value of $ z^2 $,	
		(2)
	(c) the value of $arg(z^2)$, giving your answer in radians to 2 decimal places.	(2)
	(d) Show z and z^2 on a single Argand diagram.	(1)

Leave

Question 1 continued		Le
		01
		Q1
	(Total 7 marks)	

N 3 5 3 8 7 A 0 3 2 8

Turn over

$\mathbf{M} = \begin{pmatrix} 2a & 3 \\ 6 & a \end{pmatrix}, \text{ where } a \text{ is a real constant.}$	Leave blank
--	----------------

2.

(a) Given that a = 2, find \mathbf{M}^{-1} .

		(3)
(h)	Find the values of a for which M is singular	

a for which M is singular.	(2

Question 2 continued		Le bla
		Q2
	(Total 5 marks)	

N 3 5 3 8 7 A 0 5 2 8

Turn over

2	
Э	

$$f(x) = x^3 - \frac{7}{x} + 2, \quad x > 0$$

Leave blank

(a) Show that f(x) = 0 has a root α between 1.4 and 1.5

(2)

(b) Starting with the interval [1.4, 1.5], use interval bisection twice to find an interval of width 0.025 that contains α .

(3)

(c) Taking 1.45 as a first approximation to α , apply the Newton-Raphson procedure once to $f(x) = x^3 - \frac{7}{x} + 2$ to obtain a second approximation to α , giving your answer to 3 decimal places.

(5)



Question 3 continued	Leave blank



Question 3 continued	Lea bla
Question 5 continued	



Question 3 continued	Ldbl
	(Total 10 marks)

N 3 5 3 8 7 A 0 9 2 8

9

		Leave blank
4.	$f(x) = x^3 + x^2 + 44x + 150$	
	Given that $f(x) = (x+3)(x^2 + ax + b)$, where a and b are real constants,	

(a)	find the value of a and the value of b .	(2)
(b)	Find the three roots of $f(x) = 0$.	(4)

(c) Find the sum of the three roots of $f(x) = 0$.	



Question 4 continued		Leave blank
	(Total 7 marks)	Q4

The parabola C has equation $y^2 = 20x$.	
(a) Verify that the point $P(5t^2, 10t)$ is a general point on C .	(1)
	(1)
The point A on C has parameter $t = 4$.	
The line l passes through A and also passes through the focus of C .	
(b) Find the gradient of <i>l</i> .	(4)
	(4)



Question 5 continued	Lobl
	Q5

13

Leave
blank

- **6.** Write down the 2×2 matrix that represents
 - (a) an enlargement with centre (0,0) and scale factor 8,

(1)

(b) a reflection in the *x*-axis.

(1)

Hence, or otherwise,

(c) find the matrix T that represents an enlargement with centre (0,0) and scale factor 8, followed by a reflection in the x-axis.

(2)

$$\mathbf{A} = \begin{pmatrix} 6 & 1 \\ 4 & 2 \end{pmatrix}$$
 and $\mathbf{B} = \begin{pmatrix} k & 1 \\ c & -6 \end{pmatrix}$, where k and c are constants.

(d) Find **AB**.

(3)

Given that AB represents the same transformation as T,

(e) find the value of k and the value of c.

(2)

	Leave
	blank
Question 6 continued	
	1



	Leave blank
Question 6 continued	



Question 6 continued		Lea bla
		Q6
	(Total 9 marks)	

Leave
blank

•	$f(n) = 2^n + 6^n$	
	(a) Show that $f(k+1) = 6f(k) - 4(2^k)$.	(3)
	(b) Hence, or otherwise, prove by induction that, for $n \in \mathbb{Z}^+$, $f(n)$ is divisible by 8.	(4)
_		

Question 7 continued		Le bla
	_	
		07
		Q 7

19

The rectangular hyperbola H has equation $xy = c^2$, where c is a positive constant.	
The point A on H has x -coordinate $3c$.	
(a) Write down the y-coordinate of A.	
	(1)
(b) Show that an equation of the normal to H at A is	
3y = 27x - 80c	(5)
The manual to Hat America Harring at the maint D	(3)
The normal to H at A meets H again at the point B .	
(c) Find, in terms of c , the coordinates of B .	(5)

	Leave
	blank
Question 8 continued	014111
Question o continued	
	1



Question 8 continued	I t
	_



Question 8 continued	Le bla

N 3 5 3 8 7 A 0 2 3 2 8

23

Leave blank

9. (a) Prove by induction that

$$\sum_{r=1}^{n} r^2 = \frac{1}{6} n(n+1)(2n+1)$$

(6)

Using the standard results for $\sum_{r=1}^{n} r$ and $\sum_{r=1}^{n} r^2$,

(b) show that

$$\sum_{r=1}^{n} (r+2)(r+3) = \frac{1}{3}n(n^2 + an + b),$$

where a and b are integers to be found.

(5)

(c) Hence show that

$$\sum_{r=n+1}^{2n} (r+2)(r+3) = \frac{1}{3}n(7n^2 + 27n + 26)$$

(3)

Question 9 continued	Leave blank



Question 9 continued	Leave blank



	Leave
Question 9 continued	blank
Question / continued	



Question 9 continued		Lea
	(Total 14 marks)	
	TOTAL FOR PAPER: 75 MARKS	
EN		

